

28-OCT-1998

(Jug Sleeve)

ASSURE(R) HERBICIDE

EMULSIFIABLE CONCENTRATE

GUARANTEE: Quizalofop-Ethyl.....96 g/L

FOR WEED CONTROL IN CANOLA, FLAX (INCLUDING LOW LINOLENIC ACID VARIETIES), SOYBEANS, SEED ALFALFA, LENTILS, PEAS (FIELD AND PROCESSING), AND SEEDLING LEGUMES FOR SEED PRODUCTION

AGRICULTURAL

READ THE LABEL AND ATTACHED BROCHURE BEFORE USING

REGISTRATION NO. 22111

PEST CONTROL PRODUCTS ACT

NET CONTENTS: 8 LITRES

THIS PRODUCT CONTAINS 4% OF PHENOL WHICH HAS BEEN DETERMINED TO BE OF TOXICOLOGICAL CONCERN.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN

CAUTION! MAY IRRITATE EYES, NOSE, THROAT, AND SKIN.

WARNING - EYE AND SKIN IRRITANT

May be harmful if swallowed, inhaled or absorbed through skin. Cause eye irritation.

Do not get in eyes or on clothing. Avoid contact with skin. Avoid breathing vapours or spray mist. Use adequate ventilation.

Wear goggles or face shield, chemical resistant gloves and coveralls during mixing, loading, clean up and repair.

For worker protection during application, wear a hat and coveralls.

When using, do not eat, drink or smoke. Wash skin thoroughly with soap and water after handling.

Remove contaminated clothing immediately after use. Store and wash contaminated clothing separately from household laundry before reuse

DU PONT CANADA INC.
AGRICULTURAL PRODUCTS
BOX 2300 STREETSVILLE
MISSISSAUGA, ONTARIO
L5M 2J4
1-800-667-3925

FIRST AID

IF SWALLOWED: If swallowed, DO NOT induce vomiting. Immediately give two glasses of water. Never give anything by mouth to an unconscious person. Call a physician or contact a poison control centre.

IF ON SKIN: In case of contact, immediately flush skin with plenty of water for at least 15 minutes while removing contaminated clothing and shoes. Call a physician or contact a poison control centre. Wash contaminated clothing before reuse.

IF IN EYES: In case of contact, immediately flush eyes with plenty of water for at least 15 minutes. Call a physician or contact a poison control centre.

IF INHALED: If inhaled, remove to fresh air. If not breathing, give artificial respiration. If breathing is difficult, give oxygen. Call a physician or contact a poison control centre.

Never give anything by mouth to an unconscious person. Remove victim to safe, uncontaminated area. If victim has ceased breathing clear airway and start artificial respiration. Obtain medical attention or contact a poison control centre. Take container or label or the product name and product registration number with you to emergency/hospital or physician.

TOXICOLOGICAL INFORMATION:

"THIS PRODUCT CONTAINS PETROLEUM DISTILLATES"

For medical emergencies and transportation emergencies such as spills, leaks or fires, call 1-613-348-3616 (24 hours).

STORAGE AND DISPOSAL

STORAGE:

Store product in original container only, away from fertilizer, seed, food or feed. Not for use or storage in or around the home. Keep container closed. DO NOT FREEZE.

DISPOSAL:

1. Rinse the emptied container thoroughly and add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the Provincial Regulatory Agency or the Manufacturer.

NOTICE TO USER

This control product is to be used only in accordance with the directions on this label. It is an offence under the PEST CONTROL PRODUCTS ACT to use a control product under unsafe conditions.

NOTICE TO BUYER

Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

The seller warrants that the purchase by the buyer and the use of this product, as such, will not infringe any Canadian patent.

(Directions for Use Folder))

ASSURE(R)
HERBICIDE

EMULSIFIABLE CONCENTRATE

GUARANTEE: Quizalofop-Ethyl.....96 g/L

FOR WEED CONTROL IN CANOLA, FLAX (INCLUDING LOW LINOLENIC ACID VARIETIES), SOYBEANS, SEED ALFALFA, LENTILS, PEAS (FIELD AND PROCESSING), AND SEEDLING LEGUMES FOR SEED PRODUCTION

AGRICULTURAL

READ THE LABEL AND THIS BROCHURE BEFORE USING

REGISTRATION NO. 22111

PEST CONTROL PRODUCTS ACT

This product contains 4% of phenol which has been determined to be of toxicological concern.

PRECAUTIONS:

CAUTION! KEEP OUT OF REACH OF CHILDREN
MAY IRRITATE EYES, NOSE, THROAT, AND SKIN.

WARNING - EYE AND SKIN IRRITANT

May be harmful if swallowed, inhaled or absorbed through skin. Causes eye irritation.

Do not get in eyes or on clothing. Avoid contact with skin. Avoid breathing vapours or spray mist. Use adequate ventilation.

Wear goggles or face shield, chemical resistant gloves and coveralls during mixing, loading, clean up and repair.

For worker protection during application, wear a hat and coveralls.

When using, do not eat, drink or smoke. Wash skin thoroughly with soap and water after handling.

Remove contaminated clothing immediately after use. Store and wash contaminated clothing separately from household laundry before reuse.

DU PONT CANADA INC.
AGRICULTURAL PRODUCTS
BOX 2300 STREETSVILLE

MISSISSAUGA, ONTARIO
L5M 2J4
1-800-667-3925

FIRST AID

IF SWALLOWED: If swallowed, DO NOT induce vomiting. Immediately give two glasses of water. Never give anything by mouth to an unconscious person. Call a physician or contact a poison control centre.

IF ON SKIN: In case of contact, immediately flush skin with plenty of water for at least 15 minutes while removing contaminated clothing and shoes. Call a physician or contact a poison control centre. Wash contaminated clothing before reuse.

IF IN EYES: In case of contact, immediately flush eyes with plenty of water for at least 15 minutes. Call a physician or contact a poison control centre.

IF INHALED: If inhaled, remove to fresh air. If not breathing, give artificial respiration. If breathing is difficult, give oxygen. Call a physician or contact a poison control centre.

Never give anything by mouth to an unconscious person. Remove victim to safe, uncontaminated area. If victim has ceased breathing clear airway and start artificial respiration. Obtain medical attention or contact a poison control centre. Take container or label or the product name and product registration number with you to emergency/hospital or physician.

TOXICOLOGICAL INFORMATION:

"THIS PRODUCT CONTAINS PETROLEUM DISTILLATES"

For medical emergencies and transportation emergencies such as spills, leaks or fires, call 1-613-348-3616.

IMPORTANT

Injury to or loss of desirable trees or vegetation may result from failure to observe the following: Do not apply or drain or flush equipment on or near desirable trees or plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not use on lawns, walks, driveways, tennis courts, or similar areas. Prevent drift of spray to desirable plants. Do not contaminate any body of water including irrigation water that may be used on other crops.

DO NOT APPLY BY AIR. APPLY WITH GROUND EQUIPMENT ONLY.

This product is toxic to fish. Overspray or drift to fish-bearing water must be avoided. Overspray or drift to important wildlife habitats such as shelterbelts, wetlands, sloughs or dry slough borders and woodlots should be avoided. Leave a 15 metre buffer zone between the last spray swath and the edge of any of these habitats.

Use caution when handling ASSURE® Herbicide as accidental spills or splashes may harm the finish on painted equipment.

DO NOT SUBJECT TO TEMPERATURES BELOW 0°C.

FOR USE ONLY AS AN AGRICULTURAL HERBICIDE.

GENERAL INFORMATION

ASSURE® is a selective postemergence herbicide for the control of annual and perennial grasses in flax (including low linolenic acid varieties), canola, soybeans, seed alfalfa, lentils, peas (field and processing) and seedling legumes for seed production. Recommended rates and timing for ASSURE® Herbicide application will provide control of the emerged grasses present.

ASSURE® Herbicide does not control sedges or broadleaf weeds.

Grass crops, including wheat, barley, rye, oats, and corn are highly sensitive to ASSURE® Herbicide.

When not in use, keep container closed to prevent evaporation or loss through spillage.

Under certain environmental conditions, minor leaf spotting may occur on soybeans. New growth is unaffected and there is no adverse effect on crop yield.

BIOLOGICAL ACTIVITY

ASSURE® is a systemic ACCase herbicide which is rapidly absorbed and readily translocated from the treated foliage to the root systems and growing points of the plant.

Control Symptoms - Treated plants show a reduction in growth and a loss of competitiveness. An early yellowing or browning of the younger plant tissues is followed by a progressive collapse of the remaining foliage. These symptoms will generally be observed in one to three weeks depending on the grass species treated and the environmental conditions.

DIRECTIONS FOR USE

Weeds that emerge after application may not be controlled.

POSTEMERGENCE APPLICATIONS - CANOLA, FLAX (INCLUDING LOW LINOLENIC ACID VARIETIES), SOYBEANS, SEED ALFALFA, LENTILS AND PEAS (FIELD AND PROCESSING):

Apply the specified rate of ASSURE® with CANPLUSCR® 411 or HI-MIX™ at 5-1 litres per 1000 litres of spray solution at the leaf stage of the grass indicated below:

USE RATES

	Weeds Controlled	Weed Leaf Stage at Application
0.75 L/ha	Green foxtail	2 leaf to early tillering
	Volunteer barley	2 leaf to early tillering
	Volunteer Corn	2-6 leaf
	Volunteer oats	*2 leaf to early tillering
	Volunteer wheat	2 leaf to early tillering
	Wild Oats	1-5 leaf without tillers
1.0 L/ha	Those controlled by 0.75 L/ha and Barnyard grass	*2 leaf to early tillering
	Wild Oats	*2 leaf to early tillering
1.0 L/ha	Quackgrass	2-6 leaf
1.5 L/ha	Those listed above and Quackgrass	2-6 leaf

*Best results on volunteer/wild oats if application is made before tillering begins.

Use the higher rate of CANPLUS® 411 or HI-MIX™, when wild oats or quackgrass are the main target weeds, or under conditions not conducive good growth.

When using a broadleaf herbicide, other than those registered for tank mixing with ASSURE®, wait a minimum of 24 hours after the ASSURE® application to apply the broadleaf herbicide or wait 7 days after the broadleaf application to apply ASSURE®.

TANK MIXTURES WITH BROADLEAF HERBICIDES

CANOLA - For control of broadleaf weeds and grasses in canola (rapeseed), apply a tank mixture of MUSTER(R) Herbicide plus ASSURE®. Use MUSTER® at 20 or 30 grams per hectare plus 1.0 litre per hectare ASSURE® plus CANPLUS® 411

or HI-MIX™ at 5-10 litres per 1000 litres of spray solution. Refer to the "Mixing Instructions" section of this label for specific mixing instructions.

Additional surfactant is not required when using CANPLUS® 411 or HI-MIX™. For best results follow the timing restrictions and specific weed control recommendations stated on each product's label. After application follow the sprayer clean-out directions on the MUSTER® label. Consult the MUSTER® label for restrictions and desired use rates and weeds controlled and additional restrictions regarding the use of MUSTER®.

NOTE TO BUYER/USER: READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR SPECIAL USE APPLICATIONS:

The DIRECTIONS FOR USE for this product for the use described below were developed by persons other than DuPont Canada Inc. and accepted for registration by Health Canada under the User Requested Minor Use Label Expansion Program. DuPont Canada Inc. itself makes no representation or warranty with respect to product performance (efficacy) and crop tolerance (phytotoxicity) of this product when used on the crop listed. Accordingly, the Buyer and User assume all liability arising, and agree to hold DuPont Canada Inc. harmless from any claims based on efficacy or phytotoxicity in connection with the use described below.

SEEDLING LEGUMES FOR SEED PRODUCTION ONLY:

For bird's-foot trefoil, alsike, red, white and sweet clover, and sainfoin make a postemergent broadcast application of:

ASSURE® Herbicide	1.0-1.5 litres/hectare
plus	
CANPLUS®	5-10 litres per 1000 litres spray solution
or	
HI-MIX	5-10 litres per 1000 litres spray solution

DO NOT graze or cut for hay in the year of treatment.

Refer to the weed control chart printed in the Directions for Use section of this label for application rates to control quackgrass and annual grass species. Application to the above species should be made before crop canopy closes.

Make only one application per growing season. Apply the spray mixture with ground equipment only. Use a calibrated sprayer that will provide thorough coverage of the weeds with the spray solution. Apply a minimum of 100 litres/ha of total spray volume.

ESTABLISHED CREEPING RED FESCUE FOR SEED:

For postemergent broadleaf and grassy weed control:

ASSURE® Herbicide + ALLY® Herbicide + Adjuvant

ASSURE® Herbicide	1.0-1.5 litres/hectare
plus	
ALLY® Herbicide	7.5 grams/hectare
plus	
SURE-MIX®	5 litres per 1000 litres spray solution
or	
CANPLUS® 411	5-10 litres per 1000 litres spray solution

This tankmix will control all label weeds listed on this label and the ALLY® Herbicide label.

Apply ASSURE® Herbicide + ALLY® Herbicide tank mixture postemergent to the first main flush of actively growing weeds, when established creeping red fescue is in the 2 leaf to flag leaf (shot blade) stage. Apply before the crop canopy is dense enough to prevent thorough coverage of the weeds.

Apply with ground equipment only. Make only one application per growing season. Refer to other sections of this label, and to ALLY® Herbicide label for additional application and/or use precaution instructions.

ESTABLISHED CREEPING RED FESCUE FOR SEED PRODUCTION ONLY:

For control of grassy weeds, make a postemergent broadcast application of

ASSURE® Herbicide	1.0-1.5 litres/hectare
plus	
CANPLUS® 411	5-10 litres per 1000 litres spray solution
or	
HI-MIX®	5-10 litres per 1000 litres spray solution

Do not graze or cut treated established red fescue for feed in the year of treatment.

Apply when quackgrass is in the 2-6 leaf stage and when annual grassy weeds are from the 2 leaf stage to early tillering. Application to the above species should be made before crop canopy closes. Refer to other sections of this label for a list of grassy weeds controlled as well as the

recommended ASSURE® Herbicide rates depending on the leaf stage of quackgrass or the annual grass species that are present.

Make only one application per growing season. Apply the spray mixture with ground equipment only. Use a properly calibrated sprayer that will provide thorough coverage of the weeds with the spray solution. Apply a minimum of 100 litres/ha of total spray volume.

FOR USE ON SUGARBEETS IN CANADA:

Apply ASSURE® as a postemergent broadcast application to small, emerged grassy weeds between the 2 leaf to early tillering leaf stage. Application to grassy weeds should be made before the crop canopy closes to maximize spray coverage.

To control annual grasses and volunteer cereals; apply:

ASSURE®	0.75 L/ha
plus	
SURE-MIX™	5 Litres per 1000 Litres spray solution
or	
CANPLUS®	5-10 Litres/1000 Litres spray solution

If a second flush of annual grasses or volunteer cereals is observed, apply an additional (sequential) application of ASSURE® at 0.75 L/ha + recommended adjuvant.

To control annual grasses, volunteer cereals and quackgrass, apply ASSURE® at 1.5 L/ha + recommended adjuvant.

Do not exceed an accumulative, seasonal use rate of 1.5 L/ha ASSURE® per year.

Apply the spray mixture with ground equipment only. Use a properly calibrated sprayer that will provide thorough coverage of the weeds with the spray solution. Do not apply to sugarbeets within 60-80 days of harvest.

For a complete list of grassy weeds, as well as additional application and/or use precaution instructions, please refer to other sections of this label.

SOYBEANS

ASSURE® + PINNACLE® + BASAGRAN FORTE®

For the control of annual grassy and broadleaf weeds, apply a tank mixture

of ASSURE® plus PINNACLE® plus BASAGRAN FORTE®. Use ASSURE® at 1.25 litres per hectare plus PINNACLE® at 5.5-8.0 grams per hectare plus BASAGRAN FORTE® at 1.75-2.25 litres per hectare. Add CANPLUS® 411 or HI-MIX™ at 5-10 litres per 1000 litres of spray solution. Consult the PINNACLE® and BASAGRAN FORTE® labels for the correct use rate for the broadleaf weeds present. Weeds controlled by 1.25 L/ha of ASSURE® in this tank mix are listed on this label at 1.0 L/ha (when ASSURE® is used alone). The increase in the application rate of ASSURE® in this tank mix is required for consistent grassy weed control.

For optimum leaf stage timing of broadleaf weeds, consult the PINNACLE® and BASAGRAN FORTE® labels. If leaf stages of the grass and broadleaf weeds do not coincide, a sequential application of the grass and broadleaf herbicides is required to ensure satisfactory control. Do not apply before the crop has 1-2 full trifoliate leaves.

Use the higher rate of CANPLUS® 411 or HI-MIX™ under conditions not conducive to good growth. DO NOT FREEZE.

Consult the PINNACLE® and BASAGRAN FORTE® labels for general information and use precautions.

ASSURE® + PINNACLE®

For the control of annual grassy and specific broadleaf weeds, apply a tank mixture of ASSURE® + PINNACLE®. Use ASSURE® at 1.0 litre per hectare plus PINNACLE® at 5.5-8.0 grams per hectare. Add CANPLUS® 411 or HI-MIX™ at 5-10 litres per 1000 litres of spray solution. Consult the complete ASSURE® directions for use pamphlet for grassy weeds controlled. Consult the PINNACLE® label for the correct use rate for the broadleaf weeds present.

Consult the timing section of this label for the correct timing of application to grassy weeds. For optimum leaf stage timing of broadleaf weeds, consult the PINNACLE® label. If leaf stages of the grass and broadleaf weeds do not coincide, a sequential application of the grass and broadleaf herbicides is required to ensure satisfactory control. Do not apply before the crop has 1-2 full trifoliate leaves.

Velvetleaf control may be reduced with a single tank mix application. For optimum control, make separate applications of ASSURE and PINNACLE Herbicides.

Use the higher rate of CANPLUS® 411 or HI-MIX™ under conditions not conducive to good growth. DO NOT FREEZE.

Consult the PINNACLE® label for precautions, weeds controlled and additional restrictions on the use of PINNACLE®.

SPRAY PREPARATION

EQUIPMENT - SPRAY VOLUME

Apply the spray mixture with ground equipment only. Use a properly calibrated sprayer that will provide thorough coverage of the weeds with the spray. Apply a minimum of 100 litres per hectare total spray volume. Use flat fan nozzles. DO NOT USE FLOOD JET NOZZLES.

Use 50 mesh screens or larger. A spray pressure of 210-275 kPa is recommended. Under heavy weed pressure, use increased volumes up to 400 litres total spray volume per hectare.

Avoid overlapping. Shut off spray booms while starting, turning, slowing or stopping to prevent over application.

Use spray preparation of ASSURE® within 24 hours or product degradation may occur. If spray preparation is left standing without agitation, thoroughly agitate before using.

NOTE: Extreme care must be taken to prevent drift to desirable plants or non-target agricultural land.

MIXING INSTRUCTIONS

Do not mix or load within 15 metres of all wells, including abandoned wells and drainage wells.

1. Make sure that spray tank is thoroughly cleaned before mixing.
2. Fill tank half full with water. KEEP AGITATOR RUNNING.
3. If tank mixing ASSURE with another herbicide, use the following sequence:
 - a) MUSTER® or PINNACLE®
 - b) BASAGRAN FORTE®
 - c) ASSURE®
 - d) Slowly add CANPLUS® 411 or HI-MIX™

Ensure that the herbicide is completely mixed before proceeding to the next step.

4. Add the rest of the required water to the tank. Mix well before applying to the crop.

On repeat tank loads, ensure that the amount of spray solution left in the tank from the previous load is less than 10% of volume about to be mixed.

Avoid over-filling of spray tank. Mix only enough product for the job at hand.

During spraying operation, make scheduled checks of spray equipment to help ensure proper application.

SPRAYER CLEAN-OUT

Thoroughly clean all traces of ASSURE® from application equipment immediately after use. Flush tank, pump, hoses, and boom with several changes of water after removing nozzle tips and screens (clean these parts separately). Failure to thoroughly clean the equipment may result in injury to subsequently sprayed grass crops.

USE PRECAUTIONS

CAUTION: DO NOT GRAZE THE TREATED CROPS OR HARVEST FOR FORAGE OR HAY; SUFFICIENT DATA ARE NOT AVAILABLE TO SUPPORT THIS USE.

Do not apply to soybeans within 80 days of harvest.

Do not apply to canola within 64 days of harvest.

Do not apply to flax within 82 days of harvest.

Do not apply to lentils or peas (field and processing) within 65 days of harvest.

Do not apply ASSURE® Herbicide to plants stressed by severe conditions such as drought, low fertility, or cool weather as control of weeds may be reduced.

Do not apply ASSURE® to crops stressed by severe conditions such as drought, low fertility, saline soils, water logged soils (soils at or near field capacity), disease or insect damage as crop injury may result.

Drought, disease or insect damage following application may also result in crop injury.

Rainfall within 1 hour of application may reduce effectiveness of the spray.

Avoid drift to other crops and non-target areas. Grass crops such as corn and wheat etc. are highly sensitive to ASSURE® Herbicide.

Do not tank mix ASSURE® Herbicide with any other pesticide, fertilizer, or additive except as directed in this label.

Do not contaminate irrigation water.

MANAGEMENT OF WEED RESISTANCE

Certain naturally occurring biotypes of wild oats and green foxtail have been identified in western Canada that are resistant to ASSURE® and related products (those with the same mode of action). Selection of resistant biotypes, through repeated use of these herbicides, may result in control failures. To delay selection for resistant populations of weeds, rotate the use of ASSURE® with herbicides that have a different mode of action and cultural practices that are effective in controlling these weeds.

If poor performance cannot be attributed to adverse weather conditions or improper application methods, a resistant biotype may be present. If resistance is suspected, contact your local representative of DuPont Canada Inc. for assistance. If resistance is identified, additional treatments with ASSURE® or related herbicides will be ineffective and are not recommended.

STORAGE AND DISPOSAL

STORAGE:

Store product in original container only, away from fertilizer, seed, food or feed. Not for use or storage in or around the home. Keep container **tightly** closed.

DO NOT FREEZE.

DISPOSAL:

1. Rinse the emptied container thoroughly and add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the Provincial Regulatory Agency or the Manufacturer.

NOTICE TO USER

This control product is to be used only in accordance with the directions on this label. It is an offence under the PEST CONTROL PRODUCTS ACT to use a control product under unsafe conditions.

Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

The seller warrants that the purchase by the buyer and the use of this product, as such, will not infringe any Canadian patent.

The DuPont Oval®, ASSURE®, MUSTER® and PINNACLE® are registered trademarks of E.I. du Pont de Nemours and Company. DuPont Canada Inc. is a licensed user.

HI-MIX™ is a trademark of DuPont Canada Inc.

BASAGRAN FORTE® is a registered trademark of BASF AG used under license by BASF Canada Inc.

CANPLUS® 411 is a registered trademark of ICI Americas inc.

31-DEC-1997

SUPPLEMENTAL LABEL

**For Quackgrass and Annual Grassy Weed Control in
Established Creeping Red Fescue for Seed Production**

ASSURE® HERBICIDE

EMULSIFIABLE CONCENTRATE

AGRICULTURAL

REGISTRATION NUMBER 22111

PEST CONTROL PRODUCTS ACT

GUARANTEE: Quizalofop - Ethyl 96 g/L

READ THE LABEL AND BROCHURE BEFORE USING

DUPONT CANADA INC
AGRICULTURAL PRODUCTS
BOX 2300, STREETSVILLE
MISSISSAUGA, ONTARIO
L5M 2J4
1-800-667-3925

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN
CAUTION! MAY IRRITATE EYES, NOSE, THROAT AND SKIN
WARNING - EYE AND SKIN IRRITANT

May be harmful if swallowed, inhaled or absorbed through skin. Cause: eye irritation.

Do not get in eyes or on clothing. Avoid contact with skin. Avoid breathing vapours or spray mist. Use adequate ventilation.

Wear goggles or face shield, chemical resistant gloves and coveralls during mixing, loading, clean up and repair.

For worker protection during application, wear a hat and coveralls.

When using, do not eat, drink or smoke. Wash skin thoroughly with soap and water after handling.

Remove contaminated clothing immediately after use. Store and wash contaminated clothing separately from household laundry before reuse

FIRST AID

IF SWALLOWED: If swallowed, DO NOT induce vomiting. Immediately give two glasses of water. Never give anything by mouth to an unconscious person. Call a physician or contact a poison control centre.

IF ON SKIN: In case of contact, immediately flush skin with plenty of water for at least 15 minutes while removing contaminated clothing and shoes. Call a physician or contact a poison control centre. Wash contaminated clothing before reuse.

IF IN EYES: In case of contact, immediately flush eyes with plenty of water for at least 15 minutes. Call a physician or contact a poison control centre.

IF INHALED: If inhaled, remove to fresh air. If not breathing, give

artificial respiration. If breathing is difficult, give oxygen. Call a physician or contact a poison control centre.

Never give anything by mouth to an unconscious person. Remove victim to safe, uncontaminated area. If victim has ceased breathing clear airway and start artificial respiration. Obtain medical attention or contact a poison control centre. Take container or label or the product name and product registration number with you to emergency/hospital or physician.

TOXICOLOGICAL INFORMATION:

THIS PRODUCT CONTAINS PETROLEUM DISTILLATES.

DIRECTIONS FOR USE

NOTICE TO BUYER/USER: READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR SPECIAL USE APPLICATIONS:

The DIRECTION FOR USE for this product for the use described on this Supplementary Label were developed by persons other than DuPont Canada Inc. and accepted for registration by Health Canada under the User Requested Minor Use Label Expansion Program. DuPont Canada Inc. itself makes no representation or warranty with respect to product performance (efficacy) and crop tolerance (phytotoxicity) of this product when used on the crop listed on this Supplementary Label. Accordingly, the Buyer and User assume all liability arising, and agree to hold DuPont Canada Inc. harmless from any claims based on efficacy or phytotoxicity in connection with the use described on this Supplementary Label.

POST EMERGENT APPLICATION:

Use on established creeping red fescue for seed production only.

DO NOT graze or cut treated established red fescue for feed in the year of treatment.

Apply 1.0 - 1.5 litres/ha of ASSURE® as a broadcast application. CANPLUS (R) 411 or HI-MIX (TM) should be added to the tank at a rate of 5 - 10 litres per 1000 litres of spray solution. Apply when quackgrass is in the 2 - 6 leaf stage and when annual grassy weeds are from the 2 leaf stage to early tillering. Application to the above species should be made before crop canopy closes. Refer to main label for list of grassy weeds controlled as well as the recommended ASSURE® rates depending on the leaf stage of quackgrass or the annual grass species that are present.

Make only one application per growing season.

Apply the spray mixture with ground equipment only. Use a properly calibrated sprayer that will provide thorough coverage of the weeds with

the spray solution. Apply a minimum of 100 litres/ha of total spray volume.

STORAGE:

Store product in original container only, away from fertilizer, seed, food or feed. Not for use or storage in or around the home. Keep container closed. DO NOT FREEZE.

DISPOSAL:

1. Rinse the emptied container thoroughly and add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the Provincial Regulatory Agency or the Manufacturer.

NOTICE TO USER

This control product is to be used only in accordance with the directions on this label. It is an offence under the PEST CONTROL PRODUCTS ACT to use a control product under unsafe conditions.

NOTICE TO BUYER

Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

The seller warrants that the purchase by the buyer and the use of this product, as such, will not infringe any Canadian patent.

ASSURE® is a registered trademark of E.I. du Pont de Nemour and Company. DuPont Canada Inc. is a licensed user.

CANPLUS® 411 is a registered trademark of ICI Americas inc.

HI-MIX™ is a trademark of DuPont Canada Inc.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))