

16-DEC-1997

RAMPAGE(R)

RAT AND MOUSE BAIT

KILLS RATS AND MICE

COMMERCIAL

FOR LICENSED PEST CONTROL OPERATORS

GUARANTEE:

CHOLECALCIFEROL 0.075%

READ THE LABEL BEFORE USING

KEEP OUT OF REACH OF CHILDREN

REGISTRATION NO. 19802  
PEST CONTROL PRODUCTS ACT

NET CONTENTS: (10 grams) .36 oz.

MANUFACTURED BY:

MOTOMCO LTD.

BOX 8422

Madison, WI 53708

U.S.A.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Place bait in protected places inaccessible to children and domestic animals or in tamper-resistant bait stations. Do not contaminate food or feed. Do not contaminate any body of water.

DIRECTIONS FOR USE: Use bait to control the Norway rat (*Rattus norvegicus*) the roof rat (*Rattus rattus*) and house mouse (*Mus musculus*) in and around dwellings serviced by pest control operators, farm buildings, food service establishments (non-food areas), granary bins (empty), processing plants (feed, food, non-food) non-food areas, storage areas (non-food).

For areas not directly to food processing: Use only in non-food or non-feed areas where feed, food, packaging and handling equipment are never opened or exposed.

For areas where or food is processed, served or stored: In meat and food processing plants (processing areas) use only when plant is not in operation. Remove or cover all food, packaging material and utensils before placing bait in baiting stations. Remove all baits and dead rodents before re-use of the plant. (Processing areas include storage and service.)

GENERAL: Place bait in locations frequented by the rodents, such as tunnels, feeding areas and runways. For rats place 60-100 g of bait per station. For mice set out 10 g of bait at intervals of 2-3 m. Maintain an uninterrupted supply until feeding ceases.

FIRST AID: If in eyes, flush with water for 15 minutes. If on skin, wash with soap and water. If swallowed, give patient one or two glasses of water and cause vomiting by inserting a finger down the throat, repeat until vomit is clear. Do not give anything by mouth to an unconscious person. Contact a physician or Poison Control Centre immediately.

TOXICOLOGICAL INFORMATION: If serum calcium levels are elevated with Calcitonin is effective in reducing calcium to normal levels. Continue monitoring serum calcium and treat as necessary for hypercalcemia.

DISPOSAL: Do not reuse empty container. Dispose of unused or spoiled bait in accordance with local requirements. Follow provincial instructions for any required cleaning of the container prior to its disposal. Make the empty container unsuitable for further use. Dispose of the container in accordance with provincial requirements. For more information on the disposal of unused, unwanted product and the cleanup of spills, contact the Provincial Regulatory Agency or the Manufacturer.

Rodent Disposal: Dispose of dead rodents in garbage or by burying.

NOTICE TO USER:

This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe

conditions.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))