

89.02.07

((CARTON))

SENDRAN (R) BREAK-AWAY

Collar for Cats of all Sizes

Kills Fleas & Ticks up to 4 Months

For continuous protection replace collar
when effectiveness diminishes

SOLD THROUGH VETERINARIANS

CAUTION:

DO NOT ALLOW CHILDREN TO PLAY WITH THIS COLLAR

DOMESTIC

READ LABEL BEFORE USING

GUARANTEE: Propoxur.....9.4%

CONTENTS: 1 Collar, Net Weight 8.35 g

Registration No. 18508 P.C.P. Act

Distributed by
BAYVET Division Chemagro Limited
Etobicoke, Ontario, Canada M9W 1G6

HAYER (R)

((INSERT))

CODE NO. 2383

SENDRAN (R)

Break-Away Collar
for Cats of all Sizes

DOMESTIC

GUARANTEE: Propoxur* 9.4%

REG. NO. 18508 PCP ACT

READ THE LABEL BEFORE USING

DIRECTIONS: Thread flat end of the collar through buckle hole and tighten as necessary, so that at least two ridges are covered by the buckle. Collar must be worn loosely, but securely enough to prevent easy removal. Cut off any excess length and discard in trash.

The expandable section molded into the collar will stretch under the weight of an average size cat and allow the cat to free itself if it becomes hooked or entangled by the collar. Since both the expandable section and plastic buckle are molded into the collar, the probability of neck irritation and buckle failure is reduced.

To release the collar, squeeze buckle between thumb and finger and pull sharply.

Do not use on kittens under 12 weeks of age.

The flea and tick collar starts killing fleas and ticks when it is placed around the cat's neck. Fleas will be killed and any new ones which may temporarily appear on the cat will also be killed during the 4 months. Non-engorged ticks are tough occasional pests of cats and will be killed within a few days. Collar will kill ticks encountered in first two months of wear. FOR CONTINUOUS PROTECTION, REPLACE COLLAR WHEN EFFECTIVENESS DIMINISHES.

Some cats may be sensitive to this collar. When collar is first worn, observe neck area every few days for irritation. Any collar, when fastened too tightly, may cause skin irritation. Remove collar at first sign of irritation or adverse reaction. Collar is intended for use only as an insecticide generator and is not to be taken internally by man or animals. Do not administer other cholinesterase inhibitors (e.g. oral parasiticides) while collar is worn or for a few days after removing collar.

PRECAUTIONS: Do not open inner envelope until ready to use. DO NOT ALLOW CHILDREN TO PLAY WITH UNATTACHED COLLAR. Dust will form on this collar during storage. Do not get dust or collar in mouth, harmful if swallowed. Wash hands thoroughly with soap and water after handling.

FIRST AID: In case of poisoning, call a physician immediately. If dust from collar contacts skin, wash thoroughly with soap and water. If dust contacts eyes, flush eyes with water and get medical attention.

TOXICOLOGICAL INFORMATION: Dust in eyes will cause temporary pupillary constriction*. Dust released from collar is a cholinesterase inhibitor.

NOTE TO PHYSICIAN: Atropine and homatropine* are antidotal.

SENDTRAN (R) is a Reg. TM of the parent company of Farbenfabriken Bayer, GmbH, Leverkusen.

BAYVET Division CHEMAGRO Ltd. Rexdale, Ontario M9W 1G6

PRINTED IN CANADA

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))