

91.04.29

CLEAN CROP (R)

MANEB 80W

FUNGICIDE

COMMERCIAL

CAUTION

POISON

READ LABEL BEFORE USING

GUARANTEE:

MANEB.....80%

REGISTRATION No.11034
PEST CONTROL PRODUCTS ACT

NET 10 kg

United Agri Products Canada Inc.
R.R. #2
DORCHESTER, ONTARIO
N0L 1G5

DIRECTIONS

Use MANEB 80W at 1.10 to 3.25 kg/ha or 65 grams per 10 litres except as noted below. Wash MANEB 80W through the screen or sift slowly into spray tank when partially filled. Add insecticides or nutritional sprays and spreaders last. MANEB 80W is compatible with most insecticides and may be used with most miticides. In regular spray equipment, use 800 to 1000 litres per hectare for low growing crops. Use 1000 to 1500 litres for heavy tomato and potato vines. MANEB 80W may be used in low volume or concentrate sprays by increasing the dosage accordingly, following local recommendations.

VEGETABLES:

Tomatoes - Early and late blight, Anthracnose, Septoria leaf spot, Cladosporium grey mold: Begin spraying when disease is first reported in area and repeat application at 7 to 10 day intervals or more often under severe blight conditions. Use MANEB 80W at the rate of 3.25 kg/ha. Apply at least 1500 litres of spray per hectare, per application when plants are full grown. Do not apply within 7 days of harvest.

Potatoes - Early and late blight: Apply at 7 to 10 day intervals throughout the season. Begin application early, applying 1.10 kg/ha as vines increase in size. Do not apply within 1 day of harvest.

Celery - Early and late blight: Begin spraying in the plant bed when plants first come through the ground. Make further application at 3 to 5 day intervals until plants are set in the field. Continue field application on regular weekly schedule. Do not apply within 14 days of harvest.

Carrots - Alternaria leaf blight, Cercospora leaf spot: Begin application when disease is reported in the area and continue at 7 to 10 day intervals during the growing season, or as required for adequate control. Do not apply within 5 days of harvest.

Cucumber, Muskmelon, Pumpkin and Squash - Anthracnose, Alternaria and Septoria leaf spot, Downy mildew: Start application in seeding stage repeating at weekly intervals or as necessary. Regular applications tend to suppress Powdery Mildew. Do not apply within 14 days of harvest.

Onions (Dry bulb only) - Downy mildew, Botrytis leaf blight, purple blotch: Apply when disease is reported in the area and continue at 7 to 10 day intervals during the season. Do not apply within 10 days of harvest.

PRECAUTION

May cause irritation of nose, throat, eyes and skin. Do not breathe dust or spray mist. Keep away from fire and sparks. Store in a cool dry place. Do not allow to become wet or overheated in storage. This may generate flammable vapours. Keep out of reach of children. Crush or puncture empty container and dispose of by burial or other proper means. Avoid contamination of water supplies.

FIRST AID

In case of skin contact, wash with soap and water. If taken internally induce vomiting by inserting finger in throat. Repeat until vomit fluid is clear. Drink copious quantities of water. If in eyes, flush with water for 5 to 10 minutes and get medical attention.

TOXICOLOGY

Treat as above for skin contact. If taken internally administer gastric lavage.

DISPOSAL

1. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
2. Make the empty container unsuitable for further use.
3. Dispose of the container in accordance with provincial requirements.
4. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the regional office of Environmental Protection, Environment Canada.

NOTICE TO USER - This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER - Seller's guarantee shall be limited to the terms of the label and subject thereto. The buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on these conditions.

APRIL 1991

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any

guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))