

96.08.08

SANEX(R)  
AMINE 500 FORESTRY HERBICIDE

COMMERCIAL

NET CONTENTS: 4 L (9.5 L) (20 L) (205 L)

GUARANTEE:

2,4-D (isomer specific) 470 g/L  
(present as amine)

READ THE LABEL BEFORE USING

REGISTRATION NO. 18113  
PEST CONTROL PRODUCTS ACT

WARNING POISON

SANEX AGRO INC.  
36 Head Street  
Dundas, Ontario  
L9H 3H3

DIRECTIONS:

DO NOT APPLY BY AIR

Frill or hack and squirt: Make notches (or frills) into trees with axe. Notches should angle downward to hold herbicide. For trees over 15 cm D.B.H. make a complete frill around tree as close to ground as possible. For trees less than 15 cm, make one notch per 3 cm diameter. Inject 1 mL of undiluted product, or up to a 2:1 dilution (water:2,4-D AMINE 500).

Stump treatment: To freshly cut stumps, apply undiluted or up to 2:1 dilution to the cambial area. Complete coverage of cut

surface should be made with brush or knapsack sprayer.

Lawns, turf: For annual and perennial broadleaf weeds (hedge bindweed, broadleaved plantain, burdock, dandelion, and other 2,4-D susceptible weeds, use 2.2-4.4 L per 2000 L of water to cover 1 hectare. Use sufficient water to obtain good coverage. Do not cut the grass for 4 days before or after spraying. May injure or kill bent grass and white clover. Repeat if necessary. If rain occurs within 24 hours after spraying, a second application may be necessary. On newly seeded lawns, do not spray within 4 weeks after grass is up and then only use half rate for one month.

For best results, spray in early fall and late spring when the weather promotes active weed growth and when the temperature is above 10°C. Use with caution during long hot dry spells. Avoid spraying if rain is expected shortly. Apply with pressure sprayer using a coarse spray, or use a watering can for smaller areas.

Warning: Care should be taken that the material does not drift to susceptible valuable plants. Do not use same sprayer for other agricultural purposes.

Re-entry interval for turf: Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treated areas until spray has thoroughly dried.

Time of application: For best results in selective weed control, the weeds should be actively growing. Therefore, application during the months of May, June and September is preferred. The effectiveness of a treatment in July and August will depend on general weather conditions. If it is hot and dry during that period, a second treatment may be necessary. If it rains within 24 hours after spraying, a repeat application may be required. Avoid mowing for several days before and after treatment. For ground-ivy (creeping Charlie), apply a spot treatment in June when the plants are young. A follow-up application in September is often necessary. For knotweed, apply in May or early June when the young plants come up. NEWLY SEEDED AREAS should not be sprayed within 4 weeks after grass emergence and then use only half the recommended rate for the first month.

Maximum application frequency to turf: For good turf/lawn management, normally two applications per year per treatment site are adequate. This does not include spot treatments.

Transfer System: Container larger than 20 litres: Use a transfer

system that avoids open pouring when transferring the liquid concentrate from such containers into the spray tank.

#### Protective Clothing and Equipment:

Handling the concentrate (mixing and loading): wear long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes and protective eye wear (face shield or safety glasses). Rinse gloves before removal. Coveralls or a chemical resistant apron should also be worn when open pouring from containers greater than 5L.

Handling the dilute spray solution (during application or repairing or cleaning equipment): wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes. Rinse gloves before removal. Gloves are not required during application when applicator is in an enclosed tractor.

#### Operator Use Precautions:

- Wear freshly laundered clothing and clean protective equipment daily.
- Rinse gloves before removal.
- Wash hands before eating, drinking, using tobacco or using the toilet. If herbicide penetrates clothing remove immediately, then wash thoroughly and put on clean clothing. Throw away clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate.
- After using this product, remove clothing and launder separately and promptly and thoroughly wash hands and exposed skin with soap and water. Follow manufacturer's instructions for cleaning personal protective clothing and equipment. If no such instructions for washables are provided use detergent and hot water. Keep and wash personal protective equipment separate from household laundry.
- After work, remove all clothing and shower using soap and water.

#### Re-entry

Cleaning equipment: All equipment must be cleaned thoroughly after use. Fill with hot water and detergent; flush all parts

several times. Let sprayer stand with solution for 24 hours then rinse with water. It is often more convenient to use a separate sprayer with coarse nozzles for applying weedkillers.

#### PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. Avoid breathing spray mist. Avoid spray drift. Keep in original container during storage. Do not allow spray, spray mist, visible or invisible vapours or air containing mist or vapours to contact plants other than those being sprayed. If 2,4-D or MCPA susceptible plants are growing within a distance of 400 metres from the area being sprayed, spray only when there is a light breeze away from the susceptible plants. Spray drifting can be reduced by using high volumes of water sprayed under low pressure, using coarse spray nozzles. Susceptible crops include:

beans	cabbage	cauliflower
grapes	tobacco	
sugar beets	tomatoes	
sunflowers	turnips	

as well as many other field, vegetable, ornamental and fruit crops. Do not contaminate any body of water, or watercourse, water used for livestock or for domestic purposes. Do not store open container near food, feedstuffs, seeds, fertilizers or pesticides including insecticides, fungicides and herbicides other than those containing 2,4-D or brush killers. Sprayers which have contained SANEX AMINE 500 FORESTRY should not be used to apply other pesticides to 2,4-D or MCPA susceptible crops.

#### FIRST AID:

If on skin, wash with soap and water. If in eyes, flush with clean water for 15 minutes. If irritation develops, consult a physician or contact a Poison Control Centre immediately. If swallowed, drink 1-2 glasses of water to dilute the product. Do not induce vomiting. Call a doctor or Poison Control Centre immediately.

#### TOXICOLOGICAL INFORMATION:

May cause eye irritation on contact.

#### DISPOSAL:

Rinse the emptied container thoroughly and add the rinsings to the spray mixture in the tank. Follow provincial instructions for any required additional cleaning of the container prior to its disposal. Make the container unsuitable for further use. Dispose of the container in accordance with provincial requirements. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the provincial regulatory agency or the manufacturer.

NOTICE TO USER:

This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER:

Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

(R)Registered trademark of Sanex Agro Inc.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+) )