

89.11.03

WILSON
ROTENONE SPRAY ORGANIC INSECTICIDE
WETTABLE POWDER
DOMESTIC

GUARANTEE:
Rotenone..... 5%

REG. NO. 15415 P.C.P. ACT

Low toxicity organic insecticide for use on vegetables
up to the day before harvest.

READ THE LABEL BEFORE USING

Net Weight: 250 g

WILSON LABORATORIES INC.
DUNDAS, ONTARIO L9H 3H3

Rotenone insecticide is made from the roots of tropical plants and is therefore of organic origin. It has a low toxicity to people, domestic animals and is non-toxic to honey bees. Rotenone may safely be used on edible fruits and vegetables up to one day before harvest.

DIRECTIONS: Mix 30 mL (2 level tablespoonfuls) in 3 litres of water and stir well. For squash vine borer, corn borer and fruit insects apply as stated below. For other insects apply when they appear and repeat at 5 to 7 day intervals as necessary (insect eggs are not killed and will hatch). Spray upper and lower surface of foliage and on fruit and flowers. Shake sprayer occasionally while spraying to keep spray well mixed. May be applied on edible fruits and vegetables up to 1 day before harvest.

VEGETABLES: Asparagus, Beans, Beet, Broccoli, Brussels Sprouts,

Cabbage, Cauliflower, Corn, Cucumber, Eggplant, Celery, Lettuce, Melons, Onions, Peas, Peppers, Pumpkin, Potato, Radish, Spinach, Swiss Chard, Squash, Tomato.

SMALL FRUITS: Raspberry, Blackberry, Currant, Gooseberry, Grape, Strawberry.

ALL FLOWERING PLANTS: INSECTS CONTROLLED: Aphids, Asparagus Beetle, Bean Leaf Beetle, Beet Webworm, Cabbage Looper, Cabbageworm, Colorado Potato Beetle, Corn Borer, Cucumber Beetle, Diamond-Back Moth, Flea Beetles, Hornworms(Young), Leafhoppers, Squash Vine Borer, Thrips, Pea Weevil, Fruitworms, Raspberry Sawfly, Mexican Bean Beetle, Currantworm, Grape Leaf Skeletonizer, Leafrollers.

FOR SQUASH VINE BORER: Apply weekly to the base of the plant beginning when first leaves appear.

FOR CORN BORER: Apply weekly to the centre of the leaf whorls starting about mid-June.

RASPBERRY, BLACKBERRY: Apply when blossom buds appear and repeat just before blossoms open.

CURRENTS: Apply soon after fruit set, repeat one week later.

GOOSEBERRIES: Apply when berries turn red and webbing is noticed.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Avoid breathing spray mist. Avoid contact with skin, eyes, clothing. Wash after handling. Avoid contamination of food, feed and water. Toxic to fish and hogs.

FIRST AID: If swallowed, induce vomiting and call a doctor. If on skin wash with soap and water. If in eyes, flush with water for 15 minutes and obtain medical attention.

DISPOSAL: Dispose of empty container with regular garbage.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from

reliance upon this service.

+))