

90.06.29

CLEAN CROP(R)

METHOXYCHLOR 25% EC

COMMERCIAL

READ THE LABEL BEFORE USING.

REGISTRATION NO. 10690 PEST CONTROL PRODUCTS ACT

GUARANTEE: Methoxychlor 25%

UNITED AGRI PRODUCTS

P.O. Box 22116

London, Ontario

N6C 4N0

NET: 4 L

25621

05/89

DIRECTIONS FOR USE: Use only at recommended rates. Use in sufficient water for thorough coverage (usually 1000 L of water is required to cover 1 ha). Begin treatment at first sign of insects.

GARDEN CROPS: Use 3.2-9.2 L per ha on the following crops (except asparagus).

BEANS: Bean leaf beetles, Bean weevils, Leafhoppers and Mexican bean beetles.

BEETS, CARROTS, EGGPLANTS, LETTUCE, PEPPERS, RADISHES, SPINACH & TURNIPS: Blister beetles, Flea beetles and Leafhoppers.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE & CAULIFLOWER: Cabbage worms and Flea beetles.

CUCUMBERS, PUMPKINS & SQUASH: Cucumber beetles, Flea beetles and Leafhoppers.

ONIONS: Thrips.

PEAS: Pea weevils.

POTATOES: Blister beetles, Colorado potato beetles, Flea beetles and Leafhoppers.

TOMATOES: Blister beetles, Fall armyworms, Flea beetles and Tomato fruit worms.

ASPARAGUS: Asparagus beetles - use 3.5-8.5 L per ha. Amount to be used depends on the size of the crop plants.

NOTE: Use with caution on cucurbits. Do not use on young lima bean plants. Do not feed treated crops or crop refuse to animals. Do not apply within 1 day of harvest on cucumbers, eggplants, peppers, pumpkins, squash and tomatoes; within 3 days on asparagus and cabbage; within 7 days on beans, beets, cauliflower, onions, peas, potatoes, radishes and turnips; and within 14 days on broccoli, brussels sprouts, lettuce and spinach. May be used on carrots or turnips within 1 day of harvest if tops are not to be used for food or feed.

FIELD & FORAGE CROPS: ALFALFA & CLOVER: Use 4.2-5.5 L to control Alfalfa weevils. Flea beetles and Grasshoppers. Use 8.5-11 L per ha to control Armyworms. Do not apply within 7 days of harvest or grazing. Do not apply during bloom.

TREE FRUITS: APPLES, APRICOTS, CHERRIES, PEACHES, PEARS & PLUMS: To control Cherry fruit flies and Plum curculio, use 3.7-7.5 L per 1000 L of water and give sufficient spray to cover all foliage and fruits thoroughly. Start spraying at petal fall and use 2-3 applications at 7-14 day intervals as needed. Do not apply within 7 days of harvest on apples, cherries, pears and plums; and within 21 days on apricots and peaches. Do not spray during bloom.

SMALL FRUITS: CURRANTS, GOOSEBERRIES & STRAWBERRIES: To control Currant fruit flies and Strawberry weevils, use 4.2-8.5 L in 1000 L of water per ha. Do not apply within 14 days of harvest. Do not apply during bloom.

BRUCE SPANWORMS & WINTER MOTHS ON TREE FRUITS & ORNAMENTAL TREES (B.C. only): Mix 400 mL per 100 L of water and spray trees thoroughly. Apply in the spring at the time when blossom buds on the trees are in the pre-pink stage (before blossom). Use a second spray after blossom time if required. Do not spray during bloom. Do not apply to fruit trees within the following pre-harvest intervals: 7

days - apples, cherries, pears and plums. 21 days - apricots, peaches.

For best results follow provincial recommendations for timing.

ORNAMENTALS, TREES, SHRUBS, FLOWERS & HOUSE PLANTS: To control Caterpillars, Elm leaf beetles, Flea beetles, Lace bugs, Leafhoppers, May beetles, Tent caterpillars, Thrips, Rose chafers, Rose leaf beetles, Rose slugs, Webworms and Weevils. Add 4 L of Methoxychlor to 100 L of water to make a 1% solution. Spray plants thoroughly.

ELM BARK BEETLES: Mix 8 L of METHOXYCHLOR 25% EC in 100 L of water (per tree). Apply in the late winter or early spring when trees are dormant.

PICNIC AREAS, BACKYARDS & PATIOS: To control Ants, Centipedes, Crickets, Flies, Gnats, Hornets, Mosquitoes, flying Moths and Wasps, use a 3% solution by mixing 1 L of METHOXYCHLOR 25% EC in 7 L of water. For flying insects, allow to drift over the area being treated. Repeat if necessary. For crawling insects, spray directly on pests or on the surface of the infested area.

NON CROP VEGETATION & OTHER OUTDOOR AREAS: To control Black flies, Gnats and adult Mosquitoes, use a 5% solution by mixing 1 L of METHOXYCHLOR 25% EC with 4 L of water. For Aircraft Application apply 4.2-5.5 L per ha. Apply with care near streams, lakes or ponds containing fish.

For Ground Application using a thermal fogger, apply 0.7-1.5 L per ha. Repeat as necessary.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Avoid inhaling spray mist. Avoid contact with skin, eyes and clothing. Avoid contamination of feed and food. Wash after handling or using. Avoid contamination of ponds, streams and other bodies of water which contain fish life or which may be used for irrigation or domestic purposes. Care should be taken in the disposal of empty containers.

FIRST AID: For external contact, wash with soap and water. If in eyes, flush with plenty of water. If swallowed, do not induce vomiting. Get prompt medical attention.

TOXICOLOGICAL INFORMATION: Slightly irritating to the skin. Xylene is used as a solvent.

DISPOSAL:

1. Rinse the emptied container thoroughly and add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product and the clean-up of spills contact the regional office of Environmental Protection, Environment Canada.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes any risk to persons or property arising from the use of handling and accepts the product on that condition.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))