

85.12.31

Read the Label Before Using

STANCHEM

COMMERCIAL FOG APPLICATION

SPROUT-NIP SOLUTION

SPROUT INHIBITOR FOR POTATOES IN STORAGE

For Use ONLY By Approved Operators

GUARANTEE:

Chlorpropham (Food Grade) 475 grams per litre

REGISTRATION NO. 11860 PEST CONTROL PRODUCTS ACT

A DIVISION OF PPG INDUSTRIES CANADA LTD.  
60 TITAN ROAD, TORONTO, ONTARIO M9Z 2J8  
MONTREAL, WINNIPEG, EDMONTON, CALGARY

SPROUT-NIP SOLUTION  
(Fog application)

NOTICE TO USER:

THIS CONTROL PRODUCT IS TO BE USED ONLY IN ACCORDANCE WITH THE DIRECTIONS ON THIS LABEL. IT IS AN OFFENCE UNDER THE PEST CONTROL PRODUCTS ACT TO USE A CONTROL PRODUCT UNDER UNSAFE CONDITIONS.

Sprout-Nip Solution is formulated for application as a mist or fog to inhibit sprouting of potatoes in storage. It can be used with mechanical or thermal generators. Potatoes to be treated must be firm and free from excessive dirt or other debris. Mechanically damaged potatoes cannot be treated until suberization is complete, a process which normally takes 2-3 weeks or more.

DIRECTIONS FOR USE:

Seal the potato storage building to prevent or reduce escapes. Adjust the ventilation system to ensure even distribution of moving air. Adjust the application equipment to deliver droplets in the range one to ten microns. Apply sufficient Sprout-Nip Solution to achieve a deposit on the potatoes in the range 3 to 12 p.p.m. Shut off the ventilation system for 48 hours after treatment is completed.

HANDLING PRECAUTIONS:  
Irritating to lungs during application

1. Prolonged application with thermal generators may cause fires. Remove all flammable material from the vicinity of the application point before operations commence. Keep at least two operational fire extinguishers suitable for use in confined spaces at the point of application. Do not leave equipment unattended during operation.
2. The fog from this product is irritating to lungs. Wear a nose mask fitted with a filter if it is necessary to enter the storage building during application or before the fog has settled after application.
3. This product will inhibit sprouting of potatoes used for seed. Do not apply on or near potatoes to be used for this purpose. Do not store potatoes to be used for seed in treated buildings.
4. Low doses of this product will cause internal sprouting. Potatoes stored adjacent to treated potatoes may develop internal sprouts.

FIRST AID

In case of skin contact, wash immediately with soap and water and remove contaminated clothing. If skin irritation occurs obtain medical attention. In case of eye contact, flush with plenty of water and obtain medical attention. If swallowed induce vomiting. If patient is unconscious give him air; call physician or poison control centre.

CONTAINER STORAGE AND DISPOSAL:

Store away from food, feed, seed and potatoes. Make container unusable by puncturing and crushing immediately application is completed.

WARNING

Registration and safety data of this product are under review. Directions for Use and Cautionary Statements should be carefully followed. Read the Label Before Using.

NOTICE TO BUYER:

Seller's guarantee shall be limited to the terms set out on the label and subject thereto. The buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))