

09-FEB-2006

29-JAN-2008 Notification - Change in Registrant

**LATER'S
CASORON[®] WEED PREVENTER**

Controls grasses and weeds around Roses, Rhododendrons, Evergreens,
Fruits and Woody Ornamentals.

Casoron is a research discovery of Solvay Duphar B.V.

DOMESTIC

REG. NO. 16817 P.C.P. ACT

GUARANTEE: Dichlobenil 2%

READ THE LABEL BEFORE USING
KEEP OUT OF REACH OF CHILDREN

2 kg

Notification Change

~~Spectrum Brands IP Inc.
10 Craig Street
Brantford, ON
N3R 7J1~~

Sure-Gro IP Inc.
P.O. Box 21001
Brantford ON
N3R 7W9

DIRECTIONS: Later's **CASORON Weed Preventer** controls many annual grasses, broadleaved weeds and certain perennial weeds around Woody Ornamentals, Fruit Trees, Grapes, Raspberries, High Bush Blueberries and Saskatoons.

CONTROLS:		
Annual Bluegrass	Artemisia*	Bindweed*
Blue Aster	Brackenfern	Canada Thistle*
Chickweed	Crabgrass	Couch Grass*
Dandelion	Foxtail	Groundsel
Horsetail	Knotweed	Kochia
Lambsquarters	Loosestrife	Mustard
Nutsedge*	Pigweed	Plantain
Purslane	Sheep Sorrel*	Shepherd's Purse
Smartweed	Sow Thistle	Spurge
Vetch*	Wild Buckwheat*	

*controlled with the higher rates with late Fall applications

For control of annual weeds apply **CASORON** only to prepared weed-free soil either in early spring or late fall before seeds of annual weeds germinate or after cultivation has removed weeds. Apply uniformly over the soil surface.

Do not apply until 4 weeks after transplanting. For best results **CASORON** should not be applied during periods of high soil temperatures. Follow specific instructions for the type of tree, vine or bush being treated.

BEFORE TREATMENT:

1. Remove existing weeds
2. Thoroughly work and smooth the area to be treated
3. Measure area to be treated and determine number of square metres
4. If area to be treated is large, it should be divided into smaller sections and each section treated individually.

AFTER TREATMENT:

1. If application is made on dry soil, sprinkle lightly to moisten soil, approximately 1 cm deep. Apply 1.5 – 2.5 cm of water. Avoid puddling and washing. **DO NOT FLOOD.**
2. Do not disturb soil in treated area as long as the weed control program is being maintained.

1 level tablespoon = 10 g

1 m = 1 long pace

10 m² = 5 m x 2 m

HOW TO USE:

1. Shake **CASORON Granules** on the soil at a rate of 20 g per square metres of area to be treated – 200 g treats 10 m². Do not exceed this dosage. Rates higher than this may cause injury to desirable plants.
2. As a guide for the proper application rate and uniformity of distribution of the granules, consult the diagram: Application rate (actual size). Sprinkle evenly as shown.
3. A more uniform and accurate application can be obtained by applying half the granules over the entire area and then repeating with the remaining granules.
4. CASORON granules should not be used on sloping ground where run-off can occur. The concentrated run-off could cause damage to plants and lawns below.

WOODY ORNAMENTALS (established): **CASORON** may be used for weed control around the plants listed below.

Arborvitae	Ash	Barberry
Birch (cutleaf weeping)	Boxwood	Caragana
Cedar (Eastern Red)	Cedar (White)	Crabapple
Elm	Euonymous	Forsythia SP
Heather	Holly**	Honeysuckle
Juniper	Lilac	Linden
Locust	Maple	Mock Orange
Rhododendron	Rose	Spirea
Willow	Yew (Taxus)	

**Except *I. crenata*, *I. rotunda*, *I. vomitoria*

NOTE: Do not use on Gladiolus, Mugho Pine, Firs (*Abies* species such as Balsam Fir, *Abies con-colour*), Hemlock, Ajuga, certain Hollies (*Ilex crenata*, *I. rotunda*, *I. vomitoria*), Spruce, nor on herbaceous perennials (avoid use on plants which die down to the ground in the fall) and use with caution on shallow rooted ground covers (*Euonymus* can be treated), and other plants from list above.

COUCH GRASS AND ARTEMISIA CONTROL IN ESTABLISHED WOODY ORNAMENTALS: Couch Grass and Artemisia can be controlled in the woody ornamentals listed above (except Boxwood, *Euonymus*, Forsythia, Heather) which has been planted for at least one year. Apply CASORON as a uniform coverage in late fall from October 15th to December 15th and reapply in very early spring, before May 1. Use at the rate of 25 g per square metre for each treatment.

FRUIT TREES: CASORON may be used for weed control around Apple, Cherry, Peach, Pear and Plum trees. Follow application directions. Do not apply until 4 weeks after transplanting.

HIGHBUSH BLUEBERRIES: Apply uniformly during the dormant period (late winter).

RASPBERRIES: Treat established Raspberry plantings. Apply CASORON in late fall. Do not cultivate or work into the soil. Do not apply in the spring as injury may occur.

ESTABLISHED SASKATOONS: Weeds controlled as per above listing including Chickweed, Stinkweed, Groundsel, Pigweed, Dandelion and some grasses. Apply to Saskatoons, established for one or more years. Apply with a granular applicator in the late fall but before freeze-up. Use 20-35 g for square metre of CASORON granules on a broadcast basis. Do not apply within 9 months of harvest.

GRAPES: Apply only to control weeds under grapes established for at least 2 full years. Do not treat soil the year before planting or replanting grapes.

1. Treat when vines are completely dormant.
2. Treat only healthy vines.
3. Treat only vines established for at least 2 full years.
4. Apply in cool weather on moist soil.
5. Avoid local over-application to base of them.

NOTE: If spring application is made, apply before the weeds germinate a month before the grape buds burst. Do not apply on hazy, hot days. To control seedling weeds apply in early spring to fall. To control perennial weeds apply in fall.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed. Avoid breathing dust. Avoid skin, nose or eye contact. Avoid contamination of feed and foodstuffs.

ENVIRONMENTAL HAZARDS: Toxic to fish and other aquatic life. Do not use on or around edible plants. Do not use on woody ornamentals or fruit crops other than those listed on the label. Do not use in seed beds, transplant or cutting beds. Do not use in or around greenhouses. Application in spring should be well made before bud burst. Do not plant vegetables or other sensitive crops the year following soil treatment. Do not use on light sandy soil.

FIRST AID: Call a doctor or poison control centre in case of accidental poisoning. **If swallowed**, induce vomiting, repeat until vomit fluid is clear. Rush patient to nearest hospital taking this labeled container with you. **If in eyes**, flush with clean water for 15 minutes and go to the nearest hospital at once. **If on skin**, remove contaminated clothing and wash with soap and water. Take container, label or product name and P.C.P. Registration Number with you when seeking medical attention.

STORAGE: Keep product in tightly closed and labelled original container. Store in a cool, dry place away from food.

DISPOSAL: Do not reuse empty container. Wrap container in newspaper and dispose of in household garbage.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and, subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.