

26-MAR-1997

((LABEL))

INTER-AG
Division of AgrEvo

RINSE For Dollars
RINSE For Safety
RINSE For Disposal

AGRICULTURAL

WARNING POISON

MCPA
Sodium Salt 300

10 LITRES

GUARANTEE:
MCPA 300 g/L
(Present as Sodium Salt)

REGISTRATION NO. 21485
PEST CONTROL PRODUCTS ACT

READ LABEL AND ATTACHED DIRECTIONS FOR USE BEFORE USING

AgrEvo Canada Inc.
295 Henderson Drive
Regina, SK
S4N 6C2
Telephone : 1-800-667-5959

PT#LA-M 35 11/96 Printed in the USA

PRECAUTIONS

1. Keep out of the reach of children.
2. Avoid contact with eyes, skin and clothing. May cause skin irritation.
3. Do not breathe spray mist or vapours. Avoid spray or vapour drifts onto susceptible plants such as grapes, tomatoes, tobacco, sugar beets, sunflowers, beans, turnips, cauliflower, cabbage,

alfalfa, clover, peas, potatoes, ornamental trees, shrubs or flowers, fruit trees, lawns, most vegetable crops or other desirable vegetation.

4. Do not spray when there is any danger of wind or on exceptionally hot days over 27°C.
5. Drift from spray may be reduced by using high volume sprays under low pressure, coarse sprays, and drop nozzles. Use special care in airplane application where danger from drift is greater.
6. For aerial application, exercise special care:
 - 1) Do not spray in winds exceeding 8 km per hour.
 - 2) Do not spray in dead calm near sensitive plants. A cloud of suspended droplets may drift onto sensitive plants when the wind comes up.
 - 3) Spray only when the wind is blowing away from a sensitive crop, shelter belt or garden.
7. Do not contaminate any body of water. Avoid contamination of foods.
8. Do not use in a greenhouse.
9. Keep in original container during storage.
10. Do not store near or in same room as seeds, feeds, fertilizers or pesticides.
11. Clean spray equipment thoroughly after use. Do not use spray equipment for other purposes. Do not re-use empty containers.
12. Allow 7 days between application of product and grazing/cutting for hay.

DISPOSAL INFORMATION

1. Triple rinse the emptied container thoroughly and add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.

5. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the Provincial Regulatory Agency or the Manufacturer.

FIRST AID

1. In case of poisoning, call a physician or contact a poison control centre immediately.
2. If on skin, remove contaminated clothing and wash skin thoroughly with soap and water.
3. If in eyes, flush with water immediately and continue for 5-10 minutes. Get medical attention, or contact a poison control centre.
4. If swallowed, induce vomiting then take patient to nearest hospital or doctor's office, taking the pesticide label with you, or contact a poison control centre.

NOTICE TO USER

This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

((BROCHURE))

INTER-AG
Division of AgrEvo

RINSE For Dollars
RINSE For Safety
RINSE For Disposal

AGRICULTURAL

WARNING POISON

MCPA
Sodium Salt 300

10 LITRES

GUARANTEE:
MCPA 300 g/L
(Present as Sodium Salt)

REGISTRATION NO. 21485
 PEST CONTROL PRODUCTS ACT

READ LABEL AND ENCLOSED BROCHURE PRIOR TO USING.

AgrEvo Canada Inc.
 295 Henderson Drive
 Regina, SK
 S4N 6C2
 Telephone : 1-800-667-5959

PT#LA-M 35 11/96 Printed in the USA

LABEL INDEX MCPA SODIUM SALT 300	
Wheat, Barley, Rye, Oats GO TO Section 1	Cereals underseeded to Legumes GO TO Section 6
Flax GO TO Section 2	Filling Spray Tank GO TO Section 7
Winter Wheat, Fall Rye GO TO Section 3	Precautions GO TO Section 8
Corn GO TO Section 4	First Aid GO TO Section 9
Peas (Field & Canning) GO TO Section 5	Rinsing and Disposal GO TO Section 10
SECTION 1	
CROP:	WHEAT, BARLEY, SPRING RYE, OATS
TREATMENT STAGE:	For ground or air application, treat from the 2nd leaf to early flag leaf stage.
SUSCEPTIBLE WEEDS:	Burdock, Cocklebur, Mustards (except Dog Mustard), Ragweed, Flixweed*, Kochia, Lamb's-quarters, Russian Pigweed, Shepherd's Purse, Stinkweed. * Flixweed susceptible only at seedling stage. Use rates for harder to kill weeds to kill fall germinated flixweed in the spring.
RATES TO USE:	
Small seedlings, good growing conditions:	1.25 L/ha

Weeds in bud, dry conditions, very heavy infestation:	2 L/ha
HARDER TO KILL WEEDS:	Annual Sow Thistle, Bluebur, Blue Lettuce, Canada Thistle*, Peppergrass, Curled Dock, Dog Mustard, Goose Foot, Hempnettle (suppression only), Red Root Pigweed, Russian Thistle, Perennial Sow Thistle*, Leafy Spurge*. * Top growth control only.
RATES TO USE ON HARDER TO KILL WEEDS:	
Small seedlings, growing rapidly, good growing conditions:	2 L/ha
Weeds in bud, dry or cool conditions, heavy infestation:	2.75 L/ha NOTE: These high rates may result in deformities but improved weed control should result in higher yields.
GO TO SECTION 7	
SECTION 2	
CROP:	FLAX (Do not use on low linolenic acid varieties).
TREATMENT STAGE:	Spray after plants have reached 5 cm in height and before buds appear.
SUSCEPTIBLE WEEDS:	Burdock, Cocklebur, Mustards (except Dog Mustard), Ragweed, Flixweed*, Kochia, Lamb's-quarters, Russian Pigweed, Shepherd's Purse, Stinkweed. * Flixweed susceptible only at seedling stage. Use rates for harder to kill weeds to kill fall germinated flixweed in the spring.
RATES TO USE ON SUSCEPTIBLE WEEDS:	
Small seedlings growing rapidly:	1.25 L/ha
Weeds in a bud, dry or cool weather, heavy infestations	2 L/ha

HARDER TO KILL WEEDS:	Annual Sow Thistle, Bluebur, Blue Lettuce*, Canada Thistle*, Peppergrass, Curled Dock, Dog Mustard, GooseFoot, Hempnettle (suppression only), Redroot Pigweed, Russian Thistle, Perennial Sow Thistle*, Leafy Spurge*. * Top growth control only.
RATES TO USE ON HARDER TO KILL WEEDS:	
Small seedlings, growing rapidly, good growing conditions:	2 L/ha
Weeds in bud, dry or cool weather, heavy infestation:	2.75 L/ha NOTE: These rates may result in deformities but improved and weed control should result in higher yields.
GO TO SECTION 7	
SECTION 3	
CROP:	WINTER WHEAT, FALL RYE
TREATMENT STAGE:	Treat in spring when weeds are small from 2nd leaf to flag leaf stage.
SUSCEPTIBLE WEEDS:	Burdock, Cocklebur, Mustards (except Dog Mustard), Ragweed, Flixweed*, Kochia, Lamb's-quarters, Russian Pigweed, Shepherd's Purse, Stinkweed. * Flixweed susceptible only at seedling stage. Use rates for harder to kill weeds to kill fall germinated flixweed in the spring.
RATES TO USE ON SUSCEPTIBLE WEEDS:	
Small seedlings growing rapidly:	1.25 L/ha
Weeds in bud, dry or cool weather, heavy infestation:	2 L/ha
HARDER TO KILL WEEDS:	Annual Sow Thistle, Bluebur, Blue Lettuce*, Canada Thistle*, Peppergrass, Curled Dock, Dog Mustard, Goose Foot, Hempnettle (suppression only), Redroot Pigweed, Russian Thistle, Perennial Sow Thistle*, Leafy Spurge*. * Top kill only.

RATES TO USE ON HARDER TO KILL WEEDS:	
Small seedlings, growing rapidly, good growing conditions:	2 L/ha
Weeds in bud, dry or cool weather, heavy infestation:	2.75 L/ha NOTE: These rates may result in deformities but improved weed control should result in higher yields.
GO TO SECTION 7	
SECTION 4	
CROP:	CORN
TREATMENT STAGE:	Treat when weeds are small until corn reaches 15 cm in height. When corn is taller than 15 cm, a directed spray should be used.
SUSCEPTIBLE WEEDS:	Burdock, Cocklebur, Mustards (except Dog Mustard), Ragweed, Flixweed*, Kochia, Lamb's-quarters, Russian Pigweed, Shepherd's Purse, Stinkweed. * Flixweed susceptible only at seedling stage. Use rates for harder to kill weeds to kill fall germinated flixweed in the spring.
RATES TO USE ON SUSCEPTIBLE WEEDS:	
Small seedlings, good growing conditions:	1 L/ha
Weeds in bud, dry or cool weather, heavy infestation:	1.5 L/ha
GO TO SECTION 7	
SECTION 5	
CROP:	PEAS (FIELD and CANNING)

TREATMENT STAGE:	Not recommended in Nova Scotia. Treat when pea vine length is 10 to 20 cm but before flowering begins. Use at least 180 Litres of water per hectare.
SUSCEPTIBLE WEEDS:	Mustards (except Dog Mustard), Ragweed, Flixweed*, Kochia, Lamb's-quarters, Russian Pigweed, Shepherd's Purse, Stinkweed. * Flixweed susceptible only at seedling stage. Use rates for harder to kill weeds to kill fall germinated flixweed in the spring.
RATES TO USE ON SUSCEPTIBLE WEEDS:	
	1 L/ha
HARDER TO KILL WEEDS:	Annual Sow Thistle, Bluebur, Blue Lettuce*, Canada Thistle*, Peppergrass, Curled Dock, Dog Mustard, Goosefoot, Hempnettle (suppression only), Redroot Pigweed, Russian Thistle*, Perennial Sow Thistle*, Leafy Spurge*. * Top growth control only.
RATES TO USE ON HARDER TO KILL WEEDS:	
	1.25 L/ha
GO TO SECTION 7	
SECTION 6	
CROP:	CEREALS UNDERSEEDED TO LEGUMES - ALFALFA, RED AND ALSIKE CLOVER
TREATMENT STAGE:	For emergency treatment only. Use 2,4-DB or MCPB if possible. Do not spray sweet clover or early Flemish alfalfa varieties with MCPA. Treat when cereals and weeds are up high enough to offer protection to the legume. Do not treat after the shot blade stage of cereals.
SUSCEPTIBLE WEEDS:	Mustards (except Dog Mustard), Ragweed, Flixweed, Kochia, Lamb's-quarters, Russian Pigweed, Shepherd's Purse, Stinkweed.
RATES TO USE ON SUSCEPTIBLE WEEDS:	
	1 L/ha
GO TO SECTION 7	

SECTION 7

FILLING SPRAY TANKS

1. Fill tank 1/2 full with water.
2. Add MCPA Sodium Salt 300.
3. Complete filling spray tank with water.
4. Use 60 - 240 litres of water per hectare unless other specified.
5. Do not exceed 275 kPa pressure.

GO TO SECTION 8

SECTION 8

PRECAUTIONS:

1. Keep out of reach of children.
2. Avoid contact with eyes, skin and clothing. May cause skin irritation.
3. Do not breathe spray mist or vapors. Avoid spray or vapor drifts onto susceptible plants such as grapes, tomatoes, tobacco, sugar beets, sunflowers, beans, turnips, cauliflower, cabbage, alfalfa, clover, peas, potatoes, ornamental treat, shrubs or flowers, fruit trees, lawns, most vegetable crops or other desirable vegetation.
4. Do not spray when there is any danger of wind or on exceptionally hot days over 27 degrees C.
5. Drift from spray may be reduced by using high volume sprays under low pressure, coarse sprays, and drop nozzles. Use special care in airplane application where danger from drift is greater.
6. For aerial application, exercise special care:
 - 1) Do not spray in winds exceeding 8 km per hour.
 - 2) Do not spray in dead calm near sensitive plants. A cloud of suspended droplets may drift onto sensitive plants when the wind comes up.
 - 3) Spray only when the wind is blowing away from a sensitive crop, shelter belt or garden.
7. Do not contaminate any body of water. Avoid contamination of foods.
8. Do not use in a greenhouse.
9. Keep in original container during storage.
10. Do not store near or in same room as seeds, feeds, fertilizers or pesticides.
11. Clean spray equipment thoroughly after use. Do not use spray equipment for other purposes. Do not reuse empty containers.
12. Allow 7 days between application of product and grazing/cutting for hay.

GO TO SECTION 9

FIRST AID:	<ol style="list-style-type: none"> 1. In case of poisoning, call a physician or contact a poison control centre immediately. 2. If on skin, remove contaminated clothing and wash skin thoroughly with soap and water. 3. If in eyes, flush with water immediately and continue for 5-10 minutes. Get medical attention or contact a poison control centre. 4. If swallowed, induce vomiting. Rush patient to nearest hospital or doctor's office, taking the pesticide label with you, or contact a poison control centre.
GO TO SECTION 10	
SECTION 10	
DISPOSAL INFORMATION: RINSE For Dollars RINSE For Safety RINSE For Disposal	<ol style="list-style-type: none"> 1. Triple rinse the emptied container thoroughly and add the rinsings to the spray mixture in the tank. 2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal. 3. Make the empty container unsuitable for further use. 4. Dispose of the container in accordance with provincial requirements. 5. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the Provincial Regulatory Agency or the Manufacturer.
NOTICE TO USER:	This control product is to be used only in accordance with the directions on this label. It is an offense under the Pest Control Products Act to use a control product under unsafe conditions.
NOTICE TO BUYER:	Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

Inter-Ag is a registered trademark of Hoechst Schering AgrEvo GmbH, Germany.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))