

89.06.09

weed away

Cobutox 400

EMULSIFIABLE CONCENTRATE HERBICIDE

CONTAINS 2,4-DB

For weed control in seeding - Alfalfa,  
Bird's-foot Trefoil and Pastures

AGRICULTURAL

READ THE LABEL BEFORE USING

GUARANTEE: 2,4-DB 400 g/L

REGISTRATION NO. 18230  
PEST CONTROL PRODUCTS ACT

20 LITRES

WARNING POISON

INTERPROVINCIAL COOPERATIVE LIMITED. P.O. BOX 1050,  
SASKATOON, SASKATCHEWAN

WINNIPEG TORONTO MONTREAL MONCTON

DIRECTIONS:

Rate Product/hectare

Seeding Alfalfa and Bird's-foot Trefoil (direct or underseeded)	2.75 - 3.5 L	Treat when crop is in the 1st to 4th trifoliolate leaf stage. Do not apply if crop is to be used for seed production later in the season.
Seeding Clovers (direct or Underseeded)	2.75 - 3.5 L	Treat as soon as possible after the first trifoliolate leaf appears. Do not spray sweet clovers. Red clover may be damaged.
Cereals (Wheat, oats and barley)	2.75 - 3.5 L	Apply after the crop has 5 full exposed leaves but before the early flag leaf stage.
Field Corn	2.75 - 4.25 L	Apply after crop is at least 40 cm high but before tasseling. Use drop nozzles to avoid spraying crop foliage.
Pastures containing forage legumes	2.75 - 4.25 L	Treat after cutting or grazing when growth is not above 7.5 cm in height. Damage is likely to occur if crop is sprayed beyond this stage.
Established pastures 2 years or more	2.75 - 4.25 L	Treat early in the season when weeds are in the seeding stage.

2,4-DB + MCPA TANK MIX

For improved wild mustard control in seeding alfalfa and bird's foot trefoil. When mustard is beyond the four leaf stage, spray 2,4-DB AT 2 L/ha in a tank mix with MCPA AMINE 500 at 70 mL per hectare, or equivalent MCPA Amine. Sodium Salt or Potassium Salt at 35 g active ingredient per hectare. This tank mix may increase

crop stunting. Treat mustard with 2,4-DB alone before the four leaf stage if possible.

#### ANNUALS:

Weeds controlled with 2.75 Litres of 2,4-DB. Wormseed mustard, wild mustard, ball mustard, stinkweed, ragweed, redroot pigweed, shepherd's purse, lamb's quarters. Weeds controlled with 3.5 litres of 2,4-DB, Oak-leaved goose foot, wild buckwheat, wild radish. Spray when weeds are in the 1-2 leaf stage. Weeds controlled with 4.25 litres of 2,4-DB. Lady's thumb, green smartweed (seedlings are only stunted but their competitive ability is reduced).

#### PERENNIALS:

Weeds controlled with 2.75-4.25 litres of 2,4-DB per hectare.

Weeds Spray in the stage listed below

Bull Thistle Rosette to early bud Perennial Sow Thistle Rosette  
Canada Thistle Top growth only. Spray from 15 cm to early bud  
Curled Dock Early growth Plantain Before flowering Field Bindweed  
Top growth reduced in spring. For best results, spray in late  
summer Yellow Rocket Spray between 3rd week in September and  
middle of October Chicory Rosette stage Narrow leafed Spray  
3.5 L/ha to rosette stage in the fall after Hawksbeard legume  
growth ends.

Weeds reduced by 4.25 Litres of 2,4-DB per hectare. (top growth checked only)

Dandelion Spray before bud stage Horsetail Spray when 10-12 cm high

Use lower rates for susceptible weeds in the seeding stage. Use higher rates for older weeds, heavy infestation and under poor growing conditions. Do not exceed recommended rate for each crop. Apply product in warm weather. Do not apply in drought conditions. Use at least 150-200 litres of water per hectare. Do not exceed 275 kPa of spray pressure.

#### PRECAUTIONS:

KEEP OUT OF THE REACH OF CHILDREN. Do not store near seed, feed, fertilizer, other pesticides or growing plants. Do not contaminate rivers, ponds, lakes or other bodies of water. Avoid spray drift to susceptible crops such as grapes, tomatoes, beans, turnips, cauliflower, cabbage, sugar beets and tobacco. Clean sprayer thoroughly after use. Wash hands and face after use. Change contaminated clothing. Do not freeze.

FIRST AID:

In case of accidental skin or eye contact, the affected areas should be thoroughly washed with cold water followed by a saline douche in the case of eye contact and washing with soap and water in the case of skin contact. This product contains petroleum distillates; in case of accidental swallowing, do not induce vomiting. Call a physician immediately.

DISPOSAL PROCEDURES:

Destroy empty container. Dispose of wastes by burying in a non-crop area away from all water supplies. Follow local regulations if they have been established.

NOTICE TO USER:

This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

NON GUARANTEE:

Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any

guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))