Santé

Canada

RD2009-04

Registration Decision

Gliocladium catenulatum strain J1446

(publié aussi en français)

2 April 2009

This document is published by the Health Canada Pest Management Regulatory Agency. For further information, please contact:

Publications Pest Management Regulatory Agency Health Canada 2720 Riverside Drive A.L. 6605C Ottawa, Ontario K1A 0K9

Internet: pmra_publications@hc-sc.gc.ca healthcanada.gc.ca/pmra

Facsimile: 613-736-3758

Information Service: 1-800-267-6315 or 613-736-3799 pmra infoserv@hc-sc.gc.ca



HC Pub: 8168

ISBN: 978-1-100-12222-9 (978-1-100-12223-6)

Catalogue number: H113-25/2009-4E (H113-25/2009-4E-PDF)

© Her Majesty the Queen in Right of Canada, represented by the Minister of Health Canada, 2009

All rights reserved. No part of this information (publication or product) may be reproduced or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, or stored in a retrieval system, without prior written permission of the Minister of Public Works and Government Services Canada, Ottawa, Ontario K1A 0S5.

Table of Contents

Overview
Registration Decision for <i>Gliocladium catenulatum</i> strain J1446
What Does Health Canada Consider When Making a Registration Decision?
What Is Gliocladium catenulatum strain J1446?
Health Considerations
Environmental Considerations
Value Considerations
Measures to Minimize Risk
Other Information
References

Overview

Registration Decision for Gliocladium catenulatum strain J1446

Health Canada's Pest Management Regulatory Agency (PMRA), under the authority of the <u>Pest Control Products Act</u> and in accordance with the Pest Control Products Regulations, is granting full registration for the sale and use of *Gliocladium catenulatum* J1446 Dried Cell Mass and Prestop Biofungicide WP containing the technical grade active ingredient *Gliocladium catenulatum* strain J1446 for the suppression of a variety of fungal diseases on the following greenhouse-grown vegetables, herbs and ornamentals: cucumber, tomato, pepper, lettuce, cauliflower, broccoli, oregano, basil, parsley, thyme, dill, alyssum, geranium, pansy, petunia, salvia, snapdragon, tagetes, poinsettia and saintpaulia.

Current scientific data from the applicant were evaluated to determine whether, under the proposed conditions of use, the product has value and does not present an unacceptable risk to human health or the environment.

These products were first proposed for registration in the consultation document¹: Proposed Registration Decision *Gliocladium catenulatum* strain J1446 (PRD2008-03). This Registration Decision² describes this stage of the PMRA's regulatory process for *Gliocladium catenulatum* strain J1446, and summarizes the Agency's decision and the reasons for it. Comments received were only for error correction and did not affect the risk assessment. This decision is consistent with the proposed registration decision stated in PRD2008-03.

For more details on the information presented in this Registration Decision, please refer to PRD2008-03, which contains a detailed evaluation of the information submitted in support of this registration.

What Does Health Canada Consider When Making a Registration Decision?

The key objective of the *Pest Control Products Act* is to prevent unacceptable risks to people and the environment from the use of pest control products. Health or environmental risk is considered acceptable if there is reasonable certainty that no harm to human health, future generations or the environment will result from use of or exposure to the product under its conditions of registration.³ The Act also requires that products have value⁴ when used according

[&]quot;Consultation statement" as required by subsection 28(2) of the *Pest Control Products Act*.

[&]quot;Decision statement" as required by subsection 28(5) of the *Pest Control Products Act*.

³ "Acceptable risks" as defined by subsection 2(2) of *Pest Control Products Act*.

[&]quot;Value" as defined by subsection 2(1) of *Pest Control Products Act*"...the product's actual or potential contribution to pest management, taking into account its conditions or proposed conditions of registration, and includes the product's (a) efficacy; (b) effect on host organisms in connection with which it is intended to be used; and (c) health, safety, environmental benefits and social, economic impact".

to the label directions. Conditions of registration may include special precautionary measures on the product label to further reduce risk.

To reach its decisions, the PMRA applies modern, rigorous risk-assessment methods and policies. These methods consider the unique characteristics of sensitive subpopulations in humans (e.g. children), as well as organisms in the environment (e.g. those most sensitive to environmental contaminants). These methods and policies also consider the nature of the effects observed and the uncertainties present when predicting the impact of pesticides. For more information on how the PMRA regulates pesticides, as well as on the assessment process and risk-reduction programs, please visit the PMRA's website at www.healthcanada.gc.ca/pmra.

What Is Gliocladium catenulatum strain J1446?

Gliocladium catenulatum is a fungus that grows on dead organic matter that can be found commonly in soil worldwide. G. catenulatum strain J1446 was originally isolated as a microbial pest control agent (MPCA) due to its ability to suppress soil-borne fungal diseases on plants.

Health Considerations

Can Approved Uses of Gliocladium catenulatum strain J1446 Affect Human Health?

Gliocladium catenulatum strain J1446 is unlikely to affect your health when Prestop Biofungicide WP is used according to label directions.

Exposure to *G. catenulatum* strain J1446 may occur during handling and application of Prestop Biofungicide WP. When assessing health risks, several key factors are considered: the microorganism's biological properties (e.g. production of toxic byproducts); reports of any adverse incidents; its potential to cause disease or toxicity as determined in toxicological studies and the levels to which people may be exposed, relative to exposures already encountered in nature to other isolates of the microorganism.

Toxicology studies in laboratory animals describe potential health effects from large doses in an effort to identify any potential to cause disease or toxicity. When *G. catenulatum* strain J1446 was tested on laboratory animals, there were no signs that it caused any significant toxicity or disease.

Residues in Water and Food

Dietary risks from food and water are not of concern.

The Food and Drugs Act prohibits the sale of adulterated food, that is, food containing a pesticide residue that exceeds the established maximum residue limit (MRL). Pesticide MRLs are established for *Food and Drugs Act* purposes through the evaluation of scientific data under the Pest Control Products Act. Each MRL value defines the maximum concentration in parts per million (ppm) of a pesticide allowed in or on certain foods. Food containing a pesticide residue that does not exceed the established MRL does not pose an unacceptable health risk. Strains of G. catenulatum are common in nature and the use of Prestop Biofungicide WP in greenhouses to suppress fungal plant disease on vegetables, herbs and ornamentals is not expected to significantly increase the natural environmental background levels of this microorganism. Furthermore, when G. catenulatum strain J1446 was administered orally to rats, no signs that it caused toxicity or disease were observed. Although secondary metabolites of toxicological significance have been shown to be produced by other isolates of G. catenulatum, due to the demonstrated low toxicity and absence of such metabolites in cultures of G. catenulatum strain J1446, the risks from secondary metabolites for the general population, including infants and children, are negligible. The establishment of a maximum residue limit (MRL) is therefore not required for G. catenulatum strain J1446. As well, the likelihood of residues of G. catenulatum strain J1446 contaminating drinking water supplies is negligible to nonexistent. Consequently, dietary exposure and risk are minimal to nonexistent.

Occupational Risks From Handling Prestop Biofungicide WP

Occupational risks are not of concern when Prestop Biofungicide WP is used according to label directions, which include protective measures.

Growers handling Prestop Biofungicide WP can come into direct contact with *G. catenulatum* strain J1446 on the skin, in the eyes or by inhalation. For this reason, the label specifies that growers exposed to Prestop Biofungicide WP during handling, mixing/loading, application or clean-up/repair activities must wear waterproof gloves, a long-sleeved shirt, long pants, eye goggles, shoes, socks and a dust/mist filtering respirator (MSH/NIOSH approval number prefix TC-21C) or a NIOSH-approved respirator with any –95, R-95, P-95 or HE filter. Furthermore, early-entry workers will be restricted from entering areas where Prestop Biofungicide WP has been applied as a foliar spray for a period of four hours unless wearing the indicated personal protective equipment with the exception of eye goggles and a dust/mist filtering respirator, which are required only until the spray mist has settled.

Bystander exposure is expected to be much less than that of applicators, handlers and mixers/loaders and is considered negligible. Therefore, health risks to bystanders are not of concern.

Environmental Considerations

What Happens When Prestop Biofungicide WP is Introduced Into the Environment?

Environmental risks are not of concern.

Studies designed to examine the effects of *Gliocladium catenulatum* strain J1446 on various non-target organisms were evaluated. Few adverse effects were observed in birds, freshwater fish, terrestrial arthropods (including honeybees), aquatic invertebrates, marine animals or algae.

G. catenulatum is not generally considered to be a disease-causing agent. Therefore, Prestop Biofungicide WP is expected to present a negligible risk to non-target organisms.

Value Considerations

What Is the Value of Prestop Biofungicide WP?

Prestop Biofungicide WP suppresses specific soil and seed-borne pathogens and foliar diseases on greenhouse-grown vegetables, herbs and ornamentals.

The formulated end-use product, Prestop Biofungicide WP, is a biological fungicide. When applied in a 0.5–1.0% solution as a soil media treatment, a soil drench treatment or a foliar treatment, it will suppress specific soil and seed-borne pathogens and foliar diseases on greenhouse-grown vegetables, herbs and ornamentals. It is best used as a preventative before disease is present, and reapplication is necessary every three to six weeks depending on the disease pressures in the greenhouse and the method of application.

Prestop Biofungicide WP is considered to be a low-risk product and can be used as an integral part of an integrated pest management program to reduce the reliance on chemical alternatives. There are limited fungicide products currently available to greenhouse growers; therefore, Prestop Biofungicide WP is a new product available to the greenhouse sector. It may also be used as a resistance management tool for rotation of fungicides where no other alternative products are available.

Measures to Minimize Risk

The labels of registered pesticide products include specific instructions for use. Directions include risk-reduction measures to protect human and environmental health. These directions must be followed by law.

The key risk-reduction measures on the label of Prestop Biofungicide WP to address the potential risks identified in this assessment are as follows.

Key Risk-Reduction Measures

Human Health

Because of concerns with users developing allergic reactions through repeated high exposures to *G. catenulatum* strain J1446, anyone handling, mixing/loading, applying or involved in clean-up/repair activities of Prestop Biofungicide WP must wear waterproof gloves, a long-sleeved shirt, long pants, eye goggles and a dust/mist filtering respirator (MSH/NIOSH approval number prefix TC-21C) or a NIOSH-approved respirator with any –95, R-95, P-95 or HE filter. Furthermore, early-entry workers will be restricted from entering areas where Prestop Biofungicide WP has been applied as a foliar spray for a period of four hours unless wearing the indicated personal protective equipment with the exception of eye goggles and a dust/mist filtering respirator, which are required only until the spray mist has settled.

• Environment

As a general precaution, handlers are asked not to contaminate irrigation or drinking water or aquatic habitats through equipment cleaning or waste disposal. In addition, growers must not allow effluent or runoff from greenhouses containing this product to enter lakes, streams, ponds or other water bodies.

Other Information

The relevant test data on which the decision is based (as referenced in this document) are available for public inspection, upon application, in the PMRA's Reading Room (located in Ottawa). For more information, please contact the PMRA's Pest Management Information Service.

Any person may file a notice of objection⁵ regarding this registration decision within 60 days from the date of publication of this Registration Decision. For more information regarding the basis for objecting (which must be based on scientific grounds), please refer to the PMRA's website, Requesting a Reconsideration of Decision, or contact the PMRA's Pest Management Information Service.

.

As per subsection 35(1) of the *Pest Control Products Act*.

References

A. LIST OF STUDIES/INFORMATION SUBMITTED BY REGISTRANT

1.0 The Active Ingredient, Its Properties and Uses

PMRA 1187813	1997, History of the strain Gliocladium catenulatum J1446., DACO: M1.2, M2.7.1
PMRA 1187814	Gliocladium Corda 1840., DACO: M1.2, M2.7.1
PMRA 1187815	1996, Report of identification of the strain J1446., det 225 (1996), DACO: M1.2, M2.7.1
PMRA 1187816	2000, Identification of the strain J1446., DACO: M1.2, M2.7.1
PMRA 1187817	2000, Description of Gliocladium catenulatum J1446, DACO: M1.2, M2.7.1
PMRA 1187818	Paavanen-Huhtala, S. et al, 1999, Development of strain-specific primers for a strain of Gliocladium catenulatum used in biological control., European Journal of Plant Pathology 106: 187-198, 2000., DACO: M1.2, M2.7.1
PMRA 1187819	2000, Identification of J1446 strain by RAPD-PCR, UP-PCR and by specific primers., DACO: M1.2, M2.7.1
PMRA 1187820	A discussion on the relationship of Gliocladium catenulatum J1446 to known pathogenic / toxigenic strains or species., 152-1., DACO: M1.2, M2.7.2
PMRA 1187821	1996, The role of antibiosis, mycoparasitism and lytic enzymes in the control efficacy of Gliocladium catenulatum strain J1446., DACO: M1.2, M2.7.2

2.0 Methods of Analysis

PMRA 1187764	2003, Material safety data sheet. Prestop WP, DACO: 0.9,M2.9.1
PMRA 1187932	2006, Prestop WP. Comprehensive Data Summary, DACO: M12.7

PMRA 1187933	Registrant's name and address, DACO: M2.1
PMRA 1187934	Name and address of manufacturing plant, DACO: M2.2
PMRA 1187935	Name and address of formulating plant (Prestop (EP), DACO: M2.3
PMRA 1187936	Trade name: Prestop, DACO: M2.4
PMRA 1187937	Binominal name (MPCA), DACO: M2.5
PMRA 1187938	Canadian patent status information, DACO: M2.6
PMRA 1187939	2006, Gliocladium catenulatum J1446: Summary of the characterization of the MPCA., DACO: M2.7.1, M2.7.2
PMRA 1187941	2003, Summary of available information on the taxonomy of Gliocladium catenulatum J1446., DACO: M2.7.1
PMRA 1187942	2004, Maintenance and subculturing of microbial strain used in Gliocladium catenulatum - formulations., SOP-137-1, DACO: M2.7.1
PMRA 1187943	1996, Identification of J1446 strain by RAPD-PCR, UP-PCR and by specific primers., DACO: M2.7.1
PMRA 1187944	1999, Current status of taxonomy of Trichoderma and Gliocladium. 6th International Trichoderma and Gliocladium Workshop, Espoo, Finland. 11-13.6 1999., DACO: M2.7.1
PMRA 1187945	1994, Identification of the microorganism. Accession number given by the International Depositary Authority: DSM 9212. Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purpose of Patent Procedure., DACO: M2.7.1
PMRA 1187946	Yli-Mattila, T. et al, 1997, Molecular identification of Gliocladium and Nectria isolates in Finland. Fifth European Fusarium Seminar, Szeged, Hungary Cereal Research Communications, Vol. 25 No. 3/1, pp. 293-297., Fifth European Fusarium Seminar, DACO: M2
PMRA 1187947	Park, J. et al, 1999, Phylogenetic analysis of the genus Gliocladium and its related taxa by comparing the sequences of internal transcribed spacers and 5.8S r-DNA., The Korean Journal of Mycology, Vol. 27 No 3, pp. 191-197., DACO: M2.7.1
PMRA 1187948	1996, Production of chitinase (chitobiosidase) by Gliocladium strain J1446., DACO: M2.7.2

PMRA 1187949	1996, Production of β-1,3-glucanase (laminarinase) by Gliocladium strain J1446., DACO: M2.7.2
PMRA 1187950	1995, Study of the minimum and maximum temperatures for the growth of Gliocladium catenulatum strain J1446., DACO: M2.7.2
PMRA 1187951	1997, Effect of temperature on the control efficacy of Gliocladium catenulatum J1446., kas 12-94, DACO: M2.7.2
PMRA 1187952	1997, Effect of pH on the control efficacy of Gliocladium catenulatum J1446: cauliflower/Rhizoctonia., Rhil14-94, DACO: M2.7.2
PMRA 1187954	1997, Gliocladium catenulatum J1446: cauliflower/Alternaria I., 3-93, DACO: M2.7.2
PMRA 1187955	1997, Effect of pH on the control efficacy of Gliocladium catenulatum J1446: cauliflower/Alternaria II., 5-93., DACO: M2.7.2
PMRA 1187956	1998, The persistence of Gliocladium catenulatum J1446 on leaf surface., 16-98, 24-98 and 25-98, DACO: M2.7.2
PMRA 1187957	1998, Survival of Gliocladium catenulatum J1446 on cucumber leaves., 060698JO, DACO: M2.7.2
PMRA 1187958	2004, Literature search Gliocladium (s) (phytotox? or pathog?)., 060698JO, DACO: M2.7.2
PMRA 1187959	2006, Mycology of dermatophyte infections. Dermnet NZ. http://dermnetnz.org/fungak/mycology.html, DACO: M2.7.2
PMRA 1187960	Classification based on 9th edition of the Dictionary of the Fungi. CABI Bioscience Databases. http://www.indexfungorum.org/Names/fundic.asp, DACO: M2.7.2
PMRA 1187961	2004, Summary of the literature searches on the production of gliotoxin and/or genotoxins by Gliocladium catenulatum., DACO: M2.7.2
PMRA 1187962	2000, Summary of the literature search on production of toxins/metabolites by Gliocladium catenulatum., DACO: M2.7.2
PMRA 1187963	2001, Analysis of gliotoxins in culture broth., Bel 78/2001, DACO: M2.7.2
PMRA 1187964	2001, Analysis of gliotoxins in unformulated cell mass., Bel 103/2001, DACO: M2.7.2

PMRA 1187965	2001, Analysis of gliotoxins in mineral wool cultivation pot samples., Bel 184/2001, DACO: M2.7.2
	104/2001, DACO. W12.7.2
PMRA 1187966	2000, Growing of Gliocladium treated cucumber seedlings in rockwool., 29-00, DACO: M2.7.2
PMRA 1187967	1995, Cytotoxicity test on Gliocladium catenulatum J1446 product 53/94., 1608/TT., DACO: M2.7.2
PMRA 1187969	1996, Cytotoxicity test on Gliocladium catenulatum J1446 cell mass batches 37/95 and 44/95., 1661/TT., DACO: M2.7.2
PMRA 1187970	1996, Cytotoxicity test on Gliocladium catenulatum J1446 formulation batch 41/95., DACO: M2.7.2
PMRA 1187971	Pasanen, A. et al, 1992, Laboratory experiments on membrane filter sampling of airborne mycotoxins produced by Stachybotrys atra Corda. Atmospheric Environment, Vol. 27A, No. 1: 9-13., Atmosphere Environment Vol. 22A, No. 1 pp. 9-12, 1993, DACO: M2.7.2
PMRA 1187972	2000, Gliocladium catenulatum J1446/ cell toxicity test., DACO: M2.7.2
PMRA 1187973	2003, Summary of available information on the secondary metabolite profile of Gliocladium catenulatum J1446., DACO: M2.7.2
PMRA 1187974	2000, Medical certificate., DACO: M2.7.2
PMRA 1187975	2005, Medical certificate., DACO: M2.7.2
PMRA 1187976	2006, Gliocladium catenulatum J1446: Summary of the Dacos M2.8-M2.12., DACO: M2.10, M2.11, M2.12, M2.8, M2.9
PMRA 1187977	2004, Quality control of Prestop WP production., SOP- 105-2., DACO: M2.8
PMRA 1187978	2004, Production of Prestop WP., SOP-106-2., DACO: M2.8
PMRA 1187979	2004, Production of Gliocladium catenulatum Technical Grade Active Ingredient (TGAI)., SOP- 136-1, DACO: M2.8
PMRA 1187980	2004, Product Specification Prestop, DACO: M2.9.1, M2.9.2
PMRA 1187981	2004, Five batch analysis of Prestop., REP-102-2, DACO: M2.9.2
PMRA 1187983	2004, Correlation between the viability and the control efficacy of Prestop WP., Based on kas-40-01, DACO: M2.9.2

PMRA 1187984	2005, Discussion of formation of unintentional ingredients Prestop WP - Gliocladium catenulatum, biofungicide sprayable formulation., REP-104-2, DACO: M2.9.3
PMRA 1187985	1998, The isolation of Gliocladium catenulatum J1446 from roots and growth substrates., REP-104-2, DACO: M2.10.1
PMRA 1187986	2004, Viability determination of biological pesticides., DACO: M2.10.1
PMRA 1187987	2004, Analysis of pathogenic contaminants in Prestop WP., REP-130-1, DACO: M2.10.2
PMRA 1496779	2006, Analyysitodistus (Analysis Certificate), 2006-00946-001, MRID: n.a., DACO: M2.10.2
PMRA 1496780	2006, Analyysitodistus (Analysis certificate), 2006-00945-001, MRID: n.a., DACO: M2.10.2
PMRA 1187988	2005, Storage stability of Prestop WP., REP-108-3, DACO: M2.11
PMRA 1187989	2006, Tap density of Prestop WP., REP-168-1, DACO: M2.12
PMRA 1187990	2006, Gliocladium catenulatum biofungicide (Prestop WP): Determination of long-term storage stability and physico-chemical characteristics., 1841/0009, DACO: M2.11
PMRA 1188054	2004, Summary of the literature searches on the phytotoxicity and/or pathogenicity of Gliocladium catenulatum., DACO: M2.7.2,M9.8.1
PMRA 1188055	2000, Summary of the literature search on Gliocladium catenulatum regarding pathogenicity or other interactions with plants, DACO: M2.7.2,M9.8.1
PMRA 1465323	2005, Quality control in Prestop production facilities, SOP-3.2, MRID: N/A, DACO: M2.9.3
PMRA 1465324	1999, Isolation and structure determination of TMC-151s: Novel polyketide antibiotics from Gliocladium catenulatum Gilman & Abbot TC 1280, N/A, MRID: N/A, DACO: M2.7.2
PMRA 1465325	1993, Anthrotainin, an inhibitor of substance P binding produced by Gliocladium catenulatum, N/A, MRID: N/A, DACO: M2.7.2
PMRA 1465326	2007, Storage stability of Prestop WP, 108-4, MRID: N/A, DACO: M2.11
PMRA 1474679	2006, Specification of Prestop Mix, RE-179-1, DACO: M2.9.1

PMRA 1188046 2001, Discussion on analysis of gliotoxin in mineral wool; culture broth and unformulated cell mass samples., DACO: M9.7

3.0 Impact on Human and Animal Health

PMRA 1187767	U.S. EPA, Registration Eligibility Decision Gliocladium catenulatum strain J1446 (PC Code 021009)., DACO: 12.5.4
PMRA 1187768	U.S. EPA, US EPA Pesticide Fact Sheet: Gliocladium catenulatum strain J1446, DACO: 12.5.4
PMRA 1187822	1998, The spread and persistence of Gliocladium catenulatum J1446 in the air. Part I., 20/98, DACO: M1.2,M5.0,M8.3
PMRA 1187823	1998, The spread and persistence of Gliocladium catenulatum J1446 in the air. Part II., 24/98 and 26/98, DACO: M1.2,M5.0,M8.3
PMRA 1187824	1998, The spread and persistence of Gliocladium catenulatum J1446 in the air. Part III., 28/98, DACO: M1.2,M5.0,M8.3
PMRA 1187991	2005, Summary of Human Health and Safety Testing., DACO: M4.1
PMRA 1187992	2006, Gliocladium catenulatum J1446: Summary of the infectivity and toxicity studies., DACO: M4.2.1
PMRA 1187993	1996, Gliocladium catenulatum cell mass: Acute oral toxicity and pathogenicity to the rat., RKY 135, DACO: M4.2.2
PMRA 1187994	1996, Gliocladium (J1446) preparation: Acute oral toxicity (limit) test in rats., 556576; 10919, DACO: M4.2.2
PMRA 1187996	1996, Gliocladium catenulatum cell mass: Acute pulmonary toxicity and pathogenicity to rat., RKY 137, DACO: M4.2.3
PMRA 1187997	1996, Gliocladium (J1446) preparation: Acute inhalation toxicity study in rats., 654570;10916, DACO: M4.2.3
PMRA 1187998	2006, Gliocladium catenulatum J1446: Summary of the acute infectivity study (IP), DACO: M4.3.1
PMRA 1187999	1996, Gliocladium catenulatum cell mass: Acute intra-peritoneal toxicity and pathogenicity to the rat, RKY 136, DACO: M4.3.3
PMRA 1188000	2004, Data waiver for the acute dermal toxicity study on Prestop., DACO: M4.4

PMRA 1188001	1996, Gliocladium (J1446) preparation: Acute dermal toxicity (limit) test in rats., 556576;10920, DACO: M4.4
PMRA 1188002	2002, Safety Data Sheet. Valio Skimmed Milk Powder Medium Heat, DACO: 0.9.1,M2.9.1,M4.4
PMRA 1188003	1999, Material safety data sheet. C-Sorbidex S 16603, DACO: 0.9.1,M2.9.1,M4.4
PMRA 1188004	2006, Gliocladium catenulatum J1446: Summary of the dermal irritation study., DACO: M4.5.1
PMRA 1188005	2004, Data waiver for a dermal irritation study on Prestop., DACO: M4.5.2
PMRA 1188006	1996, Gliocladium (J1446) preparation: Acute dermal irritation test in rabbits., 556576;10921, DACO: M4.5.2
PMRA 1188007	2006, Gliocladium catenulatum J1446: Summary of the skin sensitization study, DACO: M4.6
PMRA 1188009	2004, Prestop Mix: Gliocladium catenulatum J1446 biofungicide hypersensitivity reporting., DACO: M4.6
PMRA 1188011	2004, Prestop WP: Gliocladium catenulatum J1446 biofungicide hypersensitivity reporting., DACO: M4.6
PMRA 1188012	1996, Gliocladium (J1446) preparation: Buehler sensitisation test in guinea pigs., 556576;10923, DACO: M4.6
PMRA 1188013	1999, Gliocladium J1446 biopesticide: Acute eye irritation test in the rabbit., 245/187, DACO: M4.9
PMRA 1188014	1996, Gliocladium (J1446) preparation: Acute eye irritation test in rabbits., 556576;10922, DACO: M4.9
PMRA 1188015	2004, Assessment of operator, bystander and worker exposure to Prestop Mix (Gliocladium catenulatum)., DACO: M5.0
PMRA 1188016	1997, Viable residues of Gliocladium catenulatum on cucumber fruits., 11/97, DACO: M7.0
PMRA 1188017	2006, Data waiver for the food and feed residue studies., DACO: M7.0
PMRA 1188018	1997, Viable residues of Gliocladium catenulatum J1446 on cucumber fruits one day after the treatment., 11/97, DACO: M7.0

PMRA 1187967	1995, Cytotoxicity test on Gliocladium catenulatum J1446 product 53/94., 1608/TT., DACO: M2.7.2
PMRA 1187969	1996, Cytotoxicity test on Gliocladium catenulatum J1446 cell mass batches 37/95 and 44/95., 1661/TT., DACO: M2.7.2
PMRA 1187970	1996, Cytotoxicity test on Gliocladium catenulatum J1446 formulation batch 41/95., DACO: M2.7.2
PMRA 1187971	Pasanen, A. et al, 1992, Laboratory experiments on membrane filter sampling of airborne mycotoxins produced by Stachybotrys atra Corda. Atmospheric Environment, Vol. 27A, No. 1: 9-13., Atmosphere Environment Vol. 22A, No. 1 pp. 9-12, 1993, DACO: M2.7.2

4.0 Impact on the Environment

4.0 Impa	ect on the Environment
PMRA 1187948	1996, Production of chitinase (chitobiosidase) by Gliocladium strain J1446., DACO: M2.7.2
PMRA 1187949	1996, Production of -1,3-glucanase (laminarinase) by Gliocladium strain J1446., DACO: M2.7.2
PMRA 1187950	1995, Study of the minimum and maximum temperatures for the growth of Gliocladium catenulatum strain J1446., DACO: M2.7.2
PMRA 1187966	2000, Growing of Gliocladium treated cucumber seedlings in rockwool., 29-00, DACO: M2.7.2
PMRA 1188020	2005, Summary on the environmental fate of Gliocladium catenulatum J1446, DACO: M8.1
PMRA 1188021	1997, The survival of Gliocladium catenulatum J1446 in water., Lab-48-96, DACO: M8.2.2
PMRA 1188022	1998, Survival of Gliocladium catenulatum J1446 on cucumber leaves., 060698Jo., DACO: M8.3
PMRA 1188023	1998, The persistence of Gliocladium catenulatum J1446 on leaf surface., 16-98, 24-98 and 25-98., DACO: M8.3
PMRA 1188024	1996, Confidential report on the survival of J1446 in a commercial peat-based growing medium for Kemira Agro Oy., DACO: M8.3
PMRA 1188025	2001, Possible mobility of the conidia into the soil., DACO: M8.5
PMRA 1188026	2006, Gliocladium catenulatum J1446 Summary of the environmental toxicity studies., DACO: M9.1

PMRA 1188027	1997, Gliocladium catenulatum cell mass: an avian oral pathogenicity and toxicity study in the northern bobwhite., 293-109, DACO: M9.2.1
PMRA 1188028	2004, Response to PMRA data evaluation for Gliocladium catenulatum cell mass (technical product): a five-concentration toxicity and pathogenicity test with the rainbow trout (Oncorhynchus mykiss)., 293A-104, DACO: M9.4.1
PMRA 1188029	1998, Gliocladium catenulatum cell mass (technical product): a five-concentration toxicity and pathogenicity test with the rainbow trout (Oncorhynchus mykiss)., 293A-104, DACO: M9.4.1
PMRA 1188031	1998, Gliocladium catenulatum cell mass (technical product): a dietary pathogenicity and toxicity study with the honey bee (Apis mellifera)., 293-110, DACO: M9.5.1
PMRA 1188032	1998, Gliocladium catenulatum cell mass (technical product): a dietary pathogenicity and toxicity study with the ladybird beetle (Hippodamia convergens)., 293-111, DACO: M9.5.1
PMRA 1188033	1998, Gliocladium catenulatum cell mass (technical product): a dietary pathogenicity and toxicity study with green lacewing larvae (Chrysoperla carnea)., 293-113A, DACO: M9.5.1
PMRA 1188034	1998, Gliocladium catenulatum cell mass (technical product): a dietary pathogenicity and toxicity study with parasitic hymenoptera (Nasonia vitripennis)., 293-112, DACO: M9.5.1
PMRA 1188035	1998, Gliocladium catenulatum cell mass (technical product): a 21-day life-cycle toxicity and pathogenicity test with the cladoceran (Daphnia magna)., 293A-103, DACO: M9.5.2
PMRA 1188036	1997, Gliocladium catenulatum cell mass: a pathogenicity study with the earthworm in an artificial soil substrate., 293A-105, DACO: M9.6
PMRA 1188037	1997, Effect of Gliocladium catenulatum J1446 biofungicide powder on other soil microbes., DACO: M9.7
PMRA 1188038	1998, Effect of Gliocladium catenulatum J1446 incorporation on peat microflora., 061797JO, DACO: M9.7
PMRA 1188039	2001, Analysis of gliotoxin in culture broth., BEL78/2001, DACO: M9.7
PMRA 1188040	2001, Analysis of gliotoxin in unformulated cell mass, BEL103/2001, DACO: M9.7

PMRA 1188042	2001, Analysis of gliotoxin in mineral wool cultivation in pot samples., BEL184/2001, DACO: M9.7	
PMRA 1188043	1996, The role of antibiosis, mycoparasitism and lytic enzymes in the control efficacy of Gliocladium catenulatum strain 1446., DACO: M9.7	
PMRA 1188044	2003, Summary of available information on the secondary metabolite profile of Gliocladium catenulatum J1446, DACO: M9.7	
PMRA 1188046	2001, Discussion on analysis of gliotoxin in mineral wool; culture broth and unformulated cell mass samples., DACO: M9.7	
PMRA 1188047	Bioassay of gliotoxin in Gliocladium extracts based on Escherichia coli CM871., DACO: M9.7	
PMRA 1188049	2004, Literature searches conducted in 2000 and 2004 on the pathogenicity or phytotoxicity of Gliocladium catenulatum., DACO: M9.8.1	
PMRA 1188050	2005, Data waiver for terrestrial plant toxicity/pathogenicity study., DACO: M9.8.1	
PMRA 1188052	1997, Effect of high dosages of Gliocladium catenulatum J1446 on different crops., DACO: M9.8.1	
PMRA 1188054	2004, Summary of the literature searches on the phytotoxicity and/or pathogenicity of Gliocladium catenulatum., DACO: M2.7.2,M9.8.1	
PMRA 1188055	2000, Summary of the literature search on Gliocladium catenulatum regarding pathogenicity or other interactions with plants, DACO: M2.7.2,M9.8.1	
PMRA 1188056	2005, Data waiver for the aquatic plant infectivity study., DACO: M9.8.2	
PMRA 1188057	2001, Gliocladium catenulatum cell mass (technical product): Algal inhibition test., 245/205, DACO: M9.8.2	
5.0 Value		
PMRA 1187842	1998. The effect of Gliocladium catenulatum J1446 on grey mold (Botrytis cinerea) on cucumber leaves. DACO M10.2.1	
PMRA 1187843	1998. The effect of Gliocladium catenulatum J1446 on grey mold (Botrytis cinerea) on pepper leaves. DACO M10.2.1	
PMRA 1187844	1998. The effect of Gliocladium catenulatum J1446 fungal suspension on grey mold (Botrytis cinerea) on pepper leaves. DACO M10.2.1	

PMRA 1187845	1998. Control of bottom rot on lettuce caused by Rhizoctonia solani. DACO M10.2.1
PMRA 1187846	2002. Biological control agents and composts suppress Fusarium and Pythium root rots on greenhouse cucumbers. DACO M10.2.1, M10.2.2
PMRA 1187847	2003. Biological control of damping-off and root rot caused by Pythium aphanidermatum on greenhouse cucumbers. DACO M10.2.1, M10.2.2
PMRA 1187848	2003. Efficacy of biological and chemical treatments for control of Fusarium root and stem rot on greenhouse cucumber. DACO M10.2.1, M10.2.2
PMRA 1187849	2006. Efficacy of Prestop WP (Gliocladium catenulatum J1446) for the control of Fusarium solani in greenhouse grown peppers. DACO M10.2.2
PMRA 1187851	1998. The effect of Gliocladium catenulatum J1446 on Botrytis cinerea on tomato plants. DACO M10.2.2
PMRA 1187852	1998. The effect of Gliocladium catenulatum J1446 fungal suspension on grey mold (Botrytis cinerea) on pepper plants. DACO M10.2.2
PMRA 1187853	1998. Gliocladium catenulatum J1446 on the cuttings of Fuchsia. DACO M10.2.2
PMRA 1187854	1998. The effect of Gliocladium catenulatum J1446 on grey mold (Botrytis cinerea) on tomato plants. DACO M10.2.1
PMRA 1187855	1997. The effect of Gliocladium catenulatum J1446 on grey mold (Botrytis cinerea) on cucumber leaves. DACO M10.2.1
PMRA 1187856	1994. The effect of Gliocladium catenulatum J1446 biofungicide powder on Pythium damping-off in cucumber. DACO M10.2.2
PMRA 1187857	1994. The effect of Gliocladium catenulatum J1446 biofungicide powder on Pythium damping-off in cucumber. DACO M10.2.2
PMRA 1187858	1994. The effect of Gliocladium catenulatum J1446 biofungicide powder on Pythium damping-off in cucumber. DACO M10.2.2
PMRA 1187859	1996. The effect of Gliocladium catenulatum J1446 biofungicide powder on Pythium in lettuce. DACO M10.2.2
PMRA 1187860	1996. The effect of Gliocladium catenulatum J1446 biofungicide powder on Pythium in lettuce. DACO M10.2.2

PMRA 1187861	1995. The effect of Gliocladium catenulatum J1446 biofungicide powder on Pythium in lettuce. DACO M10.2.2
PMRA 1187862	1995. The effect of Gliocladium catenulatum J1446 biofungicide powder on Pythium in lettuce. DACO M10.2.2
PMRA 1187863	1995. The effect of Gliocladium catenulatum J1446 biofungicide powder on Pythium in lettuce. DACO M10.2.2
PMRA 1187864	1996. The effect of Gliocladium catenulatum J1446 biofungicide powder on Pythium damping-off in oregano. DACO M10.2.2
PMRA 1187865	1996. The effect of Gliocladium catenulatum J1446 biofungicide powder on Pythium damping-off in basilica. DACO M10.2.2
PMRA 1187867	1996. The effect of Gliocladium catenulatum J1446 biofungicide powder on Pythium damping-off in parsley. DACO M10.2.2
PMRA 1187868	1996. The effect of Gliocladium catenulatum J1446 biofungicide powder on Pythium damping-off in thyme. DACO M10.2.2
PMRA 1187869	1995. The effect of Gliocladium catenulatum J1446 biofungicide powder on Pythium damping-off in oregano. DACO M10.2.2
PMRA 1187870	1997. The effect of Gliocladium catenulatum J1446 biofungicide powder on Rhizoctonia damping-off in cauliflower. DACO M10.2.2
PMRA 1187871	1994. The effect of Gliocladium catenulatum J1446 biofungicide powder on Rhizoctonia damping-off in cauliflower. DACO M10.2.2
PMRA 1187872	1996. The effect of Gliocladium catenulatum J1446 biofungicide powder on Rhizoctonia damping-off in cauliflower. DACO M10.2.2
PMRA 1187873	1998. The effect of Gliocladium catenulatum J1446 fungal suspension on grey mold (Botrytis cinerea) on pepper plants. DACO M10.2.2
PMRA 1187874	1998. Gliocladium catenulatum J1446 on the cuttings of Fuchsia. DACO M10.2.2
PMRA 1187875	Use of Prestop mix for control of Pythium root rot in poinsettia. DACO M10.2.2
PMRA 1187876	2000. Microbial products for prevention of poinsettia root rot diseases. IR-4 report. DACO M10.2.2
PMRA 1187877	Prestop and Fusarium crown and root rot of cucumber. DACO M10.2.2

PMRA 1187878	Control of bottom rot of lettuce caused by Rhizoctonia solani. DACO M10.2.1
PMRA 1187879	1998. The effect of Gliocladium catenulatum J1446 on Botrytis cinerea on tomato plants. DACO M10.2.2
PMRA 1187880	1998. The effect of Gliocladium catenulatum J1446 fungal suspension on grey mold (Botrytis cinerea) on pepper plants. DACO M10.2.2
PMRA 1187881	1997. The effect of Gliocladium catenulatum J1446 against damping-off of cucumber on rockwool. DACO M10.2.2
PMRA 1187883	1996. The effect of Gliocladium catenulatum J1446 against damping-off of cucumber on rockwool. DACO M10.2.2
PMRA 1187885	1997. The effect of Gliocladium catenulatum J1446 on Pythium ultimum with normal and sub irrigation. DACO M10.2.2
PMRA 1187886	1997. The use of Gliocladium catenulatum J1446 biofungicide powder in the control of Phytophthora in poinsettia. DACO M10.2.2
PMRA 1187887	1998. Control of Didymella by Gliocladium catenulatum J1446 in commercial cultivation. DACO M10.2.2
PMRA 1187888	1998. The effect of Gliocladium catenulatum J1446 on Didymella of cucumber. DACO M10.2.2
PMRA 1187889	1998. The effect of Gliocladium catenulatum J1446 on Didymella of cucumber. DACO M10.2.2
PMRA 1187890	1998. The effect of Gliocladium catenulatum J1446 on Didymella of cucumber. DACO M10.2.2
PMRA 1187891	1995. Rhizoctonia damping-off of alyssum and salvia. Pythium damping-off of pansy, tomato and snapdragon. Confidential report on the evaluation of J1446 for Kemira Agro Oy. DACO M10.2.2
PMRA 1187892	1996. Phytophthora root and basal stem rot of petunia and tagetes. DACO M10.2.2
PMRA 1187893	1996. Phytophthora root and basal stem rot of petunia. DACO M10.2.2
PMRA 1187894	2001. Control of soil borne diseases of greenhouse tomato. DACO M10.2.2
PMRA 1187895	2002. Trial against Pythium ultimum on basil (cv Genovese gigante). DACO M10.2.2

PMRA 1187896	2002. Trial against Fusarium oxysporum f. sp. basilici on basil (cv Genovese gigante). DACO M10.2.2
PMRA 1187898	2002. Trial against Fusarium oxysporum f. sp. basilici on basil (cv Fine verde). DACO M10.2.2
PMRA 1187899	2005. Comparing the efficacy of the two Prestop formulations: Prestop and Prestop mix. DACO M10.2.2
PMRA 1187900	2006. The effect of Prestop WP biofungicide on basil artificially inoculated with Pythium. DACO M10.2.2
PMRA 1187901	2004. The effect of Prestop WP biofungicide on saintpaulia artificially inoculated with Pythium. DACO M10.2.2
PMRA 1187902	2004. The effect of Prestop WP biofungicide on geranium artificially inoculated with Pythium. DACO M10.2.2
PMRA 1187903	2003. Mycostop and Prestop WP on sweet pepper in Finland. DACO M10.2.2
PMRA 1187904	1998. Control of Didymella by Gliocladium catenulatum J1446 in commercial cultivation. DACO M10.2.2
PMRA 1187905	1998. The effect of Gliocladium catenulatum J1446 on Didymella of cucumber. DACO M10.2.2
PMRA 1187906	1998. The effect of Gliocladium catenulatum J1446 on Didymella of cucumber. DACO M10.2.2
PMRA 1187907	2002. Botrytis stem canker of greenhouse tomato managing the disease through cultural, biological and chemical controls. DACO M10.2.2
PMRA 1187908	Evaluation of Prestop for control of gummy stem blight on naturally infected cucumber plants in a commercial greenhouse. DACO M10.2.2
PMRA 1187909	2002. Evaluation of biological and chemical treatments for control of gummy stem blight on cucumber plants grown hydroponically in greenhouses. Pacific Agrifood Research Centre, Agriculture and Agrifood Canada Biocontrol 49:109-117. DACO M10.2.2
PMRA 1187910	2001. Biological control of stem canker of greenhouse tomatoes caused by Botrytis cinerea Can. J. Microbiol. 48:550-554. DACO M10.2.2
PMRA 1187911	2005. Effect of Gliocladium catenulatum J1446 on lettuce at a high dose. DACO M10.3.1

PMRA 1187913	1997. Effect of high doses of Gliocladium catenulatum on different crops. DACO M10.3.1
PMRA 1187931	2006. Efficacy of Prestop WP strain J1446 for the control of Fusarium solani in greenhouse grown peppers. DACO M10.3.1