Health Canada Santé Canada

Proposed New Uses and / or Uses to be Withdrawn from the Label

In this form, there are several rows that allow you to input for multiple Sites of Use and Pests. For each Site of Use, please use a separate row. However, for each Site of Use, you can list multiple pests.

◉ New	○ Withdrawn	Site of Use	BLACKBERRY:
		Pests	Fruit Rot (Botrytis cinerea)
◉ New	○ Withdrawn	Site of Use	LOGANBERRY:
		Pests	Cane Spot (Elsinoë veneta), Fruit Rot (Botrytis cinerea), Leaf Spot (Septoria rubi), Spur Blight (Didymella applanate) -
◉ New	○ Withdrawn	Site of Use	BLUEBERRY:
		Pests	Fruit Rot (Botrytis cinerea), Mummy Berry (Monilinia vaccinii-corymbosi)
◉ New	○ Withdrawn	Site of Use	STRAWBERRY:
		Pests	Gray-Mold Rot (Botrytis cinerea), Leaf Spot (Mycosphaerella fragariae)
◉ New	○ Withdrawn	Site of Use	RHUBARB (IN FORCING SHEDS):
		Pests	Leaf Rot
◉ New	○ Withdrawn	Site of Use	VEGETABLES
		Pests	CUCUMBER: Anthracnose (Colletotrichum obriculare), Scab (Cladosporium cucumerinum), Early Blight (Alternaria solani), Late Blight (Phytophthora infestans)
◉ New	○ Withdrawn	Site of Use	VEGETABLES
		Pests	Anthracnose (Colletotrichum spp.), Septoria Leaf Spot (Septoria lycopersici)
◉ New	○ Withdrawn	Site of Use	GINSENG
		Pests	rhizoctonia root rot (Rhizoctonia solani), pythium root rot and damping-off (Pythium spp.), phytophthora root rot (Phytophthora cactorum) and grey mould (Botrytis cinerea) and sup
◉ New	○ Withdrawn	Site of Use	TURF and ORNAMENTAL USES
		Pests	**Brown Patch, Damping-Off , Leaf Spot and Melting-Out , Root Rot , Petal Blight, Leaf Spot, Botrytis Flower Blight, Septoria Leaf Spot, Black Spot**