

Evaluation Report for Category B, Subcategory B.2.1, 2.3, 2.4, 3.1, 3.12 Application

Application Number:	2008-1555
Application:	New end use product
Product:	Fungitrol 930 Fungicide
Registration Number:	29901
Active ingredients (a.i.):	3-Iodo-2-propynyl butyl carbamate (IPB)
PMRA Document Number	(English PDF): 1989770

Purpose of Application

The purpose of this application was to register a new end use product, Fungitrol 930 Fungicide, that contains the active ingredient 3-iodo-2-propynyl butyl carbamate (iodocarb). Fungitrol 930 Fungicide is for use as a preservative in exterior paint films, plastics, and adhesives.

Chemistry Assessment

Fungitrol 930 Fungicide is formulated as a suspension containing 3-iodo-2-propynyl butyl carbamate at a nominal concentration of 30% w/w. This end-use product has a density of 1.16-1.23 g/mL and a pH of 6-7. With the exception of the storage stability study, the chemistry requirements for Fungitrol 930 Fungicide are complete.

Health Assessments

Fungitrol 930 Fungicide is considered to be of slight acute toxicity via the oral route, of low acute toxicity via the dermal route, and of moderate acute toxicity via the inhalation route. Fungitrol 930 Fungicide is severely irritating to the eyes, moderately irritating to the skin, and is considered to be a potential dermal sensitizer.

A risk assessment was not performed for the Fungitrol end-use products as the use pattern fits within the existing use profile for the active ingredient, 3-iodo-2-propynyl butyl carbamate (iodocarb). The potential exposure to mixers, loaders, applicator and the post-applicators is not expected to exceed that of registered uses. Label amendments outlining supplementary personal protective equipment (PPE), required due to toxicology signal words and hazard statements, were provided.

Environmental Assessment

The active ingredient 3-iodo-2-propynyl butyl carbamate is toxic to aquatic organisms. Environmental exposure to these products through manufacturing processes can be mitigated through label statements.



Due to the use pattern as a preservative to be applied to exterior dry-film paints, adhesives and plastics, direct environmental exposure to Fungitrol 930 Fungicide is considered to be negligible if used according to the label.

Value Assessment

A standardized four-week laboratory test method was established and used to evaluate nine different paint samples, six different plastic samples, and two samples each of different cements and adhesives. Long term field studies, ranging from six months to two years, were carried out in New Jersey, a climate similar to much of Canada, for exterior paint dry-film and wood-plastic composite material. As a whole, the submitted data showed that 1000-2000 ppm of iodocarb was a reasonable range for the protection of exterior paint films from mildew. Wood-plastic composite material required 1000-3000 ppm of 3-iodo-2-propynyl butyl carbamate (iodocarb), while plastics and adhesives required less 3-iodo-2-propynyl butyl carbamate (iodocarb) (250-1000 and 200-500 ppm a.i., respectively) to protect them from fungal degradation.

Conclusion

The Pest Management Regulatory Agency has completed an assessment of the information provided in support for the product, Fungitrol 930 Fungicide, and has found the information sufficient to support full registration with the condition of the submission of a storage stability study.

References

<i>PMRA #</i> 1608459 1737859	<i>Title</i> 2006, IPBC: Evaluation of the Ambient Temperature Storage Stability, 1184/102-D2149, DACO: 3.5.10 CBI 2009, Fungitrol 940 Modified Chemistry Template, DACO: 3.0 CBI
1737860 1737861 1737862 1741584	 2006, Fungitrol 940 Test - Identity Methodologies, DACO: 3.0 CBI 2009, Oxidization Statement, DACO: 3.5.8 CBI 2009, Explanation for Variance in Viscosity, DACO: 3.5.9 CBI 2009, Plant Manufacturing Procedure for Fungitrol 930 Fungicide, DACO:
1753091 1803658 1803659 1803660 1803661 1803662 1803664	 3.2.2 CBI (Submission # 2008-1555) 2009, Fungitrol 940 Analytical Data, DACO: 3.4 CBI 2002, Acute Oral Toxicity in the Rat, DACO: 4.2.1,4.6.1 2002, Acute Dermal Toxicity Study in the Rat, DACO: 4.2.2,4.6.2 2001, Acute (Four-Hour) Inhalation Study in Rats, DACO: 4.2.3,4.6.3 2002, Acute Eye Irritation Study in the Rabbit, DACO: 4.2.4,4.6.4 2002, Acute Dermal Irritation Study in the Rabbit, DACO: 4.2.5,4.6.5 2002, Acute Dermal Sensitisation Study in the Guinea Pig (Buehler Method),
1925092	DACO: 4.2.6,4.6.6 2001, 8045-003 Acute Oral LD50 Definitive Study 01-09-01 to 01-24-01 Individual Clinical Signs, DACO: 4.6.1
1925093 1925094	 2000, 8045-003 Acute Oral Prelim RF 12-12-00 to 12-19-00 Individual Clinical Signs, DACO: 4.6.1 2000, 8045-003 Acute Oral Limit Test 5000 mg-kg Individual Clinical Signs, DACO: 4.6.1

- 1925095 2000, 8046-003 Acute Dermal Limit Test 5000 mg-kg Individual Clinical Signs, DACO: 4.6.2
- 1934211 2001, Fungitrol 430 Study 026-012394 Acute Inhalation Toxicity Individual Clinical Signs, DACO: 4.6.3
- 1963048 2001, 8045-003 Acute Oral LD50 Definitive Study 01-09-01 to 01-24-01 Individual Clinical Signs, DACO: 4.6.1
- 1963049 2000, 8045-003 Acute Oral Prelim RF 12-12-00 to 12-19-00 Individual Clinical Signs, DACO: 4.6.1
- 1963050 2000, 8045-003 Acute Oral Limit Test 5000 mg-kg Individual Clinical Signs, DACO: 4.6.1
- 1591685 2004, Efficacy of Fungitrol 420S Fungicide in Plastics and Plastic Coatings. International Specialty Products, Wayne NJ; 9 p.
- 1803611 2009, Efficacy Data for Various ISP-IPBC based formulation for PMRA Registrations. ISP Report; 37 p.

ISSN: 1911-8082

[®] Her Majesty the Queen in Right of Canada, represented by the Minister of Public Works and Government Services Canada 2011

All rights reserved. No part of this information (publication or product) may be reproduced or transmitted in any form or by any means, electronic, mechanical photocopying, recording or otherwise, or stored in a retrieval system, without prior written permission of the Minister of Public Works and Government Services Canada, Ottawa, Ontario K1A 0S5.