

Proposed New Uses and / or Uses to be Withdrawn from the Label

In this form, there are several rows that allow you to input for multiple Sites of Use and Pests. For each Site of Use, please use a separate row. However, for each Site of Use, you can list multiple pests.

<input checked="" type="radio"/> New	<input type="radio"/> Withdrawn	Site of Use	Red skinned potatoes, Spring Cereals (Wheat, Barley, Rye), Winter Cereals (Wheat, Barley, Rye)
		Pests	see weed list below
<input checked="" type="radio"/> New	<input type="radio"/> Withdrawn	Site of Use	Field Corn, Established Grasses Grown for Forage and Seed Production, Established Pasture and Rangeland (without legumes), Stubble Land, Roadsides, Uncropped Land, Airport
		Pests	see weed list below
<input checked="" type="radio"/> New	<input type="radio"/> Withdrawn	Site of Use	Golf Courses (roughs and fairways only), sod farms, CONTROL OF WOODY GROWTH IN NON-GRAZED AREAS, BASAL BARK AND STUMP TREATMENT
		Pests	Annual sow-thistle, Bluebur (before the 4-leaf stage), Burdock (before the 4-leaf stage), Cocklebur, Daisy fleabane, False flax, False ragweed, Flixweed, Giant ragweed,
<input checked="" type="radio"/> New	<input type="radio"/> Withdrawn	Site of Use	weed list continued
		Pests	Goat's-beard, Kochia, Lamb's-quarters, Mustards (except Dog and Tansy), Narrow-leaved hawk's-beard (in fall, and at the 1- to 2-leaf stage in spring), Plantain, Prickly lettuce,
<input type="radio"/> New	<input type="radio"/> Withdrawn	Site of Use	weed list continued
		Pests	Ragweeds, Redroot pigweed, Russian pigweed, Russian-thistle, Shepherd's-purse, Stinging nettle, Stinkweed, Sweet Clover (seedling), Thyme-leaved spurge, Volunteer canola,
<input type="radio"/> New	<input type="radio"/> Withdrawn	Site of Use	weed list continued
		Pests	Wild radish, Wild (prairie) sunflower, Curled dock (apply before 4-leaf stage), Dog mustard, Field pepper-grass, Flixweed (if treated before bolting in spring), Groundsel,
<input type="radio"/> New	<input type="radio"/> Withdrawn	Site of Use	weed list continued
		Pests	Hairy galinsoga, Hawkweed, Heal-all, Knotweed (apply before the 4-leaf stage), Narrow-leaved hawk's-beard (if treated before bolting in spring), Oak-leaved goosefoot,
<input type="radio"/> New	<input type="radio"/> Withdrawn	Site of Use	weed list continued
		Pests	Pineappleweed, Prostrate pigweed, Purslane, Sheep sorrel, Tansy mustard, Tumble pigweed, Velvetleaf, Biennial wormwood, Blue lettuce, Bull thistle, Burdock, Buttercup,
<input type="radio"/> New	<input type="radio"/> Withdrawn	Site of Use	weed list continued
		Pests	Canada thistle, Chicory, Curled dock, Dandelion, Field bindweed, Field chickweed*, Field horsetail*, Gumweed, Hedge bindweed, Hemp-nettle* (before the 4-leaf)