

Proposed New Uses and / or Uses to be Withdrawn from the Label

In this form, there are several rows that allow you to input for multiple Sites of Use and Pests. For each Site of Use, please use a separate row. However, for each Site of Use, you can list multiple pests.

<input checked="" type="radio"/> New	<input type="radio"/> Withdrawn	Site of Use	Spring Wheat
		Pests	Green smartweed , Pale smartweed , Lady's-thumb , Cow cockle, Redroot pigweed , Flixweed, Bluebur, Shepherd's-purse , Kochia, Russian thistle, Scentless chamomile
<input checked="" type="radio"/> New	<input type="radio"/> Withdrawn	Site of Use	Weeds continued
		Pests	Volunteer sunflower, Volunteer rapeseed (including Canola) , Night flowering catchfly, Cocklebur , Velvetleaf, Ball mustard, American nightshade, Jimsonweed, Wild tomato,
<input checked="" type="radio"/> New	<input type="radio"/> Withdrawn	Site of Use	Weeds continued
		Pests	Wild buckwheat , Tartary buckwheat , Common buckwheat , Stinkweed, Wild mustard , Wormseed mustard , Lamb's-quarters , Common ragweed , Common groundsel,
<input checked="" type="radio"/> New	<input type="radio"/> Withdrawn	Site of Use	Weeds continued
		Pests	Prickly lettuce, Canada thistle , Perennial sow-thistle
<input checked="" type="radio"/> New	<input type="radio"/> Withdrawn	Site of Use	Winter Wheat
		Pests	Same weeds as above
<input checked="" type="radio"/> New	<input type="radio"/> Withdrawn	Site of Use	Durum wheat
		Pests	Same weeds as above
<input checked="" type="radio"/> New	<input type="radio"/> Withdrawn	Site of Use	BARLEY
		Pests	Same weeds as above
<input checked="" type="radio"/> New	<input type="radio"/> Withdrawn	Site of Use	OATS
		Pests	Same weeds as above
<input checked="" type="radio"/> New	<input type="radio"/> Withdrawn	Site of Use	FLAX
		Pests	Same weeds as above