

## **Evaluation Report for Category B, Subcategory 3.12 Application**

**Application Number:** 2014-1585

**Application:** Changes to product labels – New site or host

**Product:** Puma Advance EC Herbicide

**Registration Number:** 29488

**Active ingredients (a.i.):** Fenoxaprop-P-ethyl (FPF)

PMRA Document Number: 2450045

## **Background**

Puma Advance EC Herbicide (Registration Number 29488; guarantee 90 g/L fenoxaprop-Pethyl) is presently registered for post-emergent control of wild oats, green foxtail, yellow foxtail, and barnyard grass in spring wheat, durum wheat, spring barley, and certain minor use crops (i.e., seedling and established perennial ryegrass and meadow bromegrass) in the Prairie Provinces and Peace River region of British Colombia and in spring wheat and spring barley in eastern Canada. Puma Advance EC Herbicide can be applied alone or in tank mix with listed broadleaf herbicides for the control of a broader spectrum of weeds. Use of Puma Advance EC Herbicide in durum wheat in eastern Canada was recently supported (2013-4318).

## **Purpose of Application**

The purpose of this application is to amend registration of Puma Advance EC Herbicide as described below.

- > To include use in the Okanagan and Creston Flats regions of British Columbia.
- > To use in all labelled crops regardless of geography.
- > To revise labelled tank mix recommendation.

#### **Chemistry Assessment**

A chemistry assessment was not required for this application.

## **Health Assessments**

A toxicology assessment was not required for this application.

Sufficient crop field trial data for wheat and barley are on file to support the national registration for Puma Advance EC Herbicide. The addition of Okanagan and Creston Flats Regions of British Columbia and the removal of geographical restriction on the label are not expected to have significant impact on the magnitude of the residues for fenoxaprop-p-ethyl and the safener, mefenpyr-diethyl, for the registered crops. Residues in/on treated wheat and



barley will continue to be covered under the MRLs of 0.05 ppm established each for fenoxapropethyl and mefenpyr-diethyl. Dietary exposure to fenoxaprop-pethyl or mefenpyr-diethyl is not expected to increase and will not pose an unacceptable risk to any segment of the population, including infants, children, adults and seniors.

#### **Environmental Assessment**

The modifications to the Puma Advance EC Herbicide use-pattern are not expected to pose an additional environmental risk.

#### Value Assessment

Value information in the form of a scientific rationale was submitted for review. The requested amendments to the registration of Puma Advance EC Herbicide, including (1) use in the Okanagan and Creston Flats regions of British Columbia and (2) use in all labelled crops regardless of geography, are supported for the following reasons.

- > The performance of a post-emergent herbicide is unlikely to be affected by soil properties or types.
- > Herbicides containing fenoxaprop-P-ethyl have been registered for use in the Prairie Provinces and Peace River region of British Columbia and eastern Canada for many years. Acceptable product performance and crop safety have been demonstrated across a range of environmental and climatic conditions.
- Application of fenoxaprop-P-ethyl during environmental conditions that result in plant stress may cause unacceptable crop injury and yield reduction. Environmental conditions that contribute to plant stress would not be expected to be any more prevalent in the Okanagan and Creston Flats regions of British Columbia than the geographic areas presently labelled for use of Puma Advance EC Herbicide, or in eastern Canada than western Canada.

All listed tank mixtures are supported for use in small grain cereals regardless of geography since (1) these tank mixes have been registered and used in small grain cereals in the Prairie Provinces and Peace River region of British Columbia for many years and (2) the labels contain a statement that reads "Always consult Puma Advance EC Herbicide and tank mix partner labels for proper timing, weeds controlled, precautions, cropping restrictions, etc and follow the more stringent of the two".

#### Conclusion

The Pest Management Regulatory Agency has completed an assessment of the information provided in support of the product, Puma Advance EC Herbicide, and has found the information sufficient to include use in the Okanagan and Creston Flats regions of British Columbia, use in all labelled crops regardless of geography and revise labelled tank mix recommendation.

#### References

2424181 2014, Puma Advance EC and Puma Advance: Rationale for use in the Okanagan and Creston Flats regions of British Columbia, in all labelled crops regardless of geography, and revision of tank mix recommendations, DACO: 10.1, 10.2.3.1, 10.2.3.3(B), 10.3.1, and 10.3.2(A).

ISSN: 1911-8082

# 8 Her Majesty the Queen in Right of Canada, represented by the Minister of Public Works and Government Services Canada 2016

All rights reserved. No part of this information (publication or product) may be reproduced or transmitted in any form or by any means, electronic, mechanical photocopying, recording or otherwise, or stored in a retrieval system, without prior written permission of the Minister of Public Works and Government Services Canada, Ottawa, Ontario K1A 0S5.