Health Canada Santé Canada

Proposed New Uses and / or Uses to be Withdrawn from the Label

In this form, there are several rows that allow you to input for multiple Sites of Use and Pests. For each Site of Use, please use a separate row. However, for each Site of Use, you can list multiple pests.

◉ New	○ Withdrawn	Site of Use	Grass Crops for Forage and Seed Production, Stubble Land, Uncropped Land,
		Pests	annual sow thistle, bluebur, burdock (before the 4-leaf stage), cocklebur,
◉ New	○ Withdrawn	Site of Use	"
		Pests	common plantain, daisy fleabane, false flax, flixweed, goat's beard, kochia, lamb's-quarters, mustards (except dog and tansy), prickly lettuce, ragweeds, redroot pigweed
◉ New	○ Withdrawn	Site of Use	"
		Pests	narrow-leaved hawk's beard (susceptible in the fall and at 1-2 leaf in the spring), Russian pigweed, Russian thistle, shepherd's purse, stinging nettle, stinkweed, sweet clover,
◉ New	○ Withdrawn	Site of Use	"
		Pests	thyme-leaved spurge, volunteer canola, wild radish, wild sunflower, chickweed, common purslane, curled dock (before the 4-leaf stage), dandelion (in lawn), dog mustard,
◉ New	○ Withdrawn	Site of Use	"
		Pests	flixweed (spring, prior to bolting), oakleaf goosefoot, groundsel, hairy galinsoga, common hawkweed, heal-all, knotweed (before the 4-leaf stage),
◉ New	○ Withdrawn	Site of Use	"
		Pests	narrow-leaved hawk'sbeard (spring, prior to bolting), peppergrass, pineappleweed, prostrate pigweed, Russian knapweed, sheep sorrel, smartweed (Pennsylvania and Green),
◉ New	○ Withdrawn	Site of Use	"
		Pests	tansy mustard, tumble pigweed, velvetleaf, yellow rocket (before the 4-leaf stage).
◉ New	○ Withdrawn	Site of Use	"
		Pests	Biennial wormwood, blue lettuce, bull thistle, burdock, buttercup, Canada thistle, dandelion, field bindweed, gumweed, hedge bindweed, hoary cress, leafy spurge,
◉ New	○ Withdrawn	Site of Use	"
		Pests	**mouse- eared chickweed, perennial sow-thistle, tartary buckwheat, teasel, wild buckwheat, yellow rocket.**