

Proposed New Uses and / or Uses to be Withdrawn from the Label

In this form, there are several rows that allow you to input for multiple Sites of Use and Pests. For each Site of Use, please use a separate row. However, for each Site of Use, you can list multiple pests.

<input checked="" type="radio"/> New	<input type="radio"/> Withdrawn	Site of Use	BLUEBERRIES, LOWBUSH BLUEBERRIES
		Pests	most common grasses and many perennial broadleaved weeds, and such woody weeds as trailing blackberry, lamb-kill
<input checked="" type="radio"/> New	<input type="radio"/> Withdrawn	Site of Use	Established Forage Alfalfa
		Pests	dandelion, quack grass , sow thistle, narrowleaved hawk's-beard and scentless chamomile
<input checked="" type="radio"/> New	<input type="radio"/> Withdrawn	Site of Use	LOWBUSH BLUEBERRY IN FRUITING YEAR (Nova Scotia, Prince Edward Island, and New Brunswick)
		Pests	golden rod, sheep sorrel, aster and poverty oatgrass
<input checked="" type="radio"/> New	<input type="radio"/> Withdrawn	Site of Use	HIGHBUSH BLUEBERRIES - BLUECROP VARIETY ONLY (For Use in British Columbia Only)
		Pests	colonial bent grass,poverty oat grass, blue grass, sheep fescue , timothy, Northern Panicgrass, yellow rattle, goldenrod , Aster, Yarrow, Cinquefoil , Hawkweed, Fireweed ,
<input checked="" type="radio"/> New	<input type="radio"/> Withdrawn	Site of Use	weeds controled continued,
		Pests	pearly everlasting, ox-eyed daisy, Black medick, trailing blackberry, Hardhack, Lamb-kill, Wild red raspberry, poplar
<input checked="" type="radio"/> New	<input type="radio"/> Withdrawn	Site of Use	ESTABLISHED SEED ALFALFA
		Pests	dandelion, quack grass , sow thistle, narrowleaved hawk's-beard and scentless chamomile
<input type="radio"/> New	<input type="radio"/> Withdrawn	Site of Use	
		Pests	
<input type="radio"/> New	<input type="radio"/> Withdrawn	Site of Use	
		Pests	
<input type="radio"/> New	<input type="radio"/> Withdrawn	Site of Use	
		Pests	