

## **New Pests**

# **Evaluation Report for Category B, Subcategory 3.11 Application**

**Application Number:** 2017-0885 **Application:** New Pests

**Product:** PP-2525 Herbicide

**Registration Number:** 29261

**Active ingredients (a.i.):** tribenuron-methyl and thifensulfuron-methyl

PMRA Document Number: 2780610

## **Background**

PP-2525 Herbicide was first registered March 30, 2009. PP-2525 Herbicide is registered for post-emergence application to spring wheat, including durum, winter wheat and spring barley at 30 g/ha in combination with 0.2% v/v of a non-ionic surfactant for the control or suppression of several broadleaved weed species. PP-2525 Herbicide may also be applied in tank mixtures with other herbicides, including with 0.21 L/ha Perimeter II Herbicide, containing 333 g a.e./L fluroxypyr, present as 1-methylheptyl ester, and 0.2% v/v of a non-ionic surfactant. For specific details of uses, application rates and methods, precautions, restrictions, and personal protective equipment requirements, refer to the product label.

## **Purpose of Application**

The purpose of this application is to amend the registration of PP-2525 Herbicide to include a claim of control of shepherd's purse (up to 20 cm) for the tank mixture of 30 g/ha PP-2525 Herbicide plus 0.21 L/ha (70 g a.e./ha) Perimeter II Herbicide plus 0.2% v/v of a non-ionic surfactant.

## **Chemistry Assessment**

A chemistry assessment was not required as there was no change to the product formulation.

### **Health and Environmental Assessment**

Health and environmental assessments were not required as there was no change to host crops, application rates, methods, and timings.

#### Value Assessment

The addition of a claim of control of shepherd's purse for the tank mixture of PP-2525 Herbicide plus Perimeter II Herbicide plus non-ionic surfactant will provide wheat and barley growers an additional option to control this troublesome weed.



Value information was submitted as efficacy data from three replicate small-scale field trials. Two trials were conducted on spring barley and a third on fallowed land. Control of shepherd's purse was usually achieved by four weeks after application and was consistently achieved by six weeks after application.

## **Conclusion**

The PMRA has conducted an assessment of the subject application and has determined that the submitted information is adequate to support a claim of control of shepherd's purse (up to 20 cm) for PP-2525 Herbicide tank mixed with Perimeter II Herbicide and non-ionic surfactant applied in accordance with the PP-2525 Herbicide label.

#### References

## List of Studies/Information Submitted by Registrant

## Value Assessment

2729713	2017, Control of Shepherds Purse with a Tank Mix of Barricade SG +
	Perimeter II Herbicide in Cereal Crops, DACO:
	10.1,10.2,10.2.1,10.2.2,10.2.3,10.2.3.1,10.2.3.3,10.2.3.3(B),10.3.2(A)

ISSN: 1911-8082

8 Her Majesty the Queen in Right of Canada, represented by the Minister of Public Works and Government Services Canada 2017

All rights reserved. No part of this information (publication or product) may be reproduced or transmitted in any form or by any means, electronic, mechanical photocopying, recording or otherwise, or stored in a retrieval system, without prior written permission of the Minister of Public Works and Government Services Canada, Ottawa, Ontario K1A 0S5.