

GUARDSMAN

AGRICULTURAL WEEDKILLER NO.1

UNIVAR CANADA LTD.
9800 Van Horne Way,
Richmond, B.C., V6X 1W5

Net Contents : 205 Litres

SELECTIVE HERBICIDAL OIL AGRICULTURAL

READ THE LABEL BEFORE USING

REGISTRATION NO. 2076 PEST CONTROL PRODUCTS ACT

GUARANTEE: MINERAL SPIRITS 100%

DIRECTIONS:

AGRICULTURAL WEEDKILLER NO.1 is a selective herbicide specifically recommended for use in carrots, parsnips, parsley, dill, celery, cranberries, and asparagus. It is supplied ready to use.

Agricultural Weedkiller No.1 will control such weeds as redroot pigweed, lamb's-quarters, purslane, annual grasses and cottontop. A few weeds including ragweed, wild carrot, perennial grasses and cloudberry are not easily killed.

Aerial Application Information - DO NOT APPLY BY AIR

CARROTS, PARSNIPS, PARSLEY AND DILL:

Apply when plants are in the 2 to 4 leaf stage and before weeds are 10 cm high. For treatment of the whole field apply 600 to 800 litres per hectare depending on size and stand of weeds. For treatment of rows only, apply 300 to 400 litres per hectare depending on size and stand of weeds. Do not apply within 30 days of harvest.

CELERY:

(1)Seed beds - Apply after plants have emerged and as soon as weed growth warrants. Apply approximately 6 litres per 100 m². Ventilation is essential following application in the seedbeds.

(2)Field - Apply before hearts form or before plants are over 15 cm in height. This is usually within 2 to 3 weeks of transplanting. For treatment of the whole field apply 600 to 800 litres per hectare depending on size and stand of weeds. For treatment of rows only apply 300 to 400 litres of spray per hectare depending on size and stand of weeds. Follow recommendations for use closely since celery is easily damaged during any stage of growth.

CRANBERRIES:

For control of cottontop and partial control of cloudberry (baked-apple berry) in peat bogs, apply at rates of 12,000 to 16,000 litres per hectare to infested spots before cranberry buds break. For control of annual grasses, sedges, bracken fern and broadleaf seedlings apply at 3,500 to 5,00 litres per hectare. Apply in March to early April.

NOTE:

Preferably apply the celery and cranberry treatments during a cool, sunny morning when leaves are dry; do not apply during strong winds, the heat of the day, a rain, or immediately before or after a frost.

ASPARAGUS:

For control of seedling bluegrasses, chickweed, barnyard grass, and other weeds in seedbeds, apply at 500 litres per hectare as an overall spray. Apply when crop seedlings are 5 cm high and seedling weeds and grasses have 3 to 4 true leaves.

In the producing field, apply before the crop emerges when the grass seedlings are in the 2 to 3 leaf stage, or 3-4 cm in height.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. Keep in original container during storage. Do not allow spray, spray mist, visible or invisible vapours or air containing mist or vapours to contact any plants other than those being sprayed. Spray all areas uniformly. Handle and apply only as recommended and at recommended rates. Do not contaminate streams, lakes, ponds, irrigation water, water used by livestock or water used for domestic purposes. Keep product away from fire, open flame, lighted electric bulbs and other sources of heat.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at: www.croplife.ca.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

FIRST AID:

EYE CONTACT: Hold eye open and rinse slowly and gently with water for 15 -20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

SKIN OR CLOTHING CONTACT: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 -20 minutes. Call a poison control centre or doctor for treatment advice.

INHALATION: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to mouth, if possible. Call a poison control centre or doctor for further treatment advice.

INGESTION: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Treat symptomatically

DISPOSAL:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.