

2023-7083  
2024-05-31

GROUP

1B

INSECTICIDE

BASE

**LAGON® 480 E**

INSECTICIDE

COMMERCIAL

This product is not to be used in and around homes or other residential areas such as parks, school grounds, playing fields. It is not for use by homeowners or other uncertified users.



WARNING

POISON

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

ACTIVE INGREDIENT: Dimethoate 480 g/L

REGISTRATION NO. 9382 PEST CONTROL PRODUCTS ACT

IN CASE OF EMERGENCY DUE TO A MAJOR SPILL, FIRE OR POISONING  
INVOLVING THIS PRODUCT CALL DAY OR NIGHT, 1-800-561-8273

LOVELAND PRODUCTS CANADA INC.

789 Donnybrook Drive

Dorchester, ON

N0L 1G5

1-800-328-4678

NET CONTENTS: 1 L – 115 L

1/16

2023-7083  
2024-05-31

GROUP	<b>1B</b>	INSECTICIDE
-------	-----------	-------------

FRONT

**LAGON® 480 E**

INSECTICIDE

COMMERCIAL

This product is not to be used in and around homes or other residential areas such as parks, school grounds, playing fields. It is not for use by homeowners or other uncertified users.



WARNING

POISON

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

ACTIVE INGREDIENT: Dimethoate 480 g/L

REGISTRATION NO. 9382 PEST CONTROL PRODUCTS ACT

IN CASE OF EMERGENCY DUE TO A MAJOR SPILL, FIRE OR POISONING  
INVOLVING THIS PRODUCT CALL DAY OR NIGHT, 1-800-561-8273

LOVELAND PRODUCTS CANADA INC.

789 Donnybrook Drive

Dorchester, ON

N0L 1G5

1-800-328-4678

NET CONTENTS: 1 L – 115 L

1/16

2023-7083  
2024-05-31

**PRECAUTIONS**  
**HARMFUL OR FATAL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN**  
**KEEP OUT OF REACH OF CHILDREN**

Keep the following personal protective equipment immediately available for use in case of emergency (for example, a broken package, spill, or equipment breakdown): chemical-resistant coveralls, chemical-resistant gloves, chemical-resistant head gear and a respirator.

**DO NOT** apply in greenhouses.

**DO NOT** use this product in and around homes or other residential areas such as parks, school grounds and/or playing fields. It is not for use by homeowners or other uncertified users.

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

**DO NOT** allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted. It is desirable that the pilot have communication capabilities at each treatment site at the time of application. All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft and cockpit and vehicle cabs must be decontaminated regularly.

Avoid breathing vapour or spray mist. Use only with adequate ventilation. **DO NOT** use indoors. Avoid prolonged or repeated contact with skin. Wash thoroughly after handling and before eating and smoking. Avoid contamination of feed and food stuffs. Keep away from heat, sparks and open flame. Keep out of sun.

If this pest control product is to be used on a commodity that may be exported and you require information regarding Maximum Residue Limits for an importing country, please contact LOVELAND PRODUCTS CANADA INC.

**Engineering Controls and Personal Protective Equipment:**  
**A. Mixing/loading, clean-up and repair (for all uses except forestry):**

Wear chemical-resistant coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, chemical-resistant footwear, protective eyewear (goggles or face shield) and suitable respiratory protection. Suitable respiratory protection is defined as either a respirator with a NIOSH-approved organic-vapour-removing cartridge (with a prefilter) approved for pesticides or a NIOSH-approved canister approved for pesticides.

**B. Mixing/loading, clean-up and repair (forestry: hemlock, spruce and balsam fir):**

Closed mixing/loading systems are required. A closed system means removing a pesticide from its original container, rinsing, mixing, diluting, and transferring the pesticide through connecting hoses and couplings that prevent exposure to the pesticide. Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, chemical-resistant footwear and protective eyewear (goggles or face shield).

**C. Mixing/loading, clean-up and repair (forestry: Douglas-fir (seed tree), Sitka spruce (seed tree), spruce (seed tree and woodland)):**

Wear chemical-resistant coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, chemical-resistant footwear, protective eyewear (goggles or face shield) and suitable respiratory protection. Suitable respiratory protection is defined as either a respirator with a NIOSH-approved organic-vapour-removing cartridge (with a prefilter) approved for pesticides or a NIOSH-approved canister approved for pesticides.

**D. Applying by air:**

Wear coveralls over a long-sleeved shirt, long pants, socks and chemical-resistant footwear.

**E. Applying by groundboom and/or airblast:**

Use a closed cab that provides both a physical barrier and respiratory protection (for example, dust/mist filtering and/or vapour/gas purification system). The closed cab must have a chemical-resistant barrier that totally surrounds the occupant and prevents contact with pesticides outside the cab. Wear coveralls over a long-sleeved shirt, long pants, socks and chemical-resistant footwear.

**F. Applying by handheld equipment:**

Wear chemical-resistant coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, chemical-resistant footwear, and suitable respiratory protection. Suitable respiratory protection is defined as either a respirator with a NIOSH-approved organic-vapour-removing cartridge (with a prefilter) approved for pesticides or a NIOSH-approved canister approved for pesticides. Wear eye and head protection when applying above waist height, including overhead. Limit the amount of active ingredient handled per day to 2.8 kg per person (for example, approximately 2950 L at a rate of 0.96 g a.i./L) when applying by handheld equipment.

**G. Applying by manually-pressurized handwand or backpack (Douglas-fir (seed tree), Sitka spruce (seed tree), spruce (seed tree and woodland)):**

Wear chemical-resistant coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, chemical-resistant footwear and suitable respiratory protection. Suitable respiratory protection is defined as either a respirator with a NIOSH-approved

2023-7083  
2024-05-31

organic-vapour-removing cartridge (with a prefilter) approved for pesticides OR a NIOSH-approved canister approved for pesticides. Wear eye and head protection when applying above waist height, including overhead. **DO NOT** apply using mechanically-pressurized handgun equipment. Only for use with manually-pressurized handwands or backpack sprayers.

#### **H. Applying by right-of-way sprayer:**

Wear chemical-resistant coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, chemical-resistant footwear, protective eyewear (goggles or face shield) and suitable respiratory protection. Suitable respiratory protection is defined as either a respirator with a NIOSH-approved organic-vapour-removing cartridge (with a prefilter) approved for pesticides OR a NIOSH-approved canister approved for pesticides. Wear head protection when applying above waist height, including overhead.

#### **I. Applying by soil drench, soil injection or chemigation:**

Wear chemical-resistant coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, chemical-resistant footwear and suitable respiratory protection. Suitable respiratory protection is defined as either a respirator with a NIOSH-approved organic-vapour-removing cartridge (with a prefilter) approved for pesticides or a NIOSH-approved canister approved for pesticides.

#### **ENVIRONMENTAL PRECAUTIONS:**

This product contains an active ingredient and aromatic petroleum distillate, which are toxic to aquatic organisms. Observe spray buffer zones specified under DIRECTIONS FOR USE.

Toxic to certain beneficial arthropods (which may include predatory and parasitic insects, spiders, and mites). Minimize spray drift to reduce harmful effects on beneficial arthropods in habitats next to the application site such as hedgerows and woodland .

TOXIC to birds and small wild mammals.

Toxic to bees. Bees may be exposed through direct spray, spray drift, and residues on/in leaves, pollen and nectar in flowering crops and weeds. Minimize spray drift to reduce harmful effects on bees in habitats close to the application site. Avoid applications when bees are foraging in the treatment area in ground cover containing blooming weeds. To further minimize exposure to pollinators, refer to the complete guidance "Protecting Pollinators during Pesticide Spraying – Best Management Practices" on the Health Canada website ([www.canada.ca/pollinators](http://www.canada.ca/pollinators)). Follow crop specific directions for application timing.

**For applications on crops that are highly attractive to pollinators (alfalfa, clovers, canola, safflower, blueberries, cherries, peaches, pears, asparagus, and outdoor ornamentals excluding coniferous evergreens), or when using managed bees for pollination services:**

DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.

#### **For applications on all other crops:**

Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

#### **DECONTAMINATION**

If accidental spillage of LAGON 480 E Insecticide should occur, scrub contaminated area immediately with a strong laundry soap solution or use household lye. Detergents are not satisfactory for this purpose. Repeated scrubbing is necessary on plain wood surfaces. Avoid spillage on all types of floor tile or linoleum.

#### **FIRST AID**

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

**IF SWALLOWED:** Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by poison control centre or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferable by mouth-to-mouth, if possible. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

#### **TOXICOLOGICAL INFORMATION**

Dimethoate is an organophosphate that is a cholinesterase inhibitor. Typical symptoms of overexposure to cholinesterase inhibitors include headache, nausea, dizziness, sweating, salivation, and runny nose and eyes. This may progress to muscle twitching, weakness, tremor, incoordination, vomiting, abdominal cramps and diarrhea in more serious poisonings. A life-threatening poisoning is signified by loss of consciousness, incontinence, convulsions and respiratory depression with a secondary cardiovascular component. Treat symptomatically. If exposed, plasma and red blood cell cholinesterase tests may indicate degree of exposure (baseline data are useful). Atropine, only by injection, is the preferable antidote. Oximes, such as pralidoxime chloride, may be

2023-7083  
2024-05-31

therapeutic if used early; however, use only in conjunction with atropine. In cases of severe acute poisoning, use antidotes immediately after establishing an open airway and respiration. With oral exposure, the decision of whether to induce vomiting or not should be made by an attending physician. This product contains petroleum distillates. Vomiting may cause aspiration pneumonia. Treat symptomatically.

#### **STORAGE**

Store this product away from food or feed.

Do not use, pour, spill, or store near heat, sparks, or open flame. Keep out of direct sunlight. Ship and store between 4 to 30°C.

#### **DISPOSAL**

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

**NOTICE TO USER:** This pest control product is to be used only in accordance with the directions on this label. It is an offence under the *PEST CONTROL PRODUCTS ACT* to use this product in a way that is inconsistent with the directions on the label.

2023-7083  
2024-05-31

GROUP	1B	INSECTICIDE
-------	----	-------------

DATAPAK

**LAGON® 480 E**

INSECTICIDE

COMMERCIAL

This product is not to be used in and around homes or other residential areas such as parks, school grounds, playing fields. It is not for use by homeowners or other uncertified users.



WARNING

POISON

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

ACTIVE INGREDIENT: Dimethoate 480 g/L

REGISTRATION NO. 9382 PEST CONTROL PRODUCTS ACT

IN CASE OF EMERGENCY DUE TO A MAJOR SPILL, FIRE OR POISONING  
INVOLVING THIS PRODUCT CALL DAY OR NIGHT, 1-800-561-8273

LOVELAND PRODUCTS CANADA INC.  
789 Donnybrook Drive  
Dorchester, ON  
N0L 1G5  
1-800-328-4678

NET CONTENTS: 1 L – 115 L

1/16

2023-7083  
2024-05-31

**PRECAUTIONS**  
**HARMFUL OR FATAL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN**  
**KEEP OUT OF REACH OF CHILDREN**

Keep the following personal protective equipment immediately available for use in case of emergency (for example, a broken package, spill, or equipment breakdown): chemical-resistant coveralls, chemical-resistant gloves, chemical-resistant head gear and a respirator.

**DO NOT** apply in greenhouses.

**DO NOT** use this product in and around homes or other residential areas such as parks, school grounds and/or playing fields. It is not for use by homeowners or other uncertified users.

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

**DO NOT** allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted. It is desirable that the pilot have communication capabilities at each treatment site at the time of application. All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft and cockpit and vehicle cabs must be decontaminated regularly.

Avoid breathing vapour or spray mist. Use only with adequate ventilation. **DO NOT** use indoors. Avoid prolonged or repeated contact with skin. Wash thoroughly after handling and before eating and smoking. Avoid contamination of feed and food stuffs. Keep away from heat, sparks and open flame. Keep out of sun.

If this pest control product is to be used on a commodity that may be exported and you require information regarding Maximum Residue Limits for an importing country, please contact LOVELAND PRODUCTS CANADA INC.

**Engineering Controls and Personal Protective Equipment:**

**A. Mixing/loading, clean-up and repair (for all uses except forestry):**

Wear chemical-resistant coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, chemical resistant footwear, protective eyewear (goggles or face shield) and suitable respiratory protection. Suitable respiratory protection is defined as either a respirator with a NIOSH-approved organic-vapour-removing cartridge (with a prefilter) approved for pesticides or a NIOSH-approved canister approved for pesticides.

**B. Mixing/loading, clean-up and repair (forestry: hemlock, spruce and balsam fir):**

Closed mixing/loading systems are required. A closed system means removing a pesticide from its original container, rinsing, mixing, diluting, and transferring the pesticide through connecting hoses and couplings that prevent exposure to the pesticide. Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, chemical-resistant footwear and protective eyewear (goggles or face shield).

**C. Mixing/loading, clean-up and repair (forestry: Douglas-fir (seed tree), Sitka spruce (seed tree), spruce (seed tree and woodland)):**

Wear chemical-resistant coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, chemical-resistant footwear, protective eyewear (goggles or face shield) and suitable respiratory protection. Suitable respiratory protection is defined as either a respirator with a NIOSH-approved organic-vapour-removing cartridge (with a prefilter) approved for pesticides or a NIOSH-approved canister approved for pesticides.

**D. Applying by air:**

Wear coveralls over a long-sleeved shirt, long pants, socks, and chemical-resistant footwear.

**E. Applying by groundboom and/or airblast:**

Use a closed cab that provides both a physical barrier and respiratory protection (for example, dust/mist filtering and/or vapour/gas purification system). The closed cab must have a chemical resistant barrier that totally surrounds the occupant and prevents contact with pesticides outside the cab. Wear coveralls over a long-sleeved shirt, long pants, socks, and chemical-resistant footwear.

**F. Applying by handheld equipment:**

Wear chemical-resistant coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, chemical-resistant footwear, and suitable respiratory protection. Suitable respiratory protection is defined as either a respirator with a NIOSH-approved organic-vapour-removing cartridge (with a prefilter) approved for pesticides or a NIOSH-approved canister approved for pesticides. Wear eye and head protection when applying above waist height, including overhead. Limit the amount of active ingredient handled per day to 2.8 kg per person (for example, approximately 2950 L at a rate of 0.96 g a.i./L) when applying by handheld equipment.

**G. Applying by manually-pressurized handwand or backpack (Douglas-fir (seed tree), Sitka spruce (seed tree), spruce (seed tree and woodland)):**

2023-7083  
2024-05-31

During manually-pressurized handwand, or backpack application, wear chemical-resistant coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, chemical-resistant footwear and suitable respiratory protection. Suitable respiratory protection is defined as either a respirator with a NIOSH-approved organic-vapour-removing cartridge (with a prefilter) approved for pesticides OR a NIOSH-approved canister approved for pesticides. Wear eye and head protection when applying above waist height, including overhead. **DO NOT** apply using mechanically-pressurized handgun equipment. Only for use with manually-pressurized handwands or backpack sprayers.

#### **H. Applying by right-of-way sprayer:**

Wear chemical-resistant coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, chemical-resistant footwear, protective eyewear (goggles or face shield) and suitable respiratory protection. Suitable respiratory protection is defined as either a respirator with a NIOSH-approved organic-vapour-removing cartridge (with a prefilter) approved for pesticides OR a NIOSH-approved canister approved for pesticides. Wear head protection when applying above waist height, including overhead.

#### **I. Applying by soil drench, soil injection or chemigation:**

Wear chemical-resistant coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, chemical-resistant footwear and suitable respiratory protection. Suitable respiratory protection is defined as either a respirator with a NIOSH-approved organic-vapour-removing cartridge (with a prefilter) approved for pesticides or a NIOSH-approved canister approved for pesticides.

#### **ENVIRONMENTAL PRECAUTIONS:**

This product contains an active ingredient and aromatic petroleum distillate, which are toxic to aquatic organisms. Observe spray buffer zones specified under DIRECTIONS FOR USE.

Toxic to certain beneficial arthropods (which may include predatory and parasitic insects, spiders, and mites). Minimize spray drift to reduce harmful effects on beneficial arthropods in habitats next to the application site such as hedgerows and woodland.

TOXIC to birds and small wild mammals.

Toxic to bees. Bees may be exposed through direct spray, spray drift, and residues on/in leaves, pollen and nectar in flowering crops and weeds. Minimize spray drift to reduce harmful effects on bees in habitats close to the application site. Avoid applications when bees are foraging in the treatment area in ground cover containing blooming weeds. To further minimize exposure to pollinators, refer to the complete guidance "Protecting Pollinators during Pesticide Spraying – Best Management Practices" on the Health Canada website ([www.canada.ca/pollinators](http://www.canada.ca/pollinators)). Follow crop specific directions for application timing.

**For applications on crops that are highly attractive to pollinators (alfalfa, clovers, canola, safflower, blueberries, cherries, peaches, pears, asparagus, and outdoor ornamentals excluding coniferous evergreens), or when using managed bees for pollination services:**

DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.

#### **For applications on all other crops:**

Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

#### **DIRECTIONS FOR USE**

Lagon 480E INSECTICIDE is a true systemic insecticide. A systemic insecticide is absorbed into the system of the plant upon application and, as with all systemic materials, may in specific plants cause reactions which are neither predictable nor common to all members of the species.

LAGON 480 E Insecticide is generally effective in controlling aphids, and other listed insects, scales and mites.

LAGON 480 E Insecticide may be applied by spray or soil drench. LAGON 480 E Insecticide is an emulsifiable concentrate. Use designated amounts in full volume of water.

#### **TANK MIXTURES**

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Loveland Products Canada Inc. at 1-800-328-4678 for information before applying any tank mix that is not specifically recommended on this label.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

Field sprayer application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO



2023-7083  
2024-05-31

NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) fine classification. Boom height must be 60 cm or less above the crop or ground.

**Airblast application:** DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT direct spray above plants to be treated. Turn off outward pointing nozzles at row ends and outer rows. DO NOT apply when wind speed is greater than 16 km/h at the application site as measured outside of the treatment area on the upwind side.

**Chemigation:** DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) fine classification. Applications MUST be conducted WITHOUT the use of end guns.

**Aerial application:** DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply when wind speed is greater than 16 km/h at flying height at the site of application. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) fine classification. Reduce drift caused by turbulent wingtip vortices. Nozzle distribution along the spray boom length MUST NOT exceed 65% of the wing - or rotorspan.

Apply only by fixed-wing or rotary aircraft equipment, which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices.

**SPRAY BUFFER ZONES:**

A spray buffer zone is NOT required for:

- uses with hand-held application equipment permitted on this label,
- soil drench or soil incorporation.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of application	Crop	Spray Buffer Zones (metres) Required for the Protection of:			
		Freshwater habitat of Depth:		Estuarine/Marine habitats of Depth:	
		Less than 1m	Greater than 1m	Less than 1m	Greater than 1m
Field Sprayer*	Cereals (barley, oats, rye, wheat), forage crops (clover, grasses, hay, grain), alfalfa (forage and seed production), clover (sweet, red, alsike), pastures and wastelands, safflower, soybeans, bok choy (pak choy), chicory, Chinese broccoli, cole crops (Brussels sprouts), peppers, potatoes, tomatoes	2	1	1	1
	Flax, canary grass (seed production)	1	0	1	0
	Canola (rape), beans (pole, snap), beets, celery, eggplant, leafy vegetables (beet greens, kale, leaf lettuce, Swiss chard, turnip greens), Blueberry (low and highbush)	1	1	1	1
	Asparagus	3	1	2	1
	Cole crops (broccoli, cauliflower)	2	1	2	1
	Peas (field and/or canning)	1	0	1	0
	Strawberries	3	2	3	1
	Christmas Trees: Balsam fir, hemlock, pine (Mugho, red, Scots), spruce; Ornamental trees: Arborvitae, cedar, juniper, oak, pine (Mugho, red, Scots), Taxus (yew), Lilac ( <i>Syringa</i> spp.)	2	1	2	1
	Birch	1	0	1	0
	Boxwood, hemlock, holly (English, American)	1	1	1	1
	Douglas-fir (seed tree), spruce (seed tree and woodland)	10	5	10	5
	Larch	3	2	4	2
	Sitka spruce (seed tree)	15	5	10	5
	Day lilies, irises, euonymus	2	1	2	1

2023-7083  
2024-05-31

	Roses, azaleas, gardenias, camellias, gerberas, gladioli, poinsettias		1	1	1	1
Airblast	Christmas trees Balsam fir, hemlock, pine (Mugho, red, Scots), spruce; Ornamental trees: Arborvitae, cedar, juniper, oak, pine (Mugho, red, Scots), Taxus (yew)	Late growth stage	4	2	4	2
	Birch	Late growth stage	1	0	1	0
	Boxwood, hemlock, holly (English, American)	Late growth stage	2	1	2	1
	Douglas-fir (seed tree)	Late growth stage	25	15	20	15
	Larch	Late growth stage	5	3	10	4
	Lilac (Syringa spp.)	Late growth stage	3	1	3	1
	Sitka spruce (seed tree)	Late growth stage	25	15	25	15
	Spruce (seed tree and woodland)	Late growth stage	20	10	25	15
	Euonymus	Late growth stage	4	2	4	2
	Roses	Late growth stage	2	1	2	1
	Poinsettias	Late growth stage	2	1	2	1
	Blueberry (low and highbush)	Early growth stage	4	1	3	1
		Late growth stage	2	1	2	1
	Tree nuts (filberts, hazelnuts)	Early growth stage	15	5	20	10
		Late growth stage	5	3	10	4
	Peaches	Early growth stage	20	10	15	5
		Late growth stage	10	4	10	4
	Pears	Early growth stage	15	5	15	5
		Late growth stage	10	3	5	3
	Cherries (sour, sweet)	Early growth stage	10	4	10	3
		Late growth stage	5	2	4	2
Aerial	Flax, canary grass (seed production), peas (field and/or canning)	Fixed wing	1	0	1	0
		Rotary wing	1	0	1	0
	Cereals (barley, oats, rye, wheat), forage crops (clover, grasses, hay, grain), non-cropland (fence rows, field edges except adjacent to unregistered crops), pastures and wastelands, safflower, soybeans	Fixed wing	10	1	5	1
		Rotary wing	5	1	1	1

2023-7083  
2024-05-31

Alfalfa (forage and seed production), clover (sweet, red, alsike)	Fixed wing	10	1	5	1
	Rotary wing	5	1	3	1
Beans (pole, snap), soybeans	Fixed wing	5	1	5	0
	Rotary wing	3	1	1	0
Hemlock	Fixed wing	90	0	150	15
	Rotary wing	55	0	90	1
Blueberry (low and highbush)	Fixed wing	4	1	1	1
	Rotary wing	1	1	1	1

\*For field sprayer application, buffer zones can be reduced with the use of drift reducing spray shields. When using a spray boom fitted with a full shield (shroud, curtain) that extends to the crop canopy, the labelled buffer zone can be reduced by 70%. When using a spray boom where individual nozzles are fitted with cone-shaped shields that are no more than 30 cm above the crop canopy, the labelled buffer zone can be reduced by 30%.

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pesticides portion of the Canada.ca website.

#### RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, please note that LAGON 480 E Insecticide contains a Group 1B insecticide/acaricide. Any insect/mite population may contain individuals naturally resistant to LAGON 480 E Insecticide and other Group 1B insecticide/acaricide. The resistant individuals may dominate the insect/mite population if this group of insecticides/acaricides is used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but are specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay insecticide/acaricide resistance:

- Where possible, rotate the use of LAGON 480 E Insecticide or other Group 1B insecticides/acaricides with different groups that control the same pests.
- Avoid application of more than the indicated number of sprays of LAGON 480 E Insecticide or other insecticides/acaricides in the same group in a season.
- Use tank mixtures with insecticides/acaricides from a different group that is effective on the target pest when such use is permitted.
- Insecticide/acaricide use should be based on an IPM program that includes scouting and record keeping, and considers cultural, biological and other chemical control practices.
- Monitor treated pest populations for resistance development.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance -management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information and to report suspected resistance, contact Technical Service at 1-800-265-4624 or at [www.uap.ca](http://www.uap.ca).

**FOR GROUND APPLICATION:** dilute with water to 200 - 300 L/ha unless otherwise stated.

**FOR AERIAL APPLICATION:** unless otherwise stated, dilute with water to 11 - 22 L/ha for blueberries, cereals (barley, oats, wheat, rye) and peas. Dilute with water to 10-30 L/ha for other field and vegetable crops when aerial application is specified.

#### Use Precautions

Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the *National Aerial Pesticide Application Manual*, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

#### Product Specific Precautions

Read and understand the entire label before opening this product. If you have questions, call the manufacturer at (800)328-4678 or obtain technical advice from the distributor or your provincial agricultural representative. Application of this specific product must meet and/or conform to the following:

Volume: Apply the recommended rate in a minimum spray volume of 10 litres per hectare.

#### FIELD AND VEGETABLE CROPS:

Sprays should be applied when the pests first appear and repeated as the infestation warrants, or as otherwise directed. Applications should be made in sufficient water to obtain good coverage. Apply at the recommended rate; do not exceed a maximum spray volume of 1000 L per hectare unless otherwise stated. When aerial application is specified apply with aerial equipment and use 10 to 30L of water per hectare. Applications may be made by ground equipment unless otherwise specified.

FIELD AND VEGETABLE CROPS	PESTS CONTROLLED	AMOUNT PER HECTARE	AMOUNT PER 10 LITRES	PREHARVEST INTERVALS (DAYS)	MAXIMUM # OF APPLICATIONS PER YEAR
---------------------------	------------------	--------------------	----------------------	-----------------------------	------------------------------------

2023-7083

2024-05-31

Alfalfa  
(for seed and  
forage production)

Aphids, Leafhoppers reduction of alfalfa weevil larvae	425 mL	4 mL	10	2
--	--------	------	----	---

COMMENTS: Do not graze or harvest for forage within 10 days after application. Minimum application interval is 7 days. All activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.

Blotch leafminers	550 mL	5 mL	10	1
-------------------	--------	------	----	---

COMMENTS: : Do not graze or harvest for forage within 10 days after application. Use a water volume of at least 200 L/ha with ground equipment for Blotch leafminers. Do not apply during bloom. May be applied by Ground and Aerial application. All activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.

Grasshoppers - nymphs	550 mL	5 mL	10	1
Grasshoppers -adults or winged	850 - 900 mL	9 mL	10	2

COMMENTS: Do not graze or harvest for forage within 10 days after treatment or when using a rate of 850 mL or higher: Do not graze or harvest for forage within 28 days after treatment. Follow provincial forecast. Apply when insects are present as young hoppers or signs of insect damage are evident. Apply when the grasshoppers are in the 2-4 nymphal stage. Best control will be achieved when application is made prior to wing development. Under severe insect pressure application should also be made to a 15 metre strip along fence rows around the field. DO NOT APPLY TO ADJACENT UNREGISTERED CROPS. The higher rate should be used when the proportion of mature and late nymphal stages in the population are high and spray penetration is inhibited by dense crop canopy. Minimum application interval is 7 days. All activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.

Alfalfa  
(seed production)

Lygus bugs Plant bugs	1.1 L	10 mL	28	1
--------------------------	-------	-------	----	---

COMMENTS: Apply just before alfalfa begins to bloom. Allow at least 10 days before placing leaf cutting bees in the field. Do not graze or harvest for forage with 28 days after treatment. May be applied by Ground and Aerial application. All activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.

Asparagus

Asparagus aphid	2.3 L	35 mL	Not Applicable	2
-----------------	-------	-------	----------------	---

COMMENTS:  
Ground application only.  
Do not apply during the crop bloom period or during the 5-day period before the crop blooms.

For mature asparagus, applications can be made post-harvest only, until plant defoliation. For immature asparagus, begin application mid-May. Apply by boom or field sprayer using 675 L of water per hectare. For immature asparagus; **DO NOT** harvest for feed or food. Ground application ONLY. Max number of applications per year is 2. Minimum application interval is 7 days. All activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.

Barley, Oats, Wheat  
(ALSO SEE  
CEREALS below)

Thrips	1 L	10 mL	35	2
--------	-----	-------	----	---

COMMENTS: Ground Application: Use sufficient water to obtain good coverage (usually 110 – 275 L/ha). Same limitations apply for harvesting and grazing as in wheat section of this label. Ground and Aerial application. With aerial equipment use 10 to 30 L of water per hectare. Maximum number of applications per year is 2. Minimum application interval is 7 days. All activities: REI is 12 hours. Pre-grazing Interval (PGI): 14 days.

Beans (Pole, Snap)

Aphids Grasshoppers Leafhoppers	700 mL - 1 L	7 - 10 mL	7	2
---------------------------------------	--------------	-----------	---	---

2023-7083  
2024-05-31

Leafminers  
Mexican bean beetles  
Mites, Lygus bugs  
Tarnished plant bugs

COMMENTS: Use sufficient water for good coverage. Do not feed or graze treated foliage to livestock. May be applied by Ground and Aerial application. Minimum application interval is 7 days. All activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period.

Beets	Tarnished plant bugs	700 mL	7 mL	12	2
COMMENTS: Ground application ONLY. Minimum application interval is 7 days. All activities: REI is 3 days. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period.					

Cauliflower Broccoli Brussels Sprouts	Aphids	700 mL - 1 L	7 - 10 mL	7 Broccoli Cauliflower 21 Brussels Sprouts	2
---	--------	--------------	-----------	--	---

COMMENTS: To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period. Use sufficient water for good coverage. Ground Application ONLY. Minimum application interval is 7 days. For all activities: REI is 5 days.

Cauliflower Broccoli	Thrips	550 mL - 1.25 L	6 - 12 mL	7	2
To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period. Apply at head formation. Ground Application ONLY. Minimum application interval is 7 days. All activities: REI is 5 days.					

Canary Grass (Grown for seed)	Aphids	500 mL	5 mL	21	2
COMMENTS: Apply in aerial or ground application when there are more than 50 aphids per canary seed head between heading and soft dough stage. May be applied by Ground and Aerial application. May be applied a maximum of 2 times per year. Minimum application interval is 30 days. All activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period.					

Canola (rape)	Aphids Leafhoppers Grasshoppers	850 - 900 mL	9 mL	21	2
---------------	---------------------------------------	--------------	------	----	---

COMMENTS: May be applied by ground and aerial application. Minimum application interval is 7 days. All activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.

Celery	Aphids, including carrot green peach foxglove aphids	700 mL	7 mL	7	2
--------	---	--------	------	---	---

COMMENTS: Begin application 3 weeks after transplanting. Ground application ONLY. Minimum application interval is 7 days. All Activities, REI is 3 days. To protect pollinators, follow the instructions

2023-7083  
2024-05-31

regarding bees in the Environmental Precautions section. TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period.

Cereals (barley, oats, wheat, rye)	Grasshoppers	550 mL	5 mL	35	2
	-nymphs, Say stink bugs				
	Grasshoppers	850 mL - 1 L	See comments		2
	-adults				
<p>COMMENTS: Do not graze within 14 days after treatment. Do not harvest for forage within 35 days after treatment. Follow provincial forecast. Apply when the grasshoppers are in the 2-4 nymphal stage. Best control will be achieved when application is made prior to wing development. Under severe insect pressure application should also be made to a 15-metre strip along fence rows around the field. DO NOT APPLY TO ADJACENT UNREGISTERED CROPS. The higher rate should be used when the proportion of mature and late nymphal stages in the population are high and spray penetration is inhibited by dense crop canopy. May be applied by Ground and Aerial application. Minimum application interval is 7 days. All activities: REI is 12 hours.</p>					
Chicory,	Aphids	600 mL - 1 L	6 - 10 mL	7	2
Chinese Broccoli					
Bok Choi	<p>COMMENTS: Apply in sufficient water for thorough coverage. Use ground application to foliage. Apply when aphids are present. Ground application ONLY. Minimum application interval is 7 days. All activities: REI is 4 days. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period.</p>				
Eggplant	Tarnished plant bug	500 - 700 mL	5 - 7 mL	7	2
<p>COMMENTS: Apply when bugs are found and flowering is seen. Treat again in 7-10 days if bugs are found. Consult local authorities for proper timing. Do not apply when bees are foraging. Ground application ONLY. Minimum application interval is 7 days. All activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period.</p>					
Flax	Potato Aphid	427 mL	4 mL	21	1
<p>COMMENTS: Apply from late flowering to early green bole stage; in sufficient water to provide good coverage. May be applied by either ground or aerial methods. Use in 200 L/ha of water for ground and 11 - 22 L of water/ha for aerial equipment. All activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period.</p>					
Forage crops (grasses, alfalfa, hay, clover)	Aphids	425 - 550 mL	4 - 5 mL	2	2
	(Only suppression of Russian Wheat aphid is provided)				
	Grasshoppers-				
	low infestations	425 – 550 mL		2	2
	nymphs	550 mL		2	2
adults	850 mL – 1L		28	2	
	Lygus Bug	425 mL		2	2
	Plant Bug	425 mL		2	2
<p>COMMENTS: TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms. Do not graze or harvest for forage within 2 days after treatment at the 425 or 550 mL rate. Do not graze or harvest for forage within 28 days after treatment at the 850 mL – 1 L rate. May be applied by either ground or aerial methods. Minimum application interval is 7 days. All activities: REI is 12 hours.</p>					
Leafy Vegetables: Lettuce, Swiss chard,	Aphids	700 mL	7 mL	14	2
	Leafhoppers				

2023-7083  
2024-05-31

Beet greens and Turnip greens COMMENTS: Ground application ONLY. Use sufficient water for good coverage. Do not use on Head Lettuce. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period. Minimum application interval is 7 days. Interval between last application and harvest:

Lettuce	14 days	Beet greens	21 days
Swiss chard	14 days	Turnip greens	14 days

Beet Greens, Turnip Greens and Lettuce: All Activities, REI is 3 days.

Kale	Aphids	700 mL	7 mL	14	2
	Leafhoppers				

COMMENTS: Ground application ONLY. Use sufficient water for good coverage. May be applied a maximum of 2 times per year. Minimum application interval is 15 days. Interval between last application and harvest: 14 days. All Activities, REI is 3 days. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period.

Peas (canning & field)	Aphids	275 - 380 mL	2.5 - 3.6 mL	3	2
---------------------------	--------	--------------	--------------	---	---

COMMENTS: Apply by aircraft or by ground equipment. Use sufficient water for good coverage. Do not feed or graze for forage within 21 days after last application. Dilute for ground (100-300 L of water/ha) and aerial application (10-30 L of water/ha). Minimum application interval is 14 days. Maximum application rate is 0.18 kg a.i./ha. All activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period.

Peppers	Aphids, Maggots	700 mL - 1L	7 - 10 mL	3	2
---------	-----------------	-------------	-----------	---	---

COMMENTS: Use sufficient water for good coverage. Ground application ONLY. Minimum application interval is 7 days. All activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period.

Potatoes, Tomatoes (field)	Aphids	550 mL - 1.0 L	5 - 9 mL	7	2
	Leafhoppers				
	Tarnished plant bugs				

COMMENTS: Use sufficient water for good coverage. Do not apply more than 2 applications per season. Consult local agricultural authorities for proper timing. Ground application ONLY. Minimum application interval is 7 days. Maximum application rate is 0.48 kg a.i./ha All activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period.

Pastures Wastelands	Grasshoppers				
	-nymphs	550 mL		2	2
	-adults	850 mL - 1 L		See Comments	

COMMENTS: Follow provincial forecast. Apply when insects are present as young hoppers or signs of insect damage are evident. Apply when the grasshoppers are in the 2-4 nymphal stage. Best control will be achieved when application is made prior to wing development. Under severe insect pressure application should also be made to a 15-metre strip along fence rows around the field. DO NOT APPLY TO ADJACENT UNREGISTERED CROPS. The higher rate should be used when the proportion of mature and late nymphal stages in the population are high and spray penetration is inhibited by dense crop canopy. Use high rate for adult grasshopper control on wasteland. Remove cattle before spraying. Do not graze or harvest for forage within 2 days after treatment at the 550 mL rate. Do not graze or harvest for forage within 7 days after treatment at the 850 mL rate. Do not graze or harvest for forage within 28 days at the 1L rate. May be applied by ground and aerial application. Minimum application interval is 7 days. All activities: REI is 12 hours. To protect pollinators, follow the

2023-7083  
2024-05-31

instructions regarding bees in the Environmental Precautions section. TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period.

Safflower	Grasshoppers (nymphs and adults)	550 mL - 1L		21	2
<p>COMMENTS: Apply when damage is apparent and more than 15 grasshoppers per square metre are found in the crop. Repeat the application only when necessary. Use sufficient water for good coverage. May be applied by ground and aerial application. Minimum application interval is 7 days. All activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.</p>					
Soybeans	Aphids bean beetles leafhoppers tarnished plant bugs, lygus bugs	0.7 - 1 L	7 - 10 mL	30	2
<p>COMMENTS: Use sufficient water for good coverage. Do not exceed 2 applications per season. Do not feed or graze treated foliage to livestock. May be applied by Ground and Aerial application. Minimum application interval is 7 days. All activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period.</p>					
	Spider mites (Twospotted)	1 L	10 mL	30	2
<p>COMMENTS: Apply by ground application in sufficient water for thorough coverage. Do not feed or allow livestock to graze treated forage. May be applied by ground and aerial application. Minimum application interval is 7 days. All activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period.</p>					
Sweet Clover Red Clover Alsike Clover	Sweet clover weevil	850 mL - 1.1 L		28	2
<p>COMMENTS: Do not graze or harvest for forage within 28 days after treatment. May be applied by ground and aerial application. Minimum application interval is 7 days. All activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.</p>					
Wheat (ALSO SEE CEREALS above)	Wheat midge, Russian wheat aphid	1 L		21	2
<p>COMMENTS: If midge population persists at 7 days apply a second treatment. Use a water volume of at least 100 L/ha with ground equipment and 50 L/ha by air. Higher volume will provide more thorough coverage. If adult midges are present (1 midge/4-5 wheat heads), sprays should be applied when 25% of the wheat head has fully emerged from the boot but before flowering has begun. At this stage, wheat first becomes susceptible to attack by egg-laying females. Applications should be made in the late afternoon or evening when temperatures exceed 15°C and the wind speed is less than 10 km/h. Higher volume sprays will improve penetration of the crop. Proper timing of application is essential for control. Do not harvest within 21 days of the application. Do not apply to areas where bees are actively foraging or near apiaries as product is toxic to bees. Do not graze or use for forage within 7 days of application. Consult provincial authorities for further information concerning rates and time of application. Minimum application interval is 7 days. All activities: REI is 12 hours.</p>					

**FRUIT CROPS:**



2023-7083

2024-05-31

APPLY WITH GROUND EQUIPMENT ONLY UNLESS OTHERWISE NOTED. FOR DILUTE GROUND APPLICATIONS, Apply at the recommended rate; do not exceed a maximum spray volume of 1000 L per hectare unless otherwise stated. For Cherry, Peach and Pear, the rate per hectare is based on using 3000 L of spray on full sized trees, 4.5-5.5 metres high. Apply at the recommended rate; do not exceed a maximum spray volume of 3000 L per hectare unless otherwise stated. Trees smaller or larger may require more or less LAGON 480 E. Follow provincial calendar guidelines as to water volumes for the size of tree being sprayed.

FRUIT TREE CROPS	PEST CONTROLLED	AMOUNT PER HECTARE	AMOUNT PER 10 L	AMOUNT PER 1000 L	PREHARVEST INTERVALS	MAXIMUM # OF APPLICATIONS PER YEAR
Pears	Tarnished plant bugs		6.25 mL	625 mL	28	2
	COMMENTS: Pears: Apply at pre-bloom. The minimum application interval is 10 days. Maximum rate 1.44 kg a.i./ha. Thinning: REI is 28 days. Hand harvesting, hand-line irrigation: REI is 17 days. All other activities: REI is 1 day. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.					
	Pear Psylla		10 – 12.5 mL	1 – 1.25 L	28	2
	COMMENTS: Use sufficient water for good cover spray. The minimum application interval is 10 days. Maximum application rate is 1.44 kg a.i./ha. Thinning: REI is 28 days. Hand harvesting, hand-line irrigation: REI is 17 days. All other activities: REI is 1 day. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.					
Peaches (non-bearing)	Aphids		10 – 12.5 mL	1 – 1.25 L	28	2
	COMMENTS: Use sufficient water for good cover spray The minimum application interval is 10 days. Maximum application rate is 1.44 kg a.i./ha. Thinning: REI is 28 days. Hand harvesting, hand-line irrigation: REI is 17 days. All other activities: REI is 1 day. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.					
	Mites		10 – 12.5 mL	1 – 1.25 L	28	2
	COMMENTS: Use sufficient water for good cover spray The minimum application interval is 10 days. Maximum application rate is 1.44 kg a.i./ha. Thinning: REI is 28 days. Hand harvesting, hand-line irrigation: REI is 17 days. All other activities: REI is 1 day. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.					
Peaches (non-bearing)	Aphids, Mites		10 - 15 mL	1 – 1.5 L	40	2
	Tarnished plant bugs	1.7 L	5 mL	500 mL	40	2
COMMENTS: Spray when insects first appear using sufficient water for good coverage. Some defoliation may occur under cold wet soil and/or slow drying conditions during and following application. Least injury has occurred when sprays were applied in the morning while temperatures are moderate and drying conditions are fairly good. The minimum application interval is 10 days. Thinning: REI is 32 days. Hand harvesting, hand-line irrigation: REI is 20 days. All other activities: REI is 3 days. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.						
Sour Cherries	Cherry fruit fly maggots	2.25 L	5 mL	500 mL	21	2
	COMMENTS: Apply no later than 6 days after the first adult fly has been trapped and repeat in 14 days if required. Thinning: REI is 20 days. Hand harvesting, hand-line irrigation: REI is 9 days. All other activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.					
Sweet Cherries	Cherry fruit fly maggots	2.25 L	5 mL	500 mL	21	2
COMMENTS: Apply no later than 6 days after the first adult fly has been trapped and repeat in 14 days if required. Thinning: REI is 20 days. Hand harvesting, hand-line irrigation: REI is 9 days. All other activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.						

SMALL FRUIT CROPS	PESTS CONTROLLED	AMOUNT PER HECTARE	AMOUNT PER 10 LITRES	AMOUNT PER 1000 LITRES	PREHARVEST INTERVALS (DAYS)	MAXIMUM # OF APPLICATIONS PER YEAR
Blueberry Lowbush Highbush	Blueberry maggot	580 – 830 mL	6 – 8 mL		21	2
	<p>COMMENTS: Apply pesticide in sufficient water for good coverage, one week after first berries ripen. Use no more than 1000 L spray/ha. Make 1 or 2 applications between mid to late July. The minimum application interval is 10 days. May vary depending on area and season. May be applied with ground equipment or aerial equipment. Hand harvesting (high bush): REI is 9 days. Hand harvesting (low bush) and all other activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.</p>					
Strawberries (bearing)	Tarnished plant bug	2.75 L	12.5 mL	1.25 L	7	2
	<p>COMMENTS: Apply first spray when first blooms appear and the second application 10-12 days later if needed. May be applied a maximum of 2 times per year. Minimum application interval is 10 days. All Activities, REI is 2 days. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period.</p>					
Strawberries (bearing & non-bearing)	Aphids, mites	2.25 L	10 mL	1 L	7	2
	<p>COMMENTS: Use sufficient water for good cover spray. Spray when insects first appear. May be applied a maximum of 2 times per year. Minimum application interval is 10 days. All Activities, REI is 2 days. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period.</p>					

CROPS	PEST	RATE PER HECTARE	REMARKS
Filberts, Hazelnuts	Aphids	5 L	<p>Use one application per season. Apply when aphids appear. DO NOT USE WITHIN 45 DAYS OF HARVEST. Primarily for use on young plantings Thinning: REI is 34 days. Hand harvesting, hand-line irrigation: REI is 21 days. All other activities: REI is 5 days. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period.</p>

**ORNAMENTALS**

GENERAL COMMENTS ORNAMENTAL PLANTS

APPLY ONLY WITH GROUND APPLICATION EQUIPMENT

- LAGON 480 E is a true systemic insecticide. A systemic insecticide is absorbed into the system of the plant upon application, and as with all systemic materials, may cause reactions in specific plants which are not predictable nor common to all members of the species. Do not over-dose or over-spray.
- FOR PROPER TIMING of treatments for the control of specific pests on ornamental plants, consult local agricultural authorities. In general, apply adequate spray for good coverage when pests appear or when damage is first observed, unless otherwise indicated.
- EXPERIENCE has shown when dealing with species that may be injured by foliar application, soil drenches are safer on those plants for which they are recommended. When using foliar sprays, do not apply during heat of the day or when temperatures are excessively high.

2023-7083  
2024-05-31

4. DO NOT USE on ornamental plants not listed.
5. WHEN IN DOUBT about plant safety, it is advisable to spray a single branch or dip several leaves in the spray solution. A waiting period of 3 or 4 days will usually indicate if plant damage will result.
6. DO NOT SPRAY the following species unless you have tested single branches or leaves to ascertain safety:
 

American Elm	Mop-head Elm
Norway Maple	Mock Orange
Sunburst Locust	Morraine Locust

 or other ornamentals not listed in this pamphlet.
7. There has been no report of damage by LAGON 480 E to any of the common evergreen species grown in Canada provided recommended rates have been followed.
8. Do not apply foliar spray during the heat of the day or when temperatures are exceedingly high.
9. Apply at the recommended rate; do not exceed a maximum spray volume of 1000 L per hectare unless otherwise stated.

OUTDOOR FLOWERING PLANTS	PESTS CONTROLLED	AMOUNTS PER 1000 L OF WATER	AMOUNTS PER 10 L OF WATER
Azaleas	Lace bugs, Leafminers, Mites Whiteflies COMMENTS: The minimum application interval is 14 days. May be applied a maximum of 2 times per year. All Activities, REI is 2 days. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms..	1 L	10 mL
Camellias	Aphids, Mites, Camellia scales FOLIAR SPRAYS: Apply 2 sprays 6 weeks apart the first year followed by annual applications soon after first growth begins in the spring. The minimum application interval is 10 days. May be applied a maximum of 2 times per year. All Activities, REI is 2 days. SOIL DRENCH: Apply LAGON 480 E as a soil drench around the base of plants in early spring at the rate of 60 mL per 10 L of water per plant up to 2 metres tall. Increase this rate proportionately for larger plants. Soil drench: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms..	1 L	10 mL
Day Lilies	Aphids, Thrips COMMENTS: May be applied a maximum of 1 time per year. All activities: REI 7 days. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms..	2 L	20 mL
Euonymus	Aphids, Scales COMMENTS: The minimum application interval is 14 days. May be applied a maximum of 2 times per year. All activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms..	2 L	20 mL
Gardenias	Tea scales, Whiteflies COMMENTS: The minimum application interval is 14 days. May be applied a maximum of 2 times per year. All activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms..	1 L	10 mL
Gerberas	Thrips COMMENTS: May be applied a maximum of 1 time per year. All Activities, REI is 2 days. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms..	1 L	10 mL
Gladiolus	Aphids, Thrips COMMENTS: May be applied a maximum of 1 time per year. All Activities, REI is 2 days. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms..	1 L	10 mL
Iris	Aphids, Iris Borers, Thrips For borer control spray when new leaves are 12 to 15 cm tall. COMMENTS: May be applied a maximum of 1 time per year. All activities: REI is 7 days. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms..	2 L	20 mL

2023-7083  
2024-05-31

Poinsettias (outdoor plants)	Aphids, Whiteflies, Mites, Mealybugs	1 L	10 mL
COMMENTS: May be applied a maximum of 1 time per year. All Activities, REI is 2 days. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms..			
Roses (outdoor plants)	Aphids, Leafhoppers, Mites, Thrips	1 L	10 mL
Apply thoroughly during growing season in a spray. The minimum application interval is 28 days. May be applied a maximum of 2 times per year. All Activities, REI is 2 days. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms..			

OUTDOOR ORNAMENTALS AND CHRISTMAS TREES	PESTS CONTROLLED	AMOUNTS PER 1000 L OF WATER	AMOUNTS PER 10 L OF WATER
Christmas Trees: Balsam Fir, Spruce, Hemlock, Pine (Mugho, Red, Scots)	Aphids, mites, Scales, Spruce needleminers, Spruce budworms	1.5 L	15 mL
	COMMENTS: The minimum application interval is 14 days. May be applied a maximum of 2 times per year. Thinning: REI is 18 days. Hand line irrigation: REI is 3 days. All other activities: 12 hours.		
Arborvitae	Bagworms, European pine shoot moths, Nantucket pine tip moths, Zimmerman pine moths	2 L	20 mL
	COMMENTS: The minimum application interval is 14 days. May be applied a maximum of 2 times per year. Thinning: REI is 18 days. Hand line irrigation: REI is 3 days. All other activities: REI is 12 hours.		
Birch	Aphids, Bagworms, Mites	2 L	20 mL
COMMENTS: The minimum application interval is 14 days. May be applied a maximum of 2 times per year. All activities: REI is 12 hours.			
Birch	Aphids, Birch Leafminers	500 mL	5 mL
	FOLIAR SPRAY: For leaf miners apply when leaves are expanded (about mid-May); repeat the last week in June. For all activities: REI is 12 hours.		
	SOIL DRENCH: For full-season control LAGON 480 E may be used as a soil drench once in early May when the leaves start to open. Apply LAGON 480 E undiluted at a rate of 11 mL/cm of stem basal diameter (cumulative total of all stems) evenly into small shallow holes made with a sharp instrument at the dripline of the tree or clump. Close the holes with soil. Drench thoroughly with water immediately so as to saturate the root system. One soil drench treatment in May should give control of birch leafminers for the complete season. Do not make more than one soil drench treatment per season. Soil drenches may not be effective on newly-transplanted birches where the root system is not well established. Some hybrid varieties eg: cutleaf and silver birch may be damaged by excessive amounts of LAGON. Therefore, apply no more than the recommended dosage. Soil drench: REI is 12 hours. COMMENTS: To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.		
Boxwood	Mites, Leafminers, Mealybugs	1 L	10 mL
COMMENTS: For leafminers, apply in spring when leafminer flies first appear or in early summer for control of larvae in the infested leaves. The minimum application interval is 14 days. May be applied a maximum of 2 times per year. Thinning: REI is 7 days. All other activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.			
Cedar	Mites	2 L	20 mL
COMMENTS: The minimum application interval is 14 days. May be applied a maximum of 2 times per year. Thinning: REI is 18 days. Hand-line irrigation: REI is 3 days. All other activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions			

2023-7083  
2024-05-31

section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.

Douglas-Fir (seed tree)	Cone insects (cone or gall midges, cone moths, seed chalcids, scale midges). 2 L per 100 litres of water. Maximum rate 20 L of product (= 9.6 kg a.i.) per hectare. Application should be made when cones are at or near the pendant stage. Complete coverage of the cones and foliage in the cone-bearing area of the tree is important for satisfactory results. Maximum of two applications per year. Minimum application interval: 10 days Seed cone harvesting: 48 days REI. Scouting: 5 days REI. Grading, animal control, baiting: 1 day REI.		
Hemlock	Mites, Scales, Spruce needleminers Spruce budworms COMMENTS: The minimum application interval is 14 days. May be applied a maximum of 2 times per year. Thinning: REI is 13 days. All other activities: REI is 12 hours.	1 L	10 mL
Holly (English & American)	Mites, Soft Scale, Leafminers  COMMENTS: For leafminers apply in spring when leafminer flies first appear, or in early summer for the control of larvae in the infested leaves. The minimum application interval is 14 days. May be applied a maximum of 2 times per year. All activities: REI is 12 days. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.	1 L	10 mL
Juniper	Aphids, Bagworms, Midges, Mites COMMENTS: The minimum application interval is 14 days. May be applied a maximum of 2 times per year. All activities: REI is 12 hours.	2 L	20 mL
Larch	Cone Maggots, Seed Wasps  COMMENTS: One application per season; apply within 3 weeks after pollination ; spray to runoff on foliage; ground hydraulic sprayer or mist sprayer. All Activities, REI is 1 day. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.	5 L	50 mL
Lilacs (Syringa spp.)	Leafminers  COMMENTS: Apply two sprays 6 weeks apart. Application should be made as soon as any signs of the leafminer appear, or, first application should be started in early June. The minimum application interval is 10 days. May be applied a maximum of 2 times per year. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms. All activities: REI is 15 days.	1.25 L	12.5 mL
Oak	Golden oak scales COMMENTS: The minimum application interval is 14 days. May be applied a maximum of 2 times per year. Thinning: REI is 18 days. Hand-line irrigation: REI is 3 days. All other activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.	2 L	20 mL
Pine (Mugho, Red, Scots)	Aphids, Bagworms, European pine shoot moths, Nantucket pine tip moths, Zimmerman pine moths  Red and black-headed pine sawfly larvae  COMMENTS: For pine shoot moths apply early in the spring and again in mid-June, thoroughly wetting branch tips. COMMENTS: The minimum application interval is 14 days. May be applied a maximum of 2 times per year. Thinning: REI is 18 days. Hand line irrigation: REI is 3 days. All other activities: REI is 12 hours.	2 L  1 L	20 mL  10 mL
Sitka spruce (seed tree):	Sitka spruce weevils. 20.8 L per 1000 litres of water. Maximum rate 20.8 L of product (= 10 kg a.i.) per hectare. Thoroughly spray the terminal growth to the point of runoff at the time of egg laying (usually during the first half of May). Maximum of two applications per year. Minimum application interval: 10 days. Seed cone harvesting: 49 days REI. Scouting: 5 days REI. Hand-pruning, staking, tying: 27 days REI. Grading, animal control, baiting: 1 day REI.		
Spruce (seed tree and woodland use):	Seed and cone insects. 10.4 - 20.8 L per 1000 L water. Maximum rate 20.8 L of product (10 kg a.i.) per hectare. Thoroughly spray cone-bearing portions of tree to point of runoff when strobili cones are closed, and turning, but before they reach the horizontal position. Contact your local forest		

2023-7083  
2024-05-31

pest control office for more information on timing, and spray application, as well as spruce species that may be treated. Maximum of two application per year. Minimum application interval: 10 days. Seed cone harvesting: 49 days REI. Scouting: 5 days REI. Hand pruning, staking, tying: 27 days REI. Grading, animal control, baiting: 2 days REI.

Taxus	Mites, Fletcher scale, Mealybugs	2 L	20 mL
-------	----------------------------------	-----	-------

COMMENTS: For Fletcher scale, spray at the end of June and again 14 days later when crawlers are moving. May be applied a maximum of 2 times per year. Minimum application interval is 14 days. All activities: REI is 12 hours.

**NOTE TO USER: READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR THE INDICATED SPECIAL USE APPLICATIONS:**

The DIRECTIONS FOR USE for the uses described in this section of the label were developed by persons other than Loveland Products Canada Inc. under the User Requested Minor Use Label Expansion program. For these uses, Loveland Products Canada Inc. has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.

**Highbush Blueberry**

Crop	Pest	Rate per 1000 L water	Maximum No. of Applications	Reapplication Interval
Highbush Blueberry	Spotted-wing Drosophila	830 mL	2	15 days

COMMENTS: Apply post-harvest only, to control spotted-wing Drosophila (adults and larvae in fruit) that may otherwise infest adjacent crops. Timing varies, depending on variety of highbush blueberry. Foliar application only, using conventional ground application equipment. Use sufficient water volume to ensure thorough coverage, to a maximum of 1000 L/ha. REI is 12 days.

2023-7083  
2024-05-31

**RESTRICTED USES - FORESTRY USE**

**NOTICE TO USER:** This pest control product is to be used only in accordance with the directions on the label. It is an offence under the Pest Control Product Act to use this product in a way that is inconsistent with the directions on the label.  
**CAUTION:** Forest Pest Control - birds and other wildlife in treated area may be harmed.  
Apply at the recommended rate; do not exceed a maximum spray volume of 1000 L per hectare unless otherwise stated.

**RESTRICTED USE - AERIAL APPLICATION**

TREE	PESTS CONTROLLED	COMMENTS
Hemlock, Spruce, Balsam fir	Mites, Scales, Spruce needleminers, Spruce budworms	For aircraft application use 1.5 L of LAGON 480 E in 5-20 L of water/ha, applied as a uniform spray of fine droplets. Application should be made as soon as the larvae hatch. COMMENTS: May be applied a maximum of 1 time per year. Thinning: REI is 13 days. All other activities: REI is 12 hours.

**NOTE TO USER: READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR THE INDICATED SPECIAL USE**

**APPLICATIONS:**

The DIRECTIONS FOR USE for the uses described in this section of the label were developed by persons other than Loveland Products Canada Inc. under the User Requested Minor Use Label Expansion program. For these uses, Loveland Products Canada Inc. has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.

**ANNUAL CANARYGRASS**

CROP	PEST	RATE PER HA	MAXIMUM NO. OF APPLICATIONS PER YEAR	PREHARVEST INTERVALS (DAYS)	AMOUNT PER 10 LITRES
Annual canary grass	Aphids	500 mL	1	21	5 mL

COMMENTS: Apply in aerial or ground application when there are more than 50 aphids per canary seed head between heading and soft dough stage. May be applied by Ground and Aerial application.

May be applied a maximum of 1 time per year. All activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period.

Annual canarygrass seeds can be harvested for human consumption.

Annual canarygrass grown for human consumption is not to be cut for feed or grazed.

**DECONTAMINATION**

If accidental spillage of LAGON 480 E Insecticide should occur, scrub contaminated area immediately with a strong laundry soap solution or use household lye. Detergents are not satisfactory for this purpose. Repeated scrubbing are necessary on plain wood surfaces. Avoid spillage on all types of floor tile or linoleum.

**FIRST AID**

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

**IF SWALLOWED:** Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by poison control centre or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferable by mouth-to-mouth, if possible. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

2023-7083  
2024-05-31

### **TOXICOLOGICAL INFORMATION**

Dimethoate is an organophosphate that is a cholinesterase inhibitor. Typical symptoms of overexposure to cholinesterase inhibitors include headache, nausea, dizziness, sweating, salivation, and runny nose and eyes. This may progress to muscle twitching, weakness, tremor, incoordination, vomiting, abdominal cramps and diarrhea in more serious poisonings. A life-threatening poisoning is signified by loss of consciousness, incontinence, convulsions and respiratory depression with a secondary cardiovascular component. Treat symptomatically. If exposed, plasma and red blood cell cholinesterase tests may indicate degree of exposure (baseline data are useful). Atropine, only by injection, is the preferable antidote. Oximes, such as pralidoxime chloride, may be therapeutic if used early; however, use only in conjunction with atropine. In cases of severe acute poisoning, use antidotes immediately after establishing an open airway and respiration. With oral exposure, the decision of whether to induce vomiting or not should be made by an attending physician. This product contains petroleum distillates. Vomiting may cause aspiration pneumonia. Treat symptomatically.

### **STORAGE**

Store this product away from food or feed.  
Do not use, pour, spill, or store near heat, sparks, or open flame. Keep out of direct sunlight. Ship and store between 4 30°C.

### **DISPOSAL**

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

**NOTICE TO USER:** This pest control product is to be used only in accordance with the directions on this label. It is an offense under the *PEST CONTROL PRODUCTS ACT* to use this product in a way that is inconsistent with the directions on the label.