

(Container)



# 2,4-D BEE-4

## Herbicide

GROUP	4	HERBICIDE
-------	---	-----------

- Emulsifiable weed killer.
- Contains 2,4-D.
- For air or ground application to wheat, barley, rye, corn, pastures and turf, for the control of certain annual and perennial weeds and for woody plant control.

AGRICULTURAL

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING  
KEEP OUT OF REACH OF CHILDREN

GUARANTEE: 2,4-D, present as butoxyethyl ester      500 g a.e./L

REGISTRATION NO. 6330 PEST CONTROL PRODUCTS ACT

**WARNING**  **POISON**

NET CONTENTS: 20 L

**Corteva Agriscience Canada Company**  
Suite 240, 115 Quarry Park Rd. SE  
Calgary, AB  
T2C 5G9  
1-800-667-3852

™ Trademark of Corteva Agriscience and its affiliated companies.

**PRECAUTIONS**  
**MAY CAUSE SKIN IRRITATION**  
**AVOID CONTACT WITH EYES, SKIN AND CLOTHING**  
**AVOID BREATHING SPRAY MIST**  
**KEEP OUT OF REACH OF CHILDREN**

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at [www.croplife.ca](http://www.croplife.ca).

This product contains a petroleum distillate which is moderately to highly toxic to aquatic organisms. Avoid contamination of aquatic systems during application. Do not contaminate these systems through direct applications, disposal of waste or cleaning equipment.

**PROTECTIVE CLOTHING AND EQUIPMENT**  
**All Application Equipment (Ground and Aerial)**

Closed mixing/loading systems are required.

- When mixing/loading, wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, shoes and protective eye wear (face shield or safety glasses). Rinse gloves before removal.
- When applying or during clean-up and repair, wear coveralls over a long sleeved shirt, long pants, chemical resistant gloves, socks and shoes. Rinse gloves before removal.
- Gloves are not required during application when applicator is in an enclosed tractor or in an enclosed airplane cockpit.

**Application using Aerial Equipment**

- Applicators must wear coveralls over a long-sleeved shirt and long pants. Chemical-resistant gloves must also be worn during clean-up and repair activities.
- No human flaggers are permitted.

**Application using Groundboom Equipment**

- Applicators must wear coveralls over a long-sleeved shirt and long pants. Chemical-resistant gloves must also be worn during clean-up and repair activities.
- When handling more than 340 L per day workers must also use a closed cab.

**Application using Handheld Equipment**

Applicators must wear coveralls over a long-sleeved shirt, long pants, respirator and chemical-resistant gloves. Solution for annual and broadleaf plant control must have a minimum dilution volume of 100L/ha.

Refer to Directions for Use section for any additional requirements.

**OPERATOR USE PRECAUTIONS**

- Wear freshly laundered clothing and clean protective equipment daily.
- Rinse gloves before removal.
- Wash hands before eating, drinking, using tobacco or using the toilet.
- If herbicide penetrates clothing, remove immediately; then wash thoroughly and put on clean clothing. Throw away clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate.
- After using this product, remove clothing and launder separately, and promptly and thoroughly wash hands and exposed skin with soap and water. Follow manufacturer's instructions for

cleaning personal protective clothing and equipment. If no such instructions for washables are provided, use detergent and hot water. Keep and wash personal protective equipment separate from household laundry.

- After work, remove all clothing and shower using soap and water.

#### **FIRST AID**

**Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.**

**If swallowed:** Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

#### **TOXICOLOGICAL INFORMATION**

No specific antidote. Employ supportive care. Treatment should be based on judgement of the physician in response to reactions of the patient. This product may cause mild irritation to the eyes. Overexposure to 2,4-D may cause coughing, burning, dizziness or temporary loss of muscle coordination. Other possible effects of overexposure include fatigue, muscle weakness or nausea. This product contains petroleum distillates. Vomiting may cause aspiration pneumonia. Treat Symptomatically.

#### **AGRICULTURAL CHEMICAL**

Do not ship or store with food, feeds, drugs or clothing.

#### **ENVIRONMENTAL HAZARDS**

- Toxic to small mammals, birds, aquatic organisms and non-target terrestrial plants.
- Observe buffer zones specified under DIRECTIONS FOR USE.
- This product will harm other broadleaved plants in the vicinity of the treatment area. If applying this product using a handheld sprayer, do not directly spray or allow the spray to drift onto ornamentals or gardens.
- Do not spray exposed roots of trees and ornamentals.

#### **LEACHING**

The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g. sandy soil) and/or the depth to the water table is shallow.

#### **RUNOFF**

- To reduce runoff from treated areas into aquatic habitats avoid application to areas with moderate to steep slope, compacted soil, or clay.
- Avoid application of this product when heavy rain is forecast.
- Contamination of aquatic areas as a result of runoff may be reduced by including a strip of untreated vegetation between the treated area and the edge of the water body.
- To prevent runoff from domestic uses avoid spraying on driveways, sidewalks or other hard surfaces. Do not irrigate within 24 hours after application.

**DISPOSAL**

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

**NOTICE TO USER:** This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

(Booklet)



# 2,4-D BEE-4

## Herbicide

GROUP	4	HERBICIDE
-------	---	-----------

- Emulsifiable weed killer.
- Contains 2,4-D.
- For air or ground application to wheat, barley, rye, corn, pastures and turf, for the control of certain annual and perennial weeds and for woody plant control.

AGRICULTURAL

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING  
KEEP OUT OF REACH OF CHILDREN

GUARANTEE: 2,4-D, present as butoxyethyl ester      500 g a.e./L

REGISTRATION NO. 6330 PEST CONTROL PRODUCTS ACT

**WARNING**  **POISON**

NET CONTENTS: 20 L

**Corteva Agriscience Canada Company**  
Suite 240, 115 Quarry Park Rd. SE  
Calgary, AB  
T2C 5G9  
1-800-667-3852

™Trademark of Corteva Agriscience and its affiliated companies.

**PRECAUTIONS**  
**MAY CAUSE SKIN IRRITATION**  
**AVOID CONTACT WITH EYES, SKIN AND CLOTHING**  
**AVOID BREATHING SPRAY MIST**  
**KEEP OUT OF REACH OF CHILDREN**

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at [www.croplife.ca](http://www.croplife.ca).

This product contains a petroleum distillate which is moderately to highly toxic to aquatic organisms. Avoid contamination of aquatic systems during application. Do not contaminate these systems through direct applications, disposal of waste or cleaning equipment.

**PROTECTIVE CLOTHING AND EQUIPMENT**  
**All Application Equipment (Ground and Aerial)**

Closed mixing/loading systems are required.

- When mixing/loading, wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, shoes and protective eye wear (face shield or safety glasses). Rinse gloves before removal.
- When applying or during clean-up and repair, wear coveralls over a long sleeved shirt, long pants, chemical resistant gloves, socks and shoes. Rinse gloves before removal.
- Gloves are not required during application when applicator is in an enclosed tractor or in an enclosed airplane cockpit.

**Application using Aerial Equipment**

- Applicators must wear coveralls over a long-sleeved shirt and long pants. Chemical-resistant gloves must also be worn during clean-up and repair activities.
- No human flaggers are permitted.

**Application using Groundboom Equipment**

- Applicators must wear coveralls over a long-sleeved shirt and long pants. Chemical-resistant gloves must also be worn during clean-up and repair activities.
- When handling more than 340 L per day workers must also use a closed cab.

**Application using Handheld Equipment**

Applicators must wear coveralls over a long-sleeved shirt, long pants, respirator and chemical-resistant gloves. Solution for annual and broadleaf plant control must have a minimum dilution volume of 100L/ha.

Refer to Directions for Use section for any additional requirements.

**OPERATOR USE PRECAUTIONS**

- Wear freshly laundered clothing and clean protective equipment daily.
- Rinse gloves before removal.
- Wash hands before eating, drinking, using tobacco or using the toilet.
- If herbicide penetrates clothing, remove immediately; then wash thoroughly and put on clean clothing. Throw away clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate.
- After using this product, remove clothing and launder separately, and promptly and thoroughly wash hands and exposed skin with soap and water. Follow manufacturer's instructions for

cleaning personal protective clothing and equipment. If no such instructions for washables are provided, use detergent and hot water. Keep and wash personal protective equipment separate from household laundry.

- After work, remove all clothing and shower using soap and water.

#### **FIRST AID**

**Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.**

**If swallowed:** Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

#### **TOXICOLOGICAL INFORMATION**

No specific antidote. Employ supportive care. Treatment should be based on judgement of the physician in response to reactions of the patient. This product may cause mild irritation to the eyes. Overexposure to 2,4-D may cause coughing, burning, dizziness or temporary loss of muscle coordination. Other possible effects of overexposure include fatigue, muscle weakness or nausea. This product contains petroleum distillates. Vomiting may cause aspiration pneumonia. Treat Symptomatically.

#### **AGRICULTURAL CHEMICAL**

Do not ship or store with food, feeds, drugs or clothing.

#### **ENVIRONMENTAL HAZARDS**

- Toxic to small mammals, birds, aquatic organisms and non-target terrestrial plants.
- Observe buffer zones specified under DIRECTIONS FOR USE.
- This product will harm other broadleaved plants in the vicinity of the treatment area. If applying this product using a handheld sprayer, do not directly spray or allow the spray to drift onto ornamentals or gardens.
- Do not spray exposed roots of trees and ornamentals.

#### **LEACHING**

The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g. sandy soil) and/or the depth to the water table is shallow.

#### **RUNOFF**

- To reduce runoff from treated areas into aquatic habitats avoid application to areas with moderate to steep slope, compacted soil, or clay.
- Avoid application of this product when heavy rain is forecast.
- Contamination of aquatic areas as a result of runoff may be reduced by including a strip of untreated vegetation between the treated area and the edge of the water body.
- To prevent runoff from domestic uses avoid spraying on driveways, sidewalks or other hard surfaces. Do not irrigate within 24 hours after application.

## **DISPOSAL**

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

## **DIRECTIONS FOR USE**

### **USE PRECAUTIONS**

#### **WARNING**

Do not apply this product directly to or otherwise permit it to come into contact with vegetables, flowers, grapes, fruit trees, ornamentals or other desirable plants which are sensitive to 2,4-D or MCPA and do not permit spray mist containing it to drift onto them since even minute quantities of the spray may cause severe injury during both growing and dormant periods (coarse sprays are less likely to drift). Accordingly, application by airplane, ground rigs and hand dispensers should be carried out only when there is no hazard from drift. At higher temperatures, vaporization may cause injury to susceptible plants growing nearby.

Do not use on lawns or pastures until grass has become well established. Most legumes are usually damaged or killed. Do not contaminate irrigation ditches or water used for domestic purposes. Do not store near fertilizers, seeds, insecticides or fungicides.

To avoid injury to desirable plants, do not store, handle or apply other agricultural chemicals with the same containers used for this product. To avoid injury to desirable plants, containers used for this product should not be reused to store other materials. Rinse equipment and container thoroughly with water. Spray equipment used to apply 2,4-D or MCPA should not be used to apply other pesticides to susceptible crops. If it is necessary to do so, contact Corteva Agriscience Canada Company for detailed equipment clean-up procedures.

Local conditions may affect the use of herbicides. Provincial agricultural authorities issue recommendations to fit local conditions.

Be sure that use of this product conforms to all applicable regulations.

- DO NOT apply this product directly to freshwater habitats such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands, estuaries or marine habitats.
- DO NOT contaminate irrigation/drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

#### **Field sprayer application**

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) coarse classification. Boom height must be 60 cm or less above the crop or ground.



### **Aerial application**

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply when wind speed is greater than 16 km/h at flying height at the site of application. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) coarse classification. To reduce drift caused by turbulent wingtip vortices, the nozzle distribution along the spray boom length MUST NOT exceed 65% of the wing- or rotorspan.

### **Restricted Entry Interval (REI)**

All crops—12-hour REI

- Corn (field)—3-day REI
- Established grass pastures, rangeland - 3-day REI
- Non-cropland (annual and perennial weed control) – 9 day REI for scouting by foot.

### **General INSTRUCTIONS - CROP USE**

#### **Ground Application**

Apply 50 to 200 L of spray solution per hectare depending on the type of application equipment used. Use sufficient water for even distribution. Spray at low pressures (200 to 275 kPa) when the weeds are actively growing.

#### **Aerial Application (including fixed and rotary wing aircraft)**

Apply a minimum of 30 L of spray solution per hectare depending on the type of application system being used. Use boom pressures of 235 kPa or less. Avoid placing nozzles where spray will enter wing tip vortices.

### **MIXING**

#### **Transfer System**

For containers larger than 20 L, use a transfer system that avoids open pouring when transferring the liquid concentrate from such containers into the spray tank.

#### **To Prepare A Spray**

Add 1/2 the required amount of water or oil to the spray tank, then add with agitation the required amount of product as shown for the different uses and finally the balance of water or oil with continued agitation.

**WARNING:** If preparing straight oil mixtures, do not let water get into the product or the finished mixture.

**NOTE:** This product in water forms an emulsion, not a solution, which tends to separate on standing. Agitate to prevent such separation and ensure uniformity of spray mixtures.

#### **Selective Weed Control In Small Grains**

Weeds differ in their susceptibility to 2,4-D and not all types can be controlled satisfactorily in crops. The amount of product to use will depend upon the susceptibility and whether the crops will tolerate this amount. See Table 1.

#### **Wheat, Barley and Rye**

Spray from the 3 leaf expanded stage (15 cm) to just before the flag-leaf (shot-blade) stage. To avoid crop injury, do not treat during boot and flowering stages. Winter wheat and fall rye should be treated in early spring as soon as weeds appear and when the crop is in the full tillering stage to shot-blade stage. Maximum 1 application per year.

**Table 1: Suggested Amount of 2,4-D BEE-4 Herbicide For Weed Control in Crops (Not Underseeded)**

Mixed Weedy Growth	Stage of Weed Kind of Weather	L/ha	g Acid/ha	ha Treated/20 L Pail
Susceptible weeds such as: bluebur, burdock, cocklebur, field horsetail*, field pennycress, flixweed, goat's-beard, hoary cress*, kochia, lamb's-quarters, mustard (except dog and green tansy), plantain, prickly lettuce, ragweed, Russian knapweed, Russian thistle, shepherd's-purse, annual sunflower, sweet clover, vetch and wild radish	Seedling (2-4 leaves) Growing rapidly	700 mL	350	28
	Weeds in bud Dry or cool weather Heavy infestation	1.1	550	18
Hard-to-kill weeds such as: biennial wormwood, docks, dog mustard, field peppergrass, gumweed, hairy galinsoga, hedge bindweed, lady's-thumb, oak leaf goosefoot, redroot pigweed, smartweed, annual sow-thistle, tansy, tartary buckwheat, dandelion, tumbleweed, wild buckwheat*, blue lettuce*, Canada thistle*, field bindweed*, leafy spurge* and perennial sow-thistle*	Seedling (2-4 leaves) Growing rapidly	1.25†	625†	16
	Weeds in bud Dry, cool weather Heavy infestation	1.75†	875†	11

♦ Top growth control only.

∨ Better control will result from 2 treatments applied 1 week apart at 350 g/ha 2,4-D BEE-4 Herbicide.

†The higher rates may cause deformities and delayed maturity which will be offset by a higher relative yield from a less weedy crop.

**Pre-Harvest Interval**

- Do not permit lactating dairy animals to graze fields within 7 days after application.
- Do not harvest forage or cut hay within 30 days after application.
- Withdraw meat animals from treated fields at least 3 days before slaughter.

**Corn**

Postemergence: Treat with 700 mL/ha when corn plants are 15 to 20 cm tall and most weeds have germinated but are still in the seedling stage. Use drop nozzles to keep spray off corn foliage. Do not cultivate until at least 2 weeks after treatment. Maximum 1 application per year.

### **Pre-Harvest Interval**

- Do not permit lactating dairy animals to graze fields within 7 days after application.
- Do not harvest forage or cut hay within 30 days after application.
- Withdraw meat animals from treated fields at least 3 days before slaughter.

### **Rangeland, Grass Pasture**

To control susceptible weeds, use 2.0 to 3.5 L/ha in enough water for coverage. Maximum 2 applications per year.

### **Pre-Harvest Interval**

- Do not permit lactating dairy animals to graze fields within 7 days after application.
- Do not harvest forage or cut hay within 30 days after application.
- Withdraw meat animals from treated fields at least 3 days before slaughter.

### **Turf Weed Control**

To control susceptible weeds in turf use 2.0 to 3.1 L/ha in enough water for coverage.

- This product is only effective when applied to the leaves of actively growing weeds. This product will not prevent new weeds - apply only when weeds are present.
- DO NOT apply more than two broadcast applications per season. This does not include spot treatments.
- If weed populations do not warrant a broadcast application (e.g., entire lawn), consider spot treatments that target only weedy areas.
- DO NOT apply by air.
- DO NOT apply this product directly to freshwater habitats such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands, estuaries or marine habitats.
- DO NOT contaminate irrigation/drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.
- DO NOT apply to the exposed roots of trees and ornamentals.
- Field sprayer application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) coarse classification. Boom height must be 60 cm or less above the crop or ground.

### **Re-entry Interval for Turf**

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treated areas until spray has thoroughly dried.

### **Maximum Application Frequency for Turf**

For good turf/lawn management, normally two applications per year per treatment site are adequate. This does not include spot treatments.

**WARNING:** Do not use on lawns of creeping grasses such as bent, except for spot spraying or on freshly seeded turf until grass has become well established. Most legumes are usually damaged or killed.

### **Pre-Harvest Interval**

- Do not permit lactating dairy animals to graze fields within 7 days after application.
- Do not harvest forage or cut hay within 30 days after application.
- Withdraw meat animals from treated fields at least 3 days before slaughter.

## **GENERAL INSTRUCTIONS-NON-CROPLAND USE**

### **Woody Plants**

To control 2,4-D susceptible species, spray brush up to 2.0 m tall after foliage is well developed with 4.48 L in 1000 L of water. Using a high pressure spray (1000 to 1400 kPa) wet thoroughly all parts of plants including foliage, shoots, stems and bark. Maximum 2 applications per year.

Maximum Amount of Product Applied/day/person with Handheld Equipment: 5.4 L.

With good growing conditions and high soil moisture, sprays may be used up to 2 or 3 weeks before normal frost time but spraying soon after leaves are fully open in the spring usually gives best results. Control may be less effective during hot, dry weather when deep soil moisture is deficient. For 2,4-D resistant species, use Tordon 101 mixture herbicide.

#### **Roadside and General Weed Control**

For taller weed growth such as in vacant lots, on roadsides, rights-of-way, railway, hydro, roadways, highways, pipeline, industrial storage yards and along fencerows, sufficient spray should be used to wet all foliage. Use at the rate of 3.75 L in 1000 L of water. Deep-rooted perennial weeds such as bindweed and Canada thistle may require repeated applications as new growth appears. Maximum 2 applications per year.

Maximum Amount of Product Applied/day/person with Handheld Equipment: 5.4 L.

#### **BROADCAST AERIAL APPLICATION**

##### **Directions for Use**

Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. **Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.**

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices

##### **Use Precautions**

Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the National Aerial Pesticide Application Manual, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

Do not apply to any body of water. Avoid drifting of spray onto any body of water or other non-target areas. Specified buffer zones should be observed.

Coarse sprays are less likely to drift, therefore, avoid combinations of pressure and nozzle type that will result in fine particles (mist). Do not apply during periods of dead calm or when wind velocity and direction pose a risk of spray drift. Do not spray when the wind is blowing towards a nearby sensitive crop, garden, terrestrial habitat (such as shelter-belt) or aquatic habitat.

##### **Operator Precautions**

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

The field crew and the mixer/loaders must wear chemical resistant gloves, coveralls and goggles or face shield during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

**Product Specific Precautions for aerial applications to listed cereal crops, corn, pasture, rangeland and non-cropland**

Read and understand the entire label before opening this product. If you have questions, call the manufacturer at 1-800-667-3852 or obtain technical advice from the distributor or your provincial agricultural representative. Application of this specific product must meet and/or conform to the following:

- DO NOT apply when wind speed is greater than 16 kmph at flying height at the site of application. Do not spray in dead calm near sensitive plants. The “cloud” of suspended droplets may drift onto sensitive plants when the wind comes up. Spray only when wind is blowing away from a sensitive crop, shelterbelt or garden.
- Avoid direct applications to any body of water. Do not contaminate water through spray drift or by cleaning of equipment or disposal of wastes.
- Fixed and rotary-winged aerial applications to field crops should be made in a minimum of 30 L/ha total spray mixture. Use boom pressure of 235 kPa or less. Avoid placing nozzles where spray will enter wing tip vortices.
- Do not apply this product directly to, or otherwise permit it to come into direct contact with desirable crops, other desirable broadleaved plants or non-target species and do not permit spray mists to drift onto them.
- Coarse sprays are less likely to drift, therefore, avoid combinations of pressure and nozzle type that will result in fine particles (mist). Do not apply during periods of dead calm or when wind velocity and direction pose a risk of spray drift. Do not spray when the wind is blowing towards a nearby sensitive crop, garden, terrestrial habitat (such as shelter-belt) or aquatic habitat.
- Do not use human flaggers.
- Avoid Spray Drift: Apply only when there is little or no hazard from spray drift. Small quantities of the spray, which may not be visible, may seriously injure susceptible crops and damage sensitive non-target habitat. A method must be used to detect air movement, lapse conditions or temperature inversions (stable air) such as the use of balloons or a continuous smoke column at or near the spray site or a smoke generator on the spray equipment. If the smoke develops into layers or indicates a potential for hazardous spray drift, do not spray.
- Buffer Zones: Appropriate buffer zones should be established between treatment areas and aquatic systems and treatment areas and significant wildlife habitat.

**BUFFER ZONES TO PROTECT SENSITIVE HABITATS**

Buffer zones are not required for applications with handheld equipment.

Seasonal water bodies require buffer zones if there is water in them during application. Water bodies which do not fill on an annual basis need not be buffered.

**Buffer Zones - Sensitive Habitats**

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater

habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands), and estuarine/marine habitats.

**Buffer Zone Table**

Method of Application	Crop		Buffer Zones (metres) Required for the Protection of:					
			Freshwater Habitat of Depths:		Estuarine/Marine Habitats of Depths:		Terrestrial Habitat	
			Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m		
Field sprayer*	Turf		1	0	1	0	1	
	Field crops (wheat, barley and rye (cereal crops), corn		1	1	1	1	1	
	Pastures, rangeland, non-cropland (including rights-of-way** and brush control)		1	1	1	1	2	
Aerial	Field crops (wheat, barley and rye (cereal crops), corn		Fixed wing	10	0	1	0	45
			Rotary wing	10	0	1	0	40
	Pastures, rangelands		Fixed wing	15	0	1	0	60
			Rotary wing	15	0	1	0	50
	Non-cropland (including rights-of-way** and brush control)	4.5 L/ha	Fixed wing	70	0	1	0	150**
			Rotary wing	35	0	1	0	80**
		9.0 L/ha	Fixed wing	125	1	15	1	225**
			Rotary wing	50	1	5	1	100**

\* For field sprayer application, buffer zones can be reduced with the use of drift reducing spray shields. When using a spray boom fitted with a full shield (shroud, curtain) that extends to the crop canopy, the labelled buffer zone can be reduced by 70%. When using a spray boom where individual nozzles are fitted with cone-shaped shields that are no more than 30 cm above the crop canopy, the labelled buffer zone can be reduced by 30%.

\*\* For application to rights-of-way, buffer zones for protection of sensitive terrestrial habitats are not required.

When a tank mixture is used, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture.

**RESISTANCE MANAGEMENT RECOMMENDATIONS**

For resistance management, 2,4-D BEE-4 Herbicide is a Group 4 herbicide. Any weed population may contain or develop plants naturally resistant to 2,4-D BEE-4 Herbicide and other Group 4 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of 2,4-D BEE-4 Herbicide or other Group 4 herbicides with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted.
- Herbicide use should be based on an IPM program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical), cultural, biological and other chemical control practices.
- Monitor treated weed populations for resistance development.
- Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment and planting clean seed.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Corteva Agriscience Canada Company at 1-800-667-3852 or at [www.corteva.ca](http://www.corteva.ca).

**NOTICE TO USER:** This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

™ Trademark of Corteva Agriscience and its affiliated companies.