

2021-11-04
2021-1185

This label has been updated according to the re-evaluation decision of pyriproxyfen RVD2021-03. While users are encouraged to follow this updated label immediately, the previously approved label is valid until 26/02/2023 in accordance with the phase out period set out in RVD2021-03. This previously approved label will be provided upon request by emailing hc.pma.info-arla.sc@canada.ca. In your email please include the product name and Registration number of the label you are requesting.

NYGUARD IGR Concentrate

A Growth Regulator for Fleas

(Emulsifiable Concentrate)

ACTIVE INGREDIENT:

Pyriproxyfen..... 10.00%
MGK® - Registered trademark of McLaughlin Gormley King Company

COMMERCIAL

REGISTRATION NO. 25490

PEST CONTROL PRODUCTS ACT

KEEP OUT OF REACH OF CHILDREN
READ THE LABEL BEFORE USING

CAUTION EYE AND SKIN IRRITANT

Valent Canada Inc.,
201-230 Hanlon Creek Blvd.,
Guelph, Ontario
N1C 0A1
(519) 767-9262

Net Contents _____

Manufactured by:



7325 Aspen Lane North
Minneapolis, MN 55428
U.S.A.
(763) 544-0341

EPA Est. No. 1021-MN-2

Can prevent adult flea emergence indoors for 5 months (150 days).

NOTE: NYGUARD IGR Concentrate is an insect growth regulator that destroys flea populations without the direct poisoning action of conventional insecticides. NYGUARD IGR Concentrate controls flea populations by mimicking growth hormones found naturally in insects and affects the immature individuals in the population, thus preventing reproduction. In the case of fleas, it prevents the larvae from maturing into the adult biting stage. If large flea populations are present a conventional insecticide that is effective in killing these insects should be used in conjunction with NYGUARD IGR Concentrate to kill most of the adult population. NYGUARD IGR Concentrate will not affect the reproduction of fleas that are adults prior to spraying.

DIRECTIONS FOR USE

INDOOR FLEA CONTROL:

For Use in Homes, Apartment Buildings, Office Buildings, Automobiles, Buses, Boats, Ship Cabins and Holds, Trucks, Boxcars and Non-Food Areas of Hotels and Motels.

NYGUARD IGR Concentrate acts on the immature life stage of the flea preventing the adult flea from developing. Larvae crawling onto treated areas will not develop into adult fleas. Existing adult fleas and flea pupae are not affected.

If a high population of adult fleas (the biting stage) is present, the use of an adulticide may be necessary for immediate relief. For best results, use NYGUARD IGR Concentrate at the start of the flea season. When applied to surfaces where fleas are found, such as carpets, upholstered furniture and pet sleeping areas, NYGUARD IGR Concentrate can prevent adult flea emergence for 5 months (150 days). The degree of control may be reduced by heavy use, vacuuming or shampooing of treated surfaces.

Flea control on pets is important in prevention of introduction of adult fleas indoors. **DO NOT TREAT PETS WITH THIS PRODUCT.** For effective flea control, pets should be treated with a registered flea control product designed for use on pets, and treatment of the pet should be combined with the sanitation of any area used by the pet. Vacuum floors, carpets, and furniture (discard vacuum bag after use) and wash the pet's bedding, living quarters, and surrounding areas prior to treatment with this product.

Spray Preparation: Prepare a diluted spray solution by adding 5mL of NYGUARD IGR Concentrate for every 3.8 L of water. Partially fill mixing container with water, add NYGUARD IGR Concentrate, shake mixture and fill to final volume. Use spray solution within 48 hours of mixing. Shake well before each use.

Apply at the rate of 4L of diluted solution per 100m² of surface area with any low-pressure sprayer typically used for indoor applications. Treat all areas which may harbor fleas, including carpets, furniture, pet sleeping areas and throw rugs. Treat under cushions of upholstered furniture. **DO NOT** allow people or pets to enter treated areas until sprays have dried.

DO NOT apply to mattresses, linens, toys or clothing.

DO NOT apply to overhead areas or in confined spaces without appropriate respiratory and eye protection.

DO NOT allow spray to drip or allow drift onto non-target surfaces; any spray residue on non-target areas must be removed by the applicator.

DO NOT apply as a space spray.

Use a coarse droplet size and low pressure spray not exceeding 345 kPa (50 psi) to avoid splashing onto non-target surfaces.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed. Avoid contact with skin or eyes or clothing. Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. Wash with soap and warm water after use. Obtain medical attention if irritation persists. Do not use in edible product areas of food processing plants, restaurants, or other areas where food is commercially prepared or processed. **DO NOT** apply to surfaces that may come into contact with food/feed. Cover or remove all food/feed. Cover all food/feed processing surfaces, equipment and utensils or thoroughly wash following treatment. Ventilate treated areas after application either by opening windows and doors or through use of air exchange/ventilation systems confirmed to be operational. Use fans where required to aid in the circulation of air. **DO NOT** apply when people or pets are present. **BROADCAST APPLICATION:** **DO NOT** allow people or pets to enter treated areas for 2 hours or until sprays have dried. **SPOT AND CRACK AND CREVICE APPLICATION:** **DO NOT** allow people or pets to enter treated areas until sprays have dried.

FIRST AID: If in eyes: hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. **If on skin or clothing,** take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice. **If swallowed,** call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person. **If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

Take container, label, or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Treat symptomatically. Contains petroleum distillates – vomiting may cause aspiration pneumonia.

STORAGE: Store in a cool, dry place. Keep container closed.

DISPOSAL: Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank. Follow provincial instructions for any required additional cleaning of the container prior to its disposal. Make the container unsuitable for further use. Dispose of the container in accordance with provincial requirements. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

EMERGENCY TELEPHONE NUMBER:

In case of Medical emergencies or health and safety inquiries, or in case of fire, leaking or damaged containers, information may be obtained by calling **1-888-740-8712**.