

Code : YYY
NET CONTENTS: XXX

DI-THOATE 4E

EMULSIFIABLE CONCENTRATE INSECTICIDE

Uses
Farm buildings

AGRICULTURAL
GUARANTEE: Dimethoate 480 g/L

READ THE LABEL BEFORE USING

REGISTRATION NO. 14478 PEST CONTROL PRODUCTS ACT

DANGER POISON

VÉTOQUINOL N.-A. INC.
2000 ch. Georges, Lavaltrie (Québec) Canada J5T 3S5
1-800-363-1700

XXX 1 L, 4L
YYY : 02302008 (1L)
 02302014 (4L)

DIRECTIONS FOR USE:

DI-THOATE 4E is generally effective in controlling aphids, grasshoppers, house flies, leafhoppers, leafminers, mites, scales, tarnished plant bugs and thrips.

This product is not to be used around homes or other residential areas such as parks, school grounds, playing fields. It is not for use by homeowners or other uncertified users.

HOUSE FLY CONTROL: DI-THOATE 4E controls house flies for up to 6 weeks. Use a spray solution containing 250 mL of Di-Thoate 4E per 10 L of water. Repeated applications should be made when necessary. Good sanitation is an essential part of any effective fly control program.

In dairy barns, hog pens, calf barns, poultry houses and other farm buildings: Apply a residual spray to the ceiling, walls, stanchions, etc. Thoroughly wet all fly resting areas to the point of runoff. 50 L of water containing 1.0 to 1.2 L of Di-Thoate 4E will cover 50 to 100 m². For localized fly control, apply spot sprays with a knapsack or similar type sprayer to doorways, around windows, counters and other areas frequented by flies.

GENERAL OUTDOOR USE:

To control house flies around garbage cans, animal quarters, food processing plants, warehouses, loading docks and refuse areas. Spray exposed surface such as walls, fences, refuse containers and garbage. Exposure of sprayed areas to weather, particularly sunshine, reduces the duration of residual effects. Repeat applications as necessary.

PRECAUTIONS: KEEP OUT OF THE REACH OF CHILDREN. May be harmful or fatal if swallowed, inhaled or absorbed through the skin. Avoid breathing vapor or spray mist. Use only with adequate ventilation. Avoid contact with skin and eyes. Wash thoroughly after handling and before eating and smoking. Keep away from heat, sparks and open flame. Do not expose to direct sunlight. Remove animals from treatment area during application. Remove livestock and poultry from buildings when spraying. Do not use in milk processing or milk storage rooms. Do not contaminate feed, foodstuffs, feed troughs or drinking fountains. Do not mix with whitewash or apply within 2 weeks of whitewashing. Do not use in homes.

DISPOSAL: 1. Triple – or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank. 2. Follow provincial instruction for any required additional cleaning of the container prior to its disposal. 3. Make the empty container unsuitable for further use. 4. Dispose of the container in accordance with provincial requirements. 5. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

FIRST AID: If on skin or clothing, take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice. If in eyes, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. If swallowed, call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Symptoms of poisoning include nausea, vomiting, pinpoint pupils, excessive salivation, muscle twitching, convulsions and coma. This product, upon repeated or prolonged use, may cause cholinesterase inhibition.

ATROPINE IS ANTIDOTAL: Give 2 to 4 mg of atropine sulphate intramuscularly or intravenously immediately and every hour as required until pupils dilate. Pralidoxime chloride (2-Pam:PROTOPAM chloride) may be effective as an adjunct to atropine. Use according to label directions.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offense under the *Pest Control Product Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

STORAGE: Keep at room temperature (between 15 ° and 25 °C). Avoid freezing.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.