

2013-06-24
2012-4592

VET-KEM®

**BREAKAWAY®
FLEA & TICK COLLAR
FOR CATS**

4 Months Protection (symbol)

**DOMESTIC
GUARANTEE**
Propoxur...9.4%

REG. NO. 18506 P.C.P. ACT

**KEEP OUT OF REACH OF CHILDREN
READ LABEL BEFORE USING
FOR USE ONLY ON CATS**

Net Weight 10.8g

Wellmark International
100 Stone Road West, Suite 111
Guelph, ON N1G 5L3
1(800) 688-7378

GENERAL INFORMATION: For use in the control of cat and dog fleas and American dog and brown dog ticks on cats. The collar starts killing fleas and ticks when it is placed around the cat's neck. The fleas on the cat will be killed and new ones which may temporarily appear on the cat, will also be killed during the 4 months. Non-engorged ticks are tough occasional pests of cats and will be killed within a few days. Collar will kill ticks encountered in the first 2 months of wear.

DIRECTIONS: Do NOT remove the collar from its package until ready to use. Thread flat end of the collar through buckle hole and tighten as necessary, so that at least two ridges are covered by the buckle. Collars must be worn loosely, but securely enough to prevent easy removal. Leaving ample length for collar expansion as the cat grows, cut off any excessive length and discard it in household garbage. The expandable section molded into the collar will stretch under the weight of an average size cat and allow the cat to free itself if it becomes hooked or entangled by the collar. Since both the expandable section and plastic buckle are molded into the collar, the possibility of neck irritation and buckle failure is reduced. To release the collar, squeeze buckle between thumb and finger and pull sharply.

Do NOT use on sick or nursing animals, or on cats under 12 weeks of age or animals receiving drug or other pesticide treatment. Consult a veterinarian before using on aged or pregnant animals. Replace this collar no more than once every 4 months. Replace collar every 4 months for continual pest control in situations where there is an ongoing risk of pest infestation. **RETAIN PACKAGING INFORMATION FOR FUTURE REFERENCE.** Treatment must only be repeated if pest problem reoccurs.

For effective flea and tick control, treatment of the pet should be combined with sanitation of any area used by the pet. Vacuum floors, carpets and furniture (discard vacuum bag after use) and wash the pet's bedding, living quarters and surrounding areas. If pest problems persist, an insecticidal treatment of the premises may be required (Vet-Kem Siphotrol® products). A veterinarian should be consulted when pest infestation continues to be a problem on the pet.

PRECAUTIONS:

HUMANS: DO NOT ALLOW CHILDREN TO HANDLE THIS COLLAR. Dust will form on this collar during storage. Dust may irritate eyes, nose and throat. Harmful if dust is inhaled or absorbed through skin. Avoid contact with mouth, eyes, skin and clothing. After handling or applying, wash hands (or any other skin that came into contact with the product) with soap and water. Avoid product contact with food, food utensils or places where food is prepared or stored.

PETS: Harmful if collar is chewed or swallowed. Do NOT use this product at the same time or within 30 days before or after treatment with, or exposure to, other cholinesterase-inhibiting drugs, pesticides or chemicals. Dust may irritate eyes, nose and throat. Avoid letting pet inhale dust from collar. Avoid contact with cat's mouth, face, eyes and genitals. Some animals may be sensitive to this collar. Any collar, when fastened too tightly, may cause skin irritation. If irritation persists after proper adjustment, remove the collar and consult a veterinarian.

FIRST AID:

HUMANS: IF SWALLOWED, SUCKED OR CHEWED, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person. IF CONTACT WITH SKIN OR CLOTHING, take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice. IF IN EYES, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. IF INHALED, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

PETS:

Sensitivities may occur after using ANY pesticide for pets. If signs of sensitivity occur, remove the collar and bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately. If swallowed or chewed, consult a veterinarian.

Take container, label or product name and PCP Registration Number with you when seeking medical or veterinary attention.

TOXICOLOGICAL INFORMATION:

This product contains a pesticide that is a cholinesterase inhibitor (an anti-cholinesterase compound). Atropine and homatropine are antidotal. Dust in eyes will cause temporary pupillary constriction.

STORAGE: Store in original, unopened container at room temperature.

DISPOSAL: Dispose of any excess collar, packaging and unwanted (used) collars in household garbage.

100% satisfaction guaranteed.

HOT LINE 1-800-688-7378

® Registered Trademarks of Wellmark International

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.