

(Container Label)



GROUP	5	HERBICIDE
-------	---	-----------

LEXONE™ DF Toss-N-Go™ HERBICIDE

DISPERSIBLE GRANULE

FOR SALE FOR USE ON SOYBEANS * POTATOES * BARLEY * WHEAT * TOMATOES * FABABEANS
PEAS (WESTERN CANADA) * ASPARAGUS * FRUIT TREES *AND LENTILS (WESTERN CANADA)
ONLY

AGRICULTURAL

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

GUARANTEE: Metribuzin 75%

REGISTRATION NO. 21077 PEST CONTROL PRODUCTS ACT

DO NOT SELL WATER SOLUBLE BAGS SEPARATELY

NET CONTENTS: 2.5 KG
(5 X 500 g water-soluble pouches)

Corteva Agriscience Canada Company

2450, 215 – 2nd Street S.W.

Calgary, Alberta

T2P 1M4

1-800-667-3852

™Trademark of Corteva Agriscience and its affiliated companies.

PRECAUTIONS:

- **KEEP OUT OF REACH OF CHILDREN.**
- May be harmful if inhaled or swallowed.
- Avoid breathing dust or spray mist.
- Wear goggles when handling.
- Avoid contact with skin, eyes, and clothing.
- May irritate skin and eyes.
- Wash thoroughly with soap and warm water after handling.
- Wash contaminated clothing with soap and hot water before re-use.
- Do not sell soluble bags separately.
- Mixers, loaders, applicators, spray tank cleaners and other handlers must wear chemical resistant gloves, long-sleeved shirt, long pants, socks and shoes.
- Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove personal protective equipment immediately after handling this product. Wash the outside of the gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.
- Wear long pants, a long-sleeved shirt, socks and shoes and chemical-resistant gloves during mixing/loading, clean-up and repair activities

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at www.croplife.ca.

FIRST AID:

INHALATION:

Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

INGESTION:

Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

SKIN OR CLOTHING CONTACT:

Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

EYE CONTACT:

Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Treat Symptomatically.

STORAGE:

Store product in original container in a secure, dry, area away from other pesticides, fertilizers, food or feed. Not for use or storage in or around the home. Keep container tightly closed. Keep water-soluble bags in original outer bag away from moisture.

DISPOSAL:

1. Make empty container unsuitable for use.
2. Dispose of the container in accordance with provincial requirements
3. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the Provincial Regulatory Agency or the Manufacturer.

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the Pest Control Products Act to use this product in a way that is inconsistent with the directions on the label.

The seller warrants that the purchase by the buyer and the use of this product, as such, will not infringe and Canadian patent.

(Booklet)



GROUP	5	HERBICIDE
-------	---	-----------

LEXONE™ DF Toss-N-Go™ HERBICIDE

DISPERSIBLE GRANULE

FOR SALE FOR USE ON SOYBEANS * POTATOES * BARLEY * WHEAT * TOMATOES * FABABEANS
PEAS (WESTERN CANADA) * ASPARAGUS * FRUIT TREES *AND LENTILS (WESTERN CANADA)
ONLY

AGRICULTURAL

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

GUARANTEE: Metribuzin 75%

REGISTRATION NO. 21077 PEST CONTROL PRODUCTS ACT

DO NOT SELL WATER SOLUBLE BAGS SEPARATELY

NET CONTENTS: 2.5 KG
(5 X 500 g water-soluble pouches)

Corteva Agriscience Canada Company
2450, 215 – 2nd Street S.W.
Calgary, Alberta
T2P 1M4
1-800-667-3852

™Trademark of Corteva Agriscience and its affiliated companies.

PRECAUTIONS:

- **KEEP OUT OF REACH OF CHILDREN.**
- May be harmful if inhaled or swallowed
- Avoid breathing dust or spray mist.
- Wear goggles when handling.
- Avoid contact with skin, eyes, and clothing.
- May irritate skin and eyes.
- Wash thoroughly with soap and warm water after handling.
- Wash contaminated clothing with soap and hot water before re-use.
- Do not sell soluble bags separately.
- Mixers, loaders, applicators, spray tank cleaners and other handlers must wear chemical resistant gloves, long-sleeved shirt, long pants, socks and shoes.
- Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove personal protective equipment immediately after handling this product. Wash the outside of the gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.
- Wear long pants, a long-sleeved shirt, socks and shoes and chemical-resistant gloves during mixing/loading, clean-up and repair activities

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., contact 1-866-375-4648 or www.cropro.org/.

FIRST AID:

INHALATION:

Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

INGESTION:

Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

SKIN OR CLOTHING CONTACT:

Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

EYE CONTACT:

Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Treat Symptomatically.

IMPORTANT

Injury to or loss of desirable trees or other plants may result from failure to observe the following:

1. Do not apply (except as recommended for crop use), or drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not use on lawns, walks, driveways, tennis courts, or similar areas. Prevent drift of dry powder or spray to desirable plants. Do not contaminate any body of water.
2. Thoroughly clean all traces of LEXONE DF TOSS-N-GO Herbicide from application equipment immediately after use. Flush tank, pump, hoses, and boom with several changes of water after removing nozzle tips and screens (clean these parts separately). Wash away any spray mixture from the outside of spray tank, nozzle or spray rig.

NOTE: Do not use on muck soil or subsequent crops may be injured.

3. Rotation crops such as onions, celery, peppers, cole crops, lettuce, spinach, sugar beets, table beets, turnips, pumpkin, squash, cucumbers, melons, tobacco, and canola are sensitive to LEXONE™ DF Toss-N-Go™ Herbicide and may be injured if planted in LEXONE DF Toss-N-Go Herbicide treated soil both during the year of application and the following crop year.
4. Applications of LEXONE DF Toss-N-Go Herbicide made at rates above those recommended may result in burning, yellowing or stunting of the crop. If heavy rains occur soon after application, plant injury may result, especially in poorly drained areas where water may stand for several days.
5. Uneven application such as swath overlapping, variable tractor speed, spraying on turns, etc., may result in crop injury and increase chance of injury to rotation crops.

GENERAL INFORMATION

LEXONE DF Toss-N-Go Herbicide should be used only in accordance with recommendations on this label, or in separate published Corteva Agriscience Canada Company recommendations available through local dealers.

LEXONE DF Toss-N-Go Herbicide are a dry flowable granule contained within a water-soluble film. The water soluble film dissolves readily in water. Use gloves and goggles when handling the water-soluble bags. Reseal unused Toss-N-Go® bags into foil bag when not in use. Avoid exposure of the Toss-N-Go® bags to excessive moisture or physical damage during storage and prior to mixing in the spray tank. Do not reuse foil bags.

LEXONE DF Toss-N-Go Herbicide is intended for use as a spray for selective control of certain grasses and broadleaf weeds in soybeans, potatoes, tomatoes, fababeans, peas (Western Canada), asparagus, fruit trees and certain weeds in spring wheat (Western Canada only), spring barley (Western Canada only) and lentils (Western Canada only). It is non-corrosive to equipment, non-flammable and non-volatile.

WEEDS CONTROLLED:

When used as directed in Eastern Canada and B.C., LEXONE DF Toss-N-Go Herbicide controls the following grasses and broadleaf weeds:

GRASS WEEDS			BROADLEAF WEEDS	
Barnyard Grass (partial control)			Common Chickweed	Redroot Pigweed
Cheat Grass			Common Ragweed	Russian Thistle

Crab Grass	Dandelion (seedling)	Shepherd's Purse
Fall Panicum	Green Smartweed	Velvetleaf
Giant Foxtail	Jimsonweed	Wild Mustard
Green Foxtail	Lady's thumb	Yellow wood-sorrel
Johnson Grass (seedling)	Lamb's Quarters	
Witch Grass	Prickly mallow	
Yellow foxtail	Prostrate Pigweed	

For weeds controlled in Manitoba, Saskatchewan and Alberta, see crop use instructions. Additional weeds may be controlled when LEXONE DF Toss-N-Go Herbicide is used with other herbicides recommended on this label. Read product labels for further information.

DIRECTIONS FOR USE:
PREEMERGENCE USE:

Soil should be well-prepared and as free as possible from trash and clods; plant seed to specific depth on flat or raised seedbeds or injury may result. Moisture is necessary to activate this herbicide; best results are obtained if treatment is made to moist soil and moisture is supplied by rainfall or irrigation (0.5-1.0 cm) within two weeks after application. Do not apply a heavy irrigation immediately after application. If moisture is insufficient to activate the chemical or if the soil becomes crusted before crop emerges (7 to 10 days), a shallow rotary hoeing should be made before weeds become established; otherwise, avoid cultivating or disturbing the sprayed area. Do not cover treated bank with soil while cultivating between rows. The degree of control and duration of effect will vary with the amount applied, soil texture, rainfall and other conditions.

NOTE: If unusually heavy rains follow application, severe crop injury may result.

PREPLANT INCORPORATED (PPI) HERBICIDES WITH LEXONE DF Toss-N-Go Herbicide:

Herbicides recommended on this label for PPI application may not be used on peat, muck or other organic soils or on wet soils (mineral or organic) or on soils in poor working condition or on soils subject to prolonged periods of flooding. Apply with a tractor-mounted, fixed-boom sprayer properly calibrated to a constant speed and rate of delivery. Do not over-apply; misapplication or improper soil incorporation can result in erratic weed control or crop injury. Seedling disease, cold weather, deep planting (more than 5 cm), excessive moisture, high soil pH (above 7.5), high salt concentration, or drought may weaken seedlings and increase possibility of crop injury. Under these conditions delayed crop development or reduced yields may occur. Read and carefully follow incorporation instructions on all recommended herbicide labels.

USE RATES AND SOIL LIMITATIONS:

All dosages are expressed as broadcast rates. For band treatments, use proportionally less; for example for 30 cm band on 90 cm row, use 1/3 of the broadcast rate. Where a range of dosage rates is given, use the lower rate on lighter soils (low in clay or organic matter) and the higher rates on heavier soils. Do not use on sand as crop injury may result. Do not use on sandy or coarse textured soils with less than 2% organic matter.

EQUIPMENT:

Use a fixed-boom power spray properly calibrated to ensure a constant rate of application. Opening in screens should be equal to or larger than 50 mesh. Agitate by mechanical or hydraulic means in the spray tank. If by-pass or return line is used, it should terminate at bottom of tank to minimize foaming.

MIXING INSTRUCTIONS:

1. Fill tank about 1/4 full with fresh water.
2. Turn on full agitation.

3. Add the required number of LEXONE DF Toss-N-Go Herbicide. Maintain full agitation until bags are completely dissolved and product is fully dispersed.
4. Continue filling tank with water and add tank mix partners (if applicable) in the following sequence:
 - Dry formulations
 - Liquid formulations
 - Emulsifiable concentrates and surfactants
5. Continue with full agitation. On repeat tank loads, ensure that the amount of spray solution left in the tank from the previous load is less than 10% of the volume about to be mixed.

Do not mix, load or apply within 15 metres of all wells. Make scheduled checks of spray equipment. Mix only enough product for the job at hand. Avoid overfilling of spray tank.

NOTE:

Fall seeded or cover crops such as oats, and rye may be injured when seeded within the same season as the application of LEXONE DF Toss-N-Go Herbicide. Post-harvest tillage practices that provide thorough mixing of the soil may be helpful in reducing the chance of injury to succeeding crops.

USE PRECAUTIONS:

Do not mix, load, or apply within 15 meters of all wells.

Extreme care should be taken to avoid damage by drift onto plants outside the target area or onto ponds, waterways or ditches or land intended for cropping. When spraying near adjacent sensitive crops or plants, do everything possible to reduce spray drift.

Do not apply more than once per season on the same area and do not exceed the maximum use rate. Do not re-enter or allow re-entry into treated areas until 12 hours after application. Exception: If the product is soil-injected or soil-incorporated, workers may enter the treated area if there will be no contact with anything that has been treated.

It is recommended that this product not be applied in a way that will contact workers or other persons, either directly or through drift. Only handlers wearing personal protective equipment may be in the area during application.

Field sprayer application: **DO NOT** apply during periods of dead calm or when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) medium classification.

DO NOT apply by air.

Buffer zones:

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, pastures, rangelands, and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs, and wetlands) and estuarine/marine habitats.

Crop	Method of Application	Buffer Zone (metres) Required for Protection of:		
		Freshwater Habitats of Depths	Estuarine/Marine Habitats of Depths	Terrestrial Habitat

		Less than 1 m	1-3 m	Greater than 3 m	Less than 1 m	1-3 m	Greater than 3 m	
All crops except tomatoes	Field Sprayer	5	2	1	5	2	1	10
	Field Sprayer with Shrouds	2	1	0	2	1	0	3
	Field Sprayer with Cones	4	1	1	4	1	1	5
Tomatoes	Field Sprayer	10	3	1	5	2	1	15
	Field Sprayer with Shrouds	3	1	0	2	1	0	5
	Field Sprayer with Cones	5	2	1	4	1	1	10

When a tank mixture is used, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture.

ENVIRONMENTAL HAZARD

The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g., sandy soil) and/or the depth to the water table is shallow.

To reduce runoff from treated areas into aquatic habitats, consider the characteristics and conditions of the site before treatment. Site characteristics and conditions that may lead to runoff include, but are not limited to: heavy rainfall, moderate to steep slope, bare soil, poorly draining soil (e.g., soils that are compacted, fine textured, or low in organic matter such as clay).

Avoid application of this product when heavy rain is forecast.

1. SOYBEANS (Eastern Canada)

LEXONE DF Toss-N-Go Herbicide is recommended for use in soybeans:

- Preemergent to the crop.
- Preemergent to the crop tank mixed with Dual* or Pursuit*
- Preplant incorporated with Dual or Treflan* followed by LEXONE DF Toss-N-Go Herbicide preemergent.
- Preplant incorporated tank mixed with Treflan, Edge* DC, Dual or Merit* EC.

Treatments provide partial control of giant ragweed and cocklebur. American and Eastern Black nightshade is controlled by Dual®. Yellow nutsedge is controlled by Dual preplant incorporated at the high rate.

REPLANTING:

If initial seeding of soybeans fails to produce a stand, the same crop may be replanted; do not rework soil. Do not retreat field with a second application as injury to the crop may result. Do not replant treated areas to any other crop within 4 months as injury to subsequent crops may result.

NOTE:

Do not use on Altona, Vansoy, Maple Amber, Ace or Nattawa soybeans as these varieties are sensitive to LEXONE DF Toss-N-Go Herbicide and injury to the crop may result. Do not use treated vines for feed or forage. Injury to soybeans may occur if any Atrazine was applied on the soil the year before use of LEXONE DF Toss-N-Go Herbicide.

PREEMERGENCE APPLICATION:

Make a single application of LEXONE DF Toss-N-Go Herbicide as a band or broadcast spray after planting (at least 4.5 cm deep) but before soybeans emerge. Do not spray over top of emerged soybeans. In areas of heavy weed pressure, use the higher LEXONE DF Toss-N-Go Herbicide rate in the recommended range. In fields with variable soils, care should be taken to reduce treatment rate on sandy areas.

LEXONE DF Toss-N-Go Herbicide Alone - Broadcast Treatment

Apply in 150-300 L water per ha

SOIL TEXTURE¹	2 - 4% Organic Matter	Over 4% Organic Matter
Medium Soils (Loam, silt loam, silt, sandy clay, sandy clay loam)	640 - 960 g	960 g - 1.1 kg
Heavy Soils (Silty clay, silty clay loam clay, clay loam)	1.1 - 1.25 kg	1.25 - 1.5 kg

¹Do not use on sandy or coarse textured soils with less than 2% organic matter as crop injury may result.

SOYBEAN TANK MIXES

LEXONE DF Toss-N-Go Herbicide and TREFLAN

PREPLANT INCORPORATED:

Before planting, mix LEXONE DF Toss-N-Go Herbicide with water and add Treflan as directed under "Mixing Instructions" above. Immediately incorporate into soil prepared according to the Treflan label.

Do not plant seed more than 5 cm deep.

TREFLAN (PPI) and LEXONE DF Toss-N-Go Herbicide (PRE):

Apply Treflan preplant incorporated followed by LEXONE DF Toss-N-Go Herbicide preemergence. Follow directions on the Treflan label.

RATES PER HECTARE:

Apply in 150-300 L water per ha

SOIL TEXTURE¹	LEXONE DF Toss-N-Go Herbicide ²		TREFLAN 545 EC
	PPI	PRE	
Medium Soils (Loam, silt loam, sandy clay sandy clay loam)	540 g	540-640 g	1.5 L
Heavy Soils (Silty clay, silty clay loam clay, clay loam)	640 g	640-960 g	2.1 L

¹ Do not use on sandy or coarse textured soils with less than 2% organic matter as crop injury may result.

² Use the higher LEXONE DF Toss-N-Go Herbicide rate on soils with more than 4% organic matter.

LEXONE DF Toss-N-Go Herbicide AND DUAL

PREPLANT INCORPORATED:

Mix LEXONE DF Toss-N-Go Herbicide with water and add Dual as directed under "Mixing Instructions". Incorporate to a depth of 5 cm within 14 days of planting. Preplant incorporated treatment will control dominant infestations of nutsedge and/or annual grasses if higher rates of application are used.

PREEMERGENCE:

Apply a tank mixture, prepared as above, as a band or broadcast spray during or after planting, before the weeds or crop have emerged.

SPLIT APPLICATION:

Following preplant incorporation of Dual, apply LEXONE DF Toss-N-Go Herbicide as a preemergence application.

RATES PER HECTARE

Apply in 150-300 L water per ha

SOIL TEXTURE ¹	LEXONE DF Toss-N-Go Herbicide		DUAL
	PPI	PRE ²	
Medium Soils (Loam, silt loam, sandy clay, sandy clay loam)	540 g	540-640 g	1.75-2.75 L
Heavy Soils (Silty clay, silty clay, loam clay, clay loam)	640 g	640-960 g	1.75-2.75 L

¹ Do not use on sandy or coarse textured soils with less than 2% organic matter as crop injury may result.

² Use higher rate on soils with more than 4% organic matter.

LEXONE DF Toss-N-Go Herbicide plus EDGE (Eastern Canada):

PREPLANT INCORPORATED:

Mix LEXONE DF Toss-N-Go Herbicide with water and add Edge DC or Edge DF as directed in the "Mixing Instructions" sections. Refer to the Edge® labels for recommendations, directions for use, restrictions and any additional weeds not specified on this label. Tank mixes of LEXONE DF Toss-N-Go Herbicide and Edge DC or Edge DF may be applied up to 10 days before planting.

NOTE:

Read and carefully observe information on the Edge DC Herbicide or Edge DF Herbicide label.

Do not use the foliage from soybeans treated with LEXONE DF Toss-N-Go Herbicide and Edge for feed or forage.

DO NOT APPLY BY AIR.

RATES PER HECTARE:

Apply in 150-300 L water per ha

SOIL TEXTURE ¹	LEXONE DF Toss-N-Go Herbicide	EDGE DC
Light (sandy loam, loamy sand)	470 g	1.1 kg
Medium (loam, silt loam, silt sandy clay, sandy clay loam)	550 g	1.4 kg
Heavy (silty clay, silty clay loam, clay, clay loam)	700 g	1.9 kg

¹ Do not use on sandy or coarse textured soils with less than 2% organic matter.

LEXONE DF Toss-N-Go Herbicide plus LOROX® DF/LOROX® L Herbicide plus DUAL Herbicide

PREEMERGENCE APPLICATION:

Mix LEXONE DF Toss-N-Go Herbicide followed by LOROX® Herbicide and then Dual Herbicide in the spray tank with water and apply as a band or broadcast spray after planting but before soybeans emerge. Read and carefully observe information on LOROX® Herbicide and Dual Herbicide labels.

RATES PER HECTARE:

Apply in 200-300 L water per ha

SOIL TEXTURE ¹	LEXONE DF Toss-N-Go Herbicide	LOROX® DF Herbicide or LOROX® L Herbicide	DUAL Herbicide
Light (sandy loam, loamy sand)	210-400 g	1.5-1.75 kg or 1.6-1.8 L	1.75-2.1 L
Medium (loam, silt loam, silt sandy clay, sandy clay loam)	385-465 g	1.7-1.9 kg or 1.8-2.0 L	2.0-2.1 L
Heavy (silty clay, silty clay loam, clay, clay loam)	465-550 g	1.75-2.0 kg or 1.8-2.1 L	2.1-2.75 L

¹Do not use on light textured soils with less than 2% organic matter.

Use lower rates for soil texture when organic matter is 2 - 4%; higher rates when organic matter is more than 4%.

LEXONE DF Toss-N-Go Herbicide plus PURSUIT Herbicide (Eastern Ontario only):

LEXONE DF Toss-N-Go Herbicide may be tank mixed with Pursuit for a pre-emergent application to provide a broader spectrum of weed control. Add Pursuit as directed in the "Mixing Instructions" sections. Follow rates, recommendations, precautions and restrictions on the Pursuit Herbicide label.

DO NOT APPLY BY AIR.

LEXONE DF Toss-N-Go Herbicide plus MERIT EC Herbicide (Eastern Canada only):

PREPLANT INCORPORATED:

LEXONE DF Toss-N-Go Herbicide may be tank mixed with Merit EC to improve the control of cocklebur, pigweed and wild mustard, and to give more consistent control of common ragweed and yellow foxtail than Merit EC Herbicide alone. Read and carefully observe the information on the Merit EC Herbicide label.

RATES PER HECTARE

Apply in 150 - 300 L water per ha

SOIL TEXTURE ¹	LEXONE DF Toss-N-Go Herbicide	MERIT EC Herbicide
Light (sandy loam, loamy sand)	270 g	1.4 L
Medium (loam, silt loam, silt sandy clay, sandy clay loam)	330 g	1.6 L
Heavy (silty clay, silty clay loam, clay, clay loam)	400 g	1.8 L

¹Do not use on sandy or coarse textured soils with less than 2% organic matter as crop injury may result.

DO NOT APPLY BY AIR.

Do not allow livestock to graze on treated soybean vines or feed livestock treated vines or vine trash.

2. POTATOES

GENERAL INFORMATION

Do not use on sandy or coarse textured soils containing less than 1% organic matter as crop injury may result. Potato varieties vary in their resistance to LEXONE DF Toss-N-Go Herbicide. Use on limited area to determine safety before larger scale sprays are adopted. Do not use on varieties Tobique or Belleisle. The variety Shepody has shown sensitivity to LEXONE DF Toss-N-Go Herbicide applied at ground crack. An earlier preemergent application has shown greater crop safety. At the high rate, LEXONE DF Toss-N-Go Herbicide will provide partial control of quackgrass, yellow nutsedge and will improve grass control when used in combination with a recommended grass herbicide. Supplemental row cultivation helps weed control.

POTATOES IN EASTERN CANADA, B.C., AND PRAIRIE PROVINCES

LEXONE DF Toss-N-Go Herbicide Plus GRAMOXONE TANK-MIX PREEMERGENCE:

LEXONE DF Toss-N-Go Herbicide may be applied at rates of 830 g to 1.2 kg/ha in a tank mix combination with Gramoxone (200 g/L) at 2.75 L/ha as a preemergence broadcast application to potatoes after weeds have emerged for control of emerged broadleaf weeds and grasses, including quackgrass. Refer to the Gramoxone Label for recommendations, directions for use, restrictions and precautions and any additional weeds not specified on this label.

POTATOES IN EASTERN CANADA, COASTAL B.C. AND IRRIGATED CROPS IN ALBERTA AND MANITOBA

LEXONE DF TOSS-N-GO ALONE PREEMERGENCE:

Make a single application of LEXONE DF Toss-N-Go Herbicide at 640 g-1.4 kg/ha after planting (at least 5 cm deep) or hilling but before crop emerges. See paragraph on Use Rates and Soil Limitations. Use 150-300 L water per ha. Apply before emerged weeds are 3 cm high.

EARLY POSTEMERGENCE:

If weather conditions prevent a preemergent application, 700 - 960 g/ha may be applied following 3 or more successive days of sunny weather. Treat before weeds are 3 cm tall and when potatoes are less than 10 cm in height. Treatment may cause some leaf burn. Do not use on red-skinned, Shepody, or early potatoes. Do not apply within 60 days of harvest.

PREEMERGENCE plus POSTEMERGENCE:

Follow directions as indicated but do not apply more than 1.4 kg/ha per season. Do not apply within 60 days of harvest.

TANK MIXES

LEXONE DF Toss-N-Go Herbicide and DUAL Herbicide (Eastern Canada Only)

Mix LEXONE DF Toss-N-Go Herbicide with water. Add Dual as directed in the "Mixing Instructions" sections of this label.

Do not use Dual Herbicide on muck, peat muck or high organic matter soils.

Do not use Dual Herbicide on the potato variety Superior.

Do not use Dual Herbicide at ground crack or if potatoes have emerged.

Preplant incorporated:

Apply the LEXONE DF Toss-N-Go Herbicide-Dual Herbicide tank mixture to the soil and incorporate to a depth of 5 cm within 14 days of planting. Preplant incorporated treatments will control dominant infestations of nutsedge and/or annual grasses when higher rates are used.

Preemergence:

Apply the LEXONE DF Toss-N-Go Herbicide-Dual Herbicide tank mixture as a band or broadcast spray during or after planting before the weeds or crop have emerged.

Split Application:

Following preplant incorporation of Dual Herbicide, apply LEXONE DF Toss-N-Go® Herbicide as a preemergence application.

Split Application:

Apply Dual Herbicide as a preplant incorporated treatment or a preemergent treatment followed by LEXONE DF Toss-N-Go Herbicide as an early postemergent treatment. Do not use LEXONE DF Toss-N-Go Herbicide on red-skinned, Shepody, or early potatoes. Do not apply within 60 days of harvest.

RATES PER HECTARE:

Apply in 150-300 L water per ha

SOIL TEXTURE ¹	LEXONE DF Toss-N-Go Herbicide		DUAL Herbicide
	PPI or PRE	POST	
Medium to Heavy Soils	830 g - 1.2 kg	830 - 960 g	2.0-2.75 L

¹Do not use on sandy or coarse textured soils with less than 1% organic matter as crop injury may result.

POTATOES IN PRAIRIE PROVINCES AND INTERIOR B.C.

See comments under POTATOES - GENERAL INFORMATION.

EARLY POSTEMERGENCE:

Apply LEXONE DF Toss-N-Go Herbicide over the top of potato plants soon after emergence and before weeds are 4 cm high. Do not use on red-skinned, Shepody, or any early maturing varieties. Do not apply within 60 days of harvest.

Use 350 g/ha of LEXONE DF Toss-N-Go Herbicide as a broadcast spray. Weeds controlled are:

wild Mustard	Hemp-nettle	Tartary Buckwheat
Chickweed	Ball Mustard	Corn Spurry
Green Smartweed	Lamb's-quarters	Lady's-thumb
Stinkweed	Volunteer Rape	Shepherd's-purse
Redroot Pigweed		

PREPLANT INCORPORATED – LEXONE DF Toss-N-Go Herbicide and Eptam 8-E:

Apply 350-550 g/ha of LEXONE DF Toss-N-Go Herbicide with 4.25-5.5 L/ha Eptam 8-E as a pre-plant incorporated treatment in 200-300 litres of water per hectare. Mix LEXONE DF Toss-N-Go Herbicide in water and add Eptam 8-E as directed under "Mixing Instructions". For proper soil preparation, application and incorporation in the soil see the appropriate section of the Eptam 8-E label. Do not use on soil with greater than 7% organic matter or on sandy or coarse textured soils with less than 2% organic matter as

crop injury may result. Do not use on red-skinned or early potatoes or on the varieties Belleisle or Tobique.
Weeds controlled are:

Chickweed	Redroot Pigweed
Green Smartweed	Shepherd's-purse
Hemp-nettle	Stinkweed
Lady's-thumb	Wild Mustard
Lamb's-quarters	Volunteer Rape

3. FABABEANS

PREPLANT INCORPORATION:

SPRING APPLICATION:

Apply 350-550 g/ha of LEXONE DF Toss-N-Go Herbicide tank mixed* with 1.5-2 L/ha Treflan 545 in 80-100 litres of water.

FALL APPLICATION:

Apply 400-550 g/ha of LEXONE DF Toss-N-Go Herbicide TOSS-N-GO BAGS tank mixed* with 2.26 L/ha Treflan 545 in 80-100 litres of water.

*Mix LEXONE DF Toss-N-Go Herbicide in water. Add Treflan as directed under Mixing Instructions and immediately incorporate into soil prepared according to the Treflan label instructions.

This treatment may control the following weeds:

Wild Mustard	Volunteer Rape	Lamb's-quarters
Hemp-nettle	Stinkweed	Shepherd's-purse
Green Smartweed	Redroot Pigweed	Lady's-thumb
Chickweed		

Follow instructions on the Treflan label to avoid erratic weed control and/or crop injury from uneven application or improper incorporation in the soil.

Do not use LEXONE DF Toss-N-Go Herbicide on muck soils. Rotation crops such as: vegetables, canola, or tobacco may be injured if planted on treated soil within 2 years after application. Injury to succeeding crops may be reduced by a post harvest tillage of the soil.

CAUTION:

Do not graze the treated crops or cut for hay; sufficient data are not available to support such use.

4. PEAS, FIELD AND PROCESSING (Western Canada)

Apply LEXONE DF Toss-N-Go Herbicide at 280-375 g/ha as a postemergent broadcast spray in a minimum of 100 L of water/ha. Spray when weeds have emerged and are less than 5 cm in height and before pea vines are 15 cm long. Crop must be planted at least 5 cm below soil surface and not underseeded with forage. Do not use on field peas grown under irrigation.

The following weeds may be controlled by this treatment:

Ball Mustard	Lamb's-quarters
Chickweed	Stinkweed
Corn Spurry	Tartary Buckwheat

Green Smartweed
Hemp-nettle

Volunteer Rape
Wild Mustard

Use the higher rate when weed infestation is dense. Weed control may be reduced if rain falls within 6 hours after spray. Do not apply more than once per season. Do not tank mix with other herbicides or adjuvants. Do not apply on light soils containing less than 3% organic matter. Do not apply within 4-5 days after frost or periods of cool, wet or cloudy weather. Allow a 4-5 day interval between an application of LEXONE DF Toss-N-Go Herbicide and other pesticides. Do not harvest processing peas within 40 days of application. Do not harvest or graze or feed treated crop until 70 days after application.

5. SPRING WHEAT & SPRING BARLEY (Western Canada only)

POSTEMERGENCE:

Use LEXONE DF Toss-N-Go Herbicide alone or combined with Banvel® Herbicide or MCPA Amine 500. When using a tank mix, mix LEXONE DF Toss-N-Go Herbicide with water then add Banvel or MCPA as directed under Mixing Instructions".

Refer to the Banvel Herbicide or MCPA Amine 500 labels for recommendations, directions for use, restrictions and additional weeds specified on the labels.

Crop must be planted 5 cm deep and not underseeded with forages. Apply only once per season. Weed control may not be noticeable for 7 to 10 days. Temporary (7-10 days) lightening in crop colour and occasionally a slight reduction in crop height may be observed, especially if frost or abnormally high temperatures occur within 1-2 days of application. If frost occurs allow 4-5 days for crop to recover before applying LEXONE DF Toss-N-Go Herbicide.

Allow a 4-5 day interval between application of LEXONE DF Toss-N-Go Herbicide and wild oat herbicides. Do not use on Klondike variety of barley.

Treatments will control:

Ball mustard
Stinkweed
Hemp-nettle
Wild mustard

Smartweed
Corn spurry
Volunteer rape
Lamb's-quarters

Chickweed
Tartary buckwheat
Lady's-thumb

Use the higher rate for the best residual control of chickweed or when weed infestation is dense. Rainfall immediately after an application may reduce weed control. Do not apply within 60 days of harvest. Do not graze or feed crop to livestock within 30 days of application.

LEXONE DF TOSS-N-GO HERBICIDE PLUS BANVEL OR MCPA AMINE 500 - RATES PER HECTARE (Apply in 80-100 L water per ha when weeds are small)			
CROP	STAGE OF CROP GROWTH		
	2-5 leaf stage (mid-tillering) LEXONE DF TOSS-N-GO HERBICIDE - (ALONE)	2-3 leaf stage ¹ (mid-tillering) LEXONE DF TOSS-N-GO HERBICIDE + BANVEL Herbicide	3-5 leaf stage (mid-tillering) LEXONE DF TOSS-N-GO HERBICIDE + MCPA AMINE 500
SPRING BARLEY	270-350 g	270g –LEXONE DF Herbicide + 275 ml BANVEL Herbicide	275 – 350 g LEXONE DF Herbicide + 850 ml– 1.1 L MCPA AMINE 500
SPRING WHEAT	270 g	270g –LEXONE DF Herbicide + 275 ml BANVEL Herbicide	275g LEXONE DF Herbicide + 850 ml– 1.1 L MCPA AMINE 500

¹ On wheat, application may be made at the 4 leaf stage but crop tolerance may be reduced.

6. TOMATOES

LEXONE DF Toss-N-Go Herbicide may be used on tomato transplants grown for processing only. Apply as a directed spray before weeds are 4 cm high. Avoid contacting at least the top 2/3 of the tomato foliage with spray. Best results are usually obtained when application is made to well established plants about 3 weeks after transplanting. Avoid applications when tomatoes are under stress from poor growing conditions such as cool, wet and cloudy weather. Do not apply to direct-seeded tomatoes.

Apply only once per season. Do not apply within 60 days of harvest.

LEXONE DF Toss-N-Go Herbicide - RATES PER HECTARE Apply in 200-300 L Water per ha

SOIL TEXTURE¹	LEXONE DF Toss-N-Go Herbicide
Light (sandy loam, loamy sand)	320 g ¹
Medium (loam, silt loam, silt sandy clay, sandy clay loam)	640 g
Heavy (silty clay, silty clay loam, clay, clay loam)	640 g - 1.1 kg ²

¹ May not control grasses

² High rate for very weedy fields only

7. ASPARAGUS

LEXONE DF Toss-N-Go Herbicide may be used on asparagus as either a preemergence application in established asparagus or as a delayed postemergence treatment for transplanted and established asparagus for the control of annual grass and broadleaf weeds (see section "Weeds Controlled" for specific weeds).

PREEMERGENCE:

Apply 1.5 kg/ha of LEXONE DF Toss-N-Go Herbicide in 100 to 300 L of water to established asparagus as a preemergence broadcast spray with low pressure (150-275 kPa) ground equipment. Do not apply within 14 days of harvest.

POSTEMERGENCE:

Make one application per season of LEXONE DF Toss-N-Go Herbicide as a delayed postemergence treatment for transplanted and established asparagus. Apply 1.5 kg/ha of LEXONE DF Toss-N-Go Herbicide in 100 to 300 L of water with ground equipment. Application of LEXONE DF Toss-N-Go Herbicide should be made after completion of harvest when spears have developed and when weeds are well established.

8. FRUIT TREES

LEXONE DF Toss-N-Go Herbicide may be used in apple, cherry, peach and apricot trees in the year of planting for the control of annual weeds (see section "Weeds Controlled" for specific weeds). Make a single application, pre-plant incorporated, over the entire area or within the planting row. Apply 0.5 to 1.0 kg/ha of LEXONE DF Toss-N-Go Herbicide.

NOTE TO BUYER/USER: READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR SPECIAL USE APPLICATIONS.

The Directions for Use for this product for the uses described below were developed by persons other than Corteva Agriscience Canada Company and accepted for registration by Health Canada under the User Requested Minor Use Label Expansion Program. Corteva Agriscience Canada Company itself makes no representation of warranty with respect to product performance (efficacy) and crop tolerance (phytotoxicity) of this product when used on the crops listed. Accordingly, the Buyer and User assume all liability arising, and agree to hold Corteva Agriscience Canada Company harmless from any claims based on efficacy or phytotoxicity in connection with these uses described below.

ESTABLISHED FRUIT TREES (apple, peach, pear, sweet and sour cherry, plum and apricot):

Apply LEXONE DF alone at 1.0 kg/ha for broadleaf weed control in established fruit trees. Application, by ground equipment only, should be made in 150-300 L water per ha as a pre-emergent treatment. Apply as a band under the trees and make only one application per year. This will control:

BROADLEAF WEEDS:

Common chickweed	Lady's-thumb	Russian thistle
Common ragweed	Lamb's-quarters	Shepherd's-purse
Dandelion (seedling)	Prickly mallow	Velvetleaf
Green smartweed	Prostrate pigweed	Wild mustard
Jimsonweed	Redroot pigweed	Yellow wood-sorrel

ESTABLISHED FRUIT TREES (apple, peach, pear, sweet and sour cherry, plum and apricot):

LEXONE DF may be tank mixed with Dual 960E to provide broad spectrum weed control in established fruit trees. Apply LEXONE DF at 1.0 kg/ha and Dual 960E at 2.75 L/ha in 150 - 300 L water per ha as a pre-emergent treatment. Apply as a band under the trees and make only one application per year. This tank mix will control:

GRASS WEEDS:

Barnyard grass (partial control)	Giant foxtail	Yellow foxtail
Cheatgrass	Green foxtail	Yellow nutsedge
Crabgrass (smooth, hairy)	Johnson grass (seedling)	

Fall Panicum	Witchgrass	
BROADLEAF WEEDS:		
American nightshade	Jimsonweed	Russian thistle
Common chickweed	Lady's-thumb	Shepherd's-purse
Common ragweed	Lamb's-quarters	Velvetleaf
Dandelion (seedling)	Prickly mallow	Wild mustard
Eastern black nightshade	Prostrate pigweed	Yellow wood-sorrel
Green smartweed	Redroot pigweed	
ESTABLISHED FRUIT TREES (apple, peach, pear, sweet and sour cherry)		
<p>LEXONE DF may be tank mixed with SINBAR® to provide broad spectrum weed control in established fruit trees. Apply LEXONE DF at 1.0 kg/ha and SINBAR® at 0.63 kg/ha in 150-300 L of water per hectare as a pre-emergent treatment. Apply as a band under the trees and make only one application per year. This tank mix will control:</p>		
GRASS WEEDS:		
Barnyardgrass	Fall panicum	Perennial rye grass
Bluegrass	Giant foxtail	Wild barley
Cheatgrass (Downy brome)	Green foxtail	Witchgrass
Crabgrass	Johnson grass (seedling)	
BROADLEAF WEEDS:		
Annual sowthistle	Lady's-thumb	Russian thistle
Common chickweed	Lamb's-quarters	Shepherd's-purse
Common ragweed	Prickly lettuce	Stinkweed
Dandelion (seedling)	Prickly mallow	Velvetleaf
Green smartweed	Prostrate pigweed	Wild mustard
Henbit	Purslane	Yellow wood-sorrel
Jimsonweed	Redroot pigweed	
<p>Treatment usually provides partial control of quackgrass, horsenettle, and yellow nutsedge.</p> <p>DO NOT use on soil coarser than sandy loams with less than 3% organic matter.</p> <p>PROCESSING TOMATOES</p> <p>Apply the following tankmix as a post emergent broadcast application:</p>		
LEXONE DF Toss-N-Go Herbicide		200 g/ha
plus		
BRAVO® 500 Fungicide		2.4-4.0 L/ha
<p>Make only one application per growing season. Application should be made using ground equipment only.</p> <p>DO NOT APPLY THIS TANKMIX WITHIN 30 DAYS OF HARVEST.</p> <p>This tank mix will control:</p>		
DISEASES CONTROLLED:	WEEDS CONTROLLED:	
Early/Late Blight	BROADLEAF WEEDS:	

Septoria Leaf Spot	Common chickweed	Prostrate pigweed
Anthrachnose	Common ragweed	Redroot pigweed
Botrytis Grey Mold	Dandelion (seedling)	Russian thistle
	Green smartweed	Shepherd's purse
	Jimsonweed	Velvetleaf
	Lady's-thumb	Wild mustard
	Lamb's-quarters	Yellow wood-sorrel
	Prickly mallow	

NOTE: Grassy weeds listed on the LEXONE label may not be controlled at this rate.

9. LENTILS (WESTERN CANADA)

Apply 280 g/ha as a postemergent broadcast spray in a minimum of 100 L of water/ha on lentil vines less than 10 cm high and after weeds have emerged. Crop must be planted more than 5 cm below soil surface and not underseeded with forage.

Do not tank mix with other herbicides or adjuvants.

Do not apply on light soils containing less than 3% organic matter.

Do not apply more than once per season.

Do not harvest or graze or feed treated crop until 70 days after application.

The following weeds may be controlled by this treatment:

Ball Mustard	Lamb's-quarters
Chickweed	Stinkweed
Corn Spurry	Tartary Buckwheat
Green Smartweed	Volunteer Rape
Hemp-nettle	Wild Mustard

Weed control may not be noticeable for 7-10 days after application. Weed control may be reduced if rain falls within 6 hours after spray. Some reduction of colour and height may occur, especially if frost or abnormally high temperatures occur 1-2 days after application. Do not spray within 4-5 days after frost. Allow a 4-5 day interval between an application of LEXONE DF and other pesticides.

NOTE:

Application of more than the recommended rate, or after the crop stage specified above, may result in crop damage. To minimize risk, adhere very carefully to all label instructions.

SPRAY TANK CLEAN OUT:

To prevent spray tank residues from injuring sensitive crops in subsequent applications, use the following procedure to clean LEXONE DF from the spray equipment.

1. Drain tank. Thoroughly hose down the interior surfaces of the tank, then flush tank, boom and hoses with clean water for 5 minutes.
2. Remove screens, filters and nozzles and clean separately.

RESISTANCE-MANAGEMENT RECOMMENDATIONS:

For resistance management, LEXONE DF Toss-N-Go Herbicide is a Group 5 herbicide. Any weed population may contain or develop plants naturally resistant to LEXONE DF Toss-N-Go Herbicide and other Group 5 herbicides. The resistance biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but

specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- ◆ Where possible rotate the use of LEXONE DF Toss-N-Go Herbicide or other Group 5 herbicides with different herbicide groups that control the same weeds in a field.
- ◆ Use tank mixtures with herbicides from different group when such use is permitted
- ◆ Herbicide use should be based on an IPM program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical), cultural, biological and other chemical control practices.
- ◆ Monitor treated weed populations for resistance development.
- ◆ Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment and planting clean seed.
- ◆ Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- ◆ For further information or to report suspected resistance, contact your local Corteva Agriscience Canada Company representative or the Corteva Agriscience Canada Company hotline at 1-800-667-3852 for further information

STORAGE:

Store product in original container in a secure, dry, area away from other pesticides, fertilizers, food or feed. Not for use or storage in or around the home. Keep container tightly closed. Keep water-soluble bags in original outer bag away from moisture.

DISPOSAL:

1. Make empty container unsuitable for use.
2. Dispose of the container in accordance with provincial requirements
3. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the Provincial Regulatory Agency or the Manufacturer.

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the Pest Control Products Act to use this product in a way that is inconsistent with the directions on the label.

The seller warrants that the purchase by the buyer and the use of this product, as such, will not infringe and Canadian patent.

TMTrademark of Corteva Agriscience and its affiliated companies.

*All other products are trademarks of their respective companies

<<Water Soluble Bag>>

LEXONE™ DF TOSS-N-GO™ HERBICIDE
DISPERSIBLE GRANULES
KEEP OUT OF REACH OF CHILDREN

SEE DIRECTIONS FOR USE BOOKLET FOR COMPLETE PRECAUTIONARY STATEMENTS

GUARANTEE: Metribuzin 75%

DISPERSIBLE GRANULES

NET CONTENTS: 500 GRAMS

CAUTION