This label has been updated according to the re-evaluation decision of Pyrethrins RVD2023-06 and Piperonyl Butoxide RVD2023-07. While users are encouraged to follow this updated label immediately, the previously approved label is valid until "02/03/2025" in accordance with the phase out period set out in RVD2023-06 and RVD2023-07. This previously approved label will be provided upon request by emailing hc.pmra.info-arla.sc@canada.ca</u>. In your email please include the product name and Registration number of the label you are requesting.

VET-KEM[®]

OVITROL[™] MOUSSE

Kills dog fleas, cat fleas, brown dog ticks and American dog ticks.

FOR USE ONLY ON DOGS & CATS DO NOT USE ON PUPPIES OR KITTENS LESS THAN 8 WEEKS OLD DO NOT USE ON ANY OTHER ANIMAL

With Precor[®] Insect Growth Regulator (IGR) Kills adult fleas Kills flea eggs Prevents hatching of flea eggs laid on your pet for up to 45 days

DOMESTIC

ACTIVE INGREDIENTS (S) Methoprene 0.5% Pyrethrins 0.4% Piperonyl Butoxide 5.0%

REG.NO. 22678 P.C.P. ACT

Contains sodium nitrite at 0.10% as a preservative

READ THE LABEL BEFORE USING KEEP OUT OF REACH OF CHILDREN

CAUTION



EXPLOSIVE

Wellmark International 100 Stone Road West, Suite 111, Guelph, ON N1G 5L3 2023-1821 2024-02-01

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

FIRST AID:

HUMANS: If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice. If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to- mouth, if possible. Call a poison control centre or doctor. DO NOT induce vomiting unless told to do so by a poison control centre or doctor. DO NOT give any liquid to the person. DO NOT give anything by mouth to an unconscious person.

PETS: Sensitivities may occur after using ANY pesticide for pets. Discontinue use if dermal or nasal irritation develops. Although the Vet-Kem Ovitrol Mousse is specifically formulated for these animals, some particularly sensitive cats may exhibit temporary and non-harmful salivation when exposed to products containing pyrethrins. In cases of known sensitivity, the use of the lower treatment rates is recommended. If signs of sensitivity occur bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

Take container, label or product name and PCP Registration Number with you when seeking medical or veterinary attention.

TOXICOLOGICAL INFORMATION: This product contains petroleum distillates. Vomiting may cause aspiration pneumonia. Treat symptomatically.

PRECAUTIONS:

PHYSICAL AND CHEMICAL HAZARDS:

Caution Explosive: Contents under pressure. Keep away from heat or open flame. **DO NOT** puncture container. **DO NOT** expose to temperatures over 50°C.

HUMANS: KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed. Avoid contact with eyes, mouth, skin and clothing. Wear rubber gloves, during application of this product. After handling or applying, wash hands (or any other skin that came into contact with the product) with soap and water. Avoid contact with pet until dry. Avoid product contact with food, food utensils or places where food is prepared or stored. **PETS:** Harmful if swallowed. Avoid contact with pet's mouth, face, eyes and genitals.

Side Effects:

Monitor your dog after application. Side effects may include: skin irritation such as redness or scratching; changes in behaviour such as agitation or lethargy; or gastrointestinal effects such as vomiting or loss of appetite. If these or other side effects occur consult your veterinarian or registrant at 1-800-688-7378.

Monitor your cat after application. Side effects may include: skin irritation such as scratching or hair loss at the application site, or changes in behaviour such as agitation or lethargy. Gastrointestinal effects such as drooling, vomiting, or loss of appetite may also occur. If these or other side effects occur consult your veterinarian or registrant at 1-800-688-7378.

DIRECTIONS FOR USE:

Vet-Kem Ovitrol Mousse is a contact and residual insecticidal foam which kills adult cat fleas, dog fleas, American dog and brown dog ticks. Effectively breaks the flea life cycle and prevents the growth of pre-adult stages of the flea. Its unique combination of ingredients kills adult fleas for 7 to 10 days and keeps the flea eggs from hatching for up to 45 days.

FOR USE ONLY ON DOGS AND CATS

Use Limitations

DO NOT use on puppies or kittens less than 8 weeks old.

DO NOT USE ON ANY OTHER ANIMAL.

Consult a veterinarian before using on sick, aged, pregnant or nursing animals or animals receiving drug or other pesticide treatment.

This product contains Pyrethrins, Piperonyl Butoxide and (S) Methoprene. **DO NOT** apply another pest control product such as a shampoo, spot on, collar or powder which contains Pyrethrins, Piperonyl Butoxide and (S) Methoprene to the treated animal when using Vet-Kem Ovitrol Mousse.

DO NOT use this product more often than once every month. Treatment must only be repeated if pest problem reoccurs.

To kill dog fleas, cat fleas, brown dog ticks and American dog ticks:

DO NOT administer by mouth. Use ONLY in a well-ventilated area, preferably outdoors. Shake can well before using. Hold can about 2 cm above pet's back and spray the entire body of the pet, beginning at the head and working towards the tail. Ticks are difficult to control and must be sprayed directly. Pay special attention to neck, hindquarters, areas under legs, around the tail and between toes. **DO NOT** apply to pet's face, eyes, nose, mouth or genitals. Rub the foam thoroughly into the pet's coat using a comb, brush or fingertips in an amount which moistens, but does not soak the coat. Wipe off any excess product from the pet with a dry cloth and then comb or brush. Using this technique, the application of 2 to 4 grams of Vet-Kem Ovitrol Mousse per kilogram of body weight (i.e., 1 to 2 strips of foam down the pet's body length) will provide effective long-term flea control.

For effective flea and tick control, treatment of the pet should be combined with sanitation of any area used by the pet. Vacuum floors, carpets and furniture (discard

2023-1821 2024-02-01

vacuum bag after use) and wash the pet's bedding, living quarters and surrounding areas. If pest problems persist, a premise treatment with an insecticide labelled for that use may be required (eg. Vet-Kem Siphotrol[®] products). A veterinarian should be consulted when pest infestation continues to be a problem on the pet. The use of an effective product for controlling adult and larva flea populations in the pet's environment will reduce the possibility of re-infestation of the animal.

STORAGE: Store this product away from food or feed. Contents under pressure. **DO NOT** puncture container. **DO NOT** expose to temperature over 50°C. Keep away from heat or open flames.

DISPOSAL: DO NOT reuse the empty containers. Dispose in household garbage. Unused or partially used products should be disposed at provincially or municipally designated hazardous waste disposal sites.

100% satisfaction guaranteed. HOT LINE 1-800-688-7378

® Registered Trademarks of Wellmark International Ovitrol is a trademark of Wellmark International