

2020-08-25
Sub.No. 2020-3528

Group 11 Insecticide

FORAY 76B
Biological Insecticide Aqueous Suspension

For Forests and Woodlands

Restricted

GUARANTEE:

Bacillus thuringiensis subsp. *kurstaki* strain ABTS-351, slurry 73.88%

Potency: 16,700 Cabbage Looper Units (CLU)/mg of product (equivalent to 20 billion CLU/L.).

The percent active ingredient does not indicate product performance and potency measurements are not federally standardized.

Warning, contains the allergen sulfites.

REGISTRATION NO. 24976 PEST CONTROL PRODUCTS ACT

POTENTIAL SENSITIZER
CAUTION EYE IRRITANT
READ THE LABEL BEFORE USING

Lot. No.
Net Contents (Litres)

Date of Manufacture:
This product should be used within 6 months of the date of manufacture.

Registrant:
Valent BioSciences LLC
1910 Innovation Way, Suite 100
Libertyville, IL 60048
USA 1-800-323-9597

Canadian Agent:
Valent Canada, Inc.
201-230 Hanlon Creek Blvd
Guelph, Ontario N1C 0A1

PRECAUTIONS:

KEEP OUT OF REACH OF UNAUTHORIZED PERSONNEL.

MAY CAUSE SENSITIZATION

CAUTION EYE IRRITANT

Avoid contact with skin, eyes, and clothing. Avoid breathing dust/spray mist. Wear a long sleeved shirt, long pants, waterproof gloves, shoes and socks, eye goggles and NIOSH-approved respirator with any N-95, R-95, or P-95 filter for biological products when handling, mixing/loading or applying the product and during all clean-up/repair activities. When applicators use closed systems (e.g. enclosed cabs) the personal protective equipment requirements may be reduced or modified. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

FIRST AID	
If on skin or clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.
If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes.. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing the eye Call a poison control centre or doctor for treatment advice.
If inhaled	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.
If swallowed	Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. DO NOT give anything by mouth to an unconscious person.
General	Seek medical attention if irritation or signs of toxicity occur and persist or is severe. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

Treat symptomatically.

STORAGE

In order to ensure microbial purity and potency, Foray 76B should be stored in the original container at 0 - 25°C and used within 6 months of the date of manufacture. After extended storage, vigorously shake or stir contents to assure a uniform suspension.

DISPOSAL:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the

location of the nearest collection site.

Before taking the container to the collection site:

Triple- or pressure-rinse the empty container. Add the rinsing to the spray mixture in the tank. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

DIRECTION FOR USE:

DO NOT apply this product through any type of irrigation system.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes. As this pesticide is not registered for control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT allow spray mixture to stand in the tank for more than 12 hours. Recirculate mixture before loading and/or application.

Foray 76B contains the spores and insecticidal crystals of a naturally occurring strain of *Bacillus thuringiensis* subspecies *kurstaki*.

Foray 76B is highly specific to some lepidopteran larvae. It is a stomach poison only and therefore must be ingested by the larvae to be effective. After ingestion of a sufficient dose, feeding ceases and death will follow in 2-5 days.

In general, larvae should be treated when they are newly hatched. Young larvae (early instars) are most susceptible to the effects of *Bacillus thuringiensis* subspecies *kurstaki* and less damage will have been caused by larval feeding. Larvae must be feeding actively when treatments are made since ingestion is necessary for effective results. The direct target for application is the foliage, buds, and flowers (objects of larval feeding). Thorough spray coverage with well-distributed deposits of Foray 76B is necessary to assure that the feeding larvae ingest a toxic dose.

Foray 76B is a water based formulation which may be applied undiluted or diluted with clean water. Dilute with minimal quantities of water only when required to improve deposit. Do not use petroleum-based solvents. Foray 76B does not settle out rapidly, so continual agitation is not recommended and should be avoided. Use diluted mixture of Foray 76B within 12 hours of mixing and recirculate mixture before loading and/or application.

2020-08-25

Sub.No. 2020-3528

Even though Foray 76B deposits are relatively resistant to wash-off, avoid application when significant rainfall is imminent. For early morning applications, foliage should not be wet with dew to the point of runoff.

Consult with recognized forest pest control authority or Valent BioSciences LLC representative regarding appropriate timing and method of application. The timing and number of applications required for effective control will depend upon foliage development and larval activity. For effective control, applications should coincide with sufficient foliage development (shoot elongation or leaf expansion) to ensure maximum spray deposit. To the extent possible, apply treatments after egg hatch has been completed to assure that the maximum number of larvae is present during the treatment period. If Foray 76B is applied to later larval instars and/or extremely high larval populations, use the higher label rates and/or additional applications.

To control eastern spruce budworm, treat prior to 5th instar larval development and when bud flush/shoot development is sufficient to allow good deposit on emerging needles.

Timing of treatment of the two-year cycle spruce budworm depends on the year of its life cycle. For first year larvae treat early instars when bud flush/shoot development is sufficient to allow good deposit on emerging needles. For second year larvae treat larvae at peak 4th instar when bud flush/shoot development is sufficient to allow good deposit on emerging needles.

For consistently acceptable control of western spruce budworm, two-year cycle spruce budworm and western blackhead budworm, higher rates and volumes of spray may be required to achieve satisfactory deposit in mountainous terrain.

Treat against gypsy moth when larvae are in 2nd and 3rd instar and when leaf expansion is 40-50%. When egg hatch is extended, 2 or more applications, 7-10 days apart, may be required.

For best control of eastern hemlock looper, apply when larvae are young (early instars) before extensive damage occurs and/or when a majority of buds have flushed or shed bud caps. Repeat applications at intervals to maintain control, usually 3-14 days depending on larval growth and weather. If control is sought with a single spray, wait until egg hatch is essentially complete.

To control whitemarked tussock moth in softwood species, make the first application at late egg hatch and follow up with a second application 5-7 days later.

For control of forest tent caterpillar apply undiluted Foray 76B at a rate of 12-20 BCLU /ha (0.6 - 1.0L product/ha) when larvae are approximately 1.3 cm in length and when leaves of deciduous trees are approximately 50 % expanded; apply by air or ground; if necessary repeat application in 7 - 10 days.

For control of Jackpine budworms, apply undiluted at 20-30 BCLU /ha to early larval instars; double applications several days apart may be considered. For consistent control of very high larval populations, one single application at higher rates up to 60 BCLU /ha may be required.

For control of Eastern and Western Blackheaded Budworm apply two applications at 30 BCLU /ha or one application at 60 BCLU /ha. For two applications at 30 BCLU /ha make the first application at the late egg hatch and follow up with a second application 3 to 7 days later. For a single application at 60 BCLU /ha, apply after 100% egg hatch.

GROUND APPLICATION:

With ground equipment, provide thorough coverage while minimizing runoff. Dilute Foray 76B only to provide sufficient volume for thorough coverage.

For mist blower applications, add the recommended volume of Foray 76B to water, at a mix ratio ranging from 1:20 to 1:50. Do not exceed 150 litres/hectare total mixture.

For high volume hydraulic sprayers, add the recommended volume of Foray 76B to water at a mix ratio ranging from 1:50 to 1:500. Do not exceed 1500 litres/hectare total mixture.

AERIAL APPLICATION INSTRUCTIONS:

Apply undiluted only by fixed-wing or rotary aircraft equipment that has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application by using appropriate marking devices and/or electronic guidance equipment.

Use Precautions:

Apply only when meteorological conditions at the treatment site allow for complete and even coverage. Apply only when meteorological conditions are in compliance with local and /or provincial authorities.

Use Precautions:

DO NOT allow the pilot to mix product to be loaded onto the aircraft. However, loading of premixed product with a closed system is permitted. It is desirable that the pilot has communication capabilities at each treatment site at the time of application.

The field crew and the mixer/loaders must wear the personal protective equipment described in the PRECAUTIONS section of this label. When handlers/loaders use closed systems to load product onto the aircraft, the handler requirement for eye goggles and a NIOSH-approved

2020-08-25

Sub.No. 2020-3528

respirator/mask with any N-95, R-95, or P-95 filter for biological products may be waived. When reduced personal protective equipment is worn, the respirator/mask and eye goggles must be immediately available for use in an emergency such as a spill or equipment breakdown.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing must be washed before reuse. Decontaminate aircraft cockpit and vehicle cabs regularly

Product Precautions:

Read and understand the entire label before opening this product. If you have questions, call the manufacturer at 1-800-323-9597 or obtain technical advice from the distributor or from your provincial agricultural or forestry representative. Application of this specific product must meet and/or conform to the aerial uses and rates on this label.

RESTRICTED USES: FORESTS, WOODLANDS AND RESIDENTIAL AREAS

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the Pest Control Products Act to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

NATURE OF RESTRICTION: This product is to be used only in the manner authorized. Consult local pesticide regulatory authorities about use permits which may be required.

Forestry Use: Ground/Aerial Application for sites greater than 500 ha

Woodlands Use: Aerial Application for sites 500 ha or less

Residential Use: Aerial Application

Consult local Transport Canada office regarding air regulations for aerial applications for urban and/or forestry applications. Foray 76B may be used for aerial application in urban areas for treatment of residential areas and municipal recreational areas (including parks, parkland, vacant lots, shelterbelts, and rights of way under municipal jurisdiction).

Undiluted applications are recommended for Eastern and Western Spruce Budworm, Two-year Cycle Spruce Budworm, Gypsy Moth, Jackpine Budworm, Eastern Hemlock Looper, Whitemarked Tussock moth and Forest Tent Caterpillar, and Eastern and Western Blackheaded Budworm. However, Foray 76B may be diluted with water and applied at the rates indicated in the dose rates table. Total volume of spray material to be applied per hectare depends upon target pest, target foliage, weather, spray equipment and droplet size.

Recommended droplet size spectra are from 30-80 microns for coniferous foliage and 50-150 microns for broadleaf foliage.

DOSE RATES: AERIAL AND GROUND APPLICATION¹

Forests, Woodlands, Farm Woodlots and Residential Areas

Target Insect	Rate Litres/hectare	Dose BCLU /hectare
Eastern Spruce Budworm	0.75-1.5 ²	15-30
Western Spruce Budworm	1.5-2.0	30-40
Gypsy Moth	1.5-2.5	30-50
Jackpine Budworm	1.0-3.0 ³	20-60
Eastern Hemlock Looper	1.5-2.0	30-40
Whitemarked Tussock Moth	1.5-2.5	30-50
Forest Tent Caterpillar	0.6-1.0 ²	12-20
Eastern Blackheaded Budworm	1.5-3.0	30-60
Two-year-cycle budworm	1.5 – 2.0	30 - 40
Western Blackhead Budworm	1.5 – 3.0	30 - 60

¹ See the RESTRICTED USES section of this label for aerial application to forests, woodlands, and residential areas, and ground application to forests.

² For satisfactory control of Forest Tent Caterpillar or Eastern Spruce Budworm at lower rates and volumes (less than or equal to 1 litre/hectare; 20 BCLU /ha.), Foray 76B must be applied using enhanced atomization in order to ensure generation of sufficient spray droplets for impingement on conifer foliage (balsam fir, black spruce). This requires use of Micronaire AU4000 atomizers with appropriate flow rate (2 L/min/head or less) and blade angle settings to ensure a minimum rotation of 9000 rpm.

³ For control of Jackpine budworms, apply undiluted at 20-30 BCLU /ha to early larval instars; double applications several days apart may be considered. For consistent control of very high larval populations, one single application at higher rates up to 60 BCLU /ha may be required.

RESISTANCE MANAGEMENT:

For resistance management, please note that Foray 76B contains a Group 11 insecticide. Any insect population may contain individuals naturally resistant to Foray 76B and other Group 11 insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same site. Other resistance mechanisms that are not linked to site of action but are specific for individual chemicals, such as enhanced metabolism, may also exist. The following appropriate resistance management strategies should be followed to delay insecticide resistance:

- Where possible, rotate the use of Foray 76B or other Group 11 insecticides with different groups that control the same pests in a site.
- Insecticide use should be based on an Integrated Pest Management program that includes scouting, record keeping, and considers cultural, biological and other chemical control practices.
- Monitor treated pest populations for resistance development.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance management and/or integrated pest management recommendations for the specific site and pest problems in your area.

2020-08-25

Sub.No. 2020-3528

- For further information or to report suspected resistance, contact Valent BioSciences LLC at 1-800-323-9597

NOTICE TO USER:

This control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.