

This label has been updated according to the re-evaluation decision of Pyrethrins RVD2023-06 and Piperonyl Butoxide RVD2023-07. While users are encouraged to follow this updated label immediately, the previously approved label is valid until "02/03/2025" in accordance with the phase out period set out in RVD2023-06 and RVD2023-07. This previously approved label will be provided upon request by emailing hc.pmra.info-arla.sc@canada.ca. In your email please include the product name and Registration number of the label you are requesting.

PYROCID[®] Pressurized Multi-Purpose Spray 7416

[Pyrethrins]

COMMERCIAL

For indoor use at listed structures housing livestock or poultry, listed food/feed handling establishments, listed modes of transport and other listed structures to kill listed pests. **DO NOT** apply in residential areas

For outdoor use around listed structures to kill adult mosquitoes, biting midges (no-see-ums) and flies.

For use on beef and dairy cattle (lactating and nonlactating), horses, goats, sheep and hogs to kill listed pests.

For indoor use at poultry houses and for use on poultry to kill Poultry lice, Darkling beetles, Bed bugs and Mites

REGISTRATION NO. 24729

PEST CONTROL PRODUCTS ACT

READ THE LABEL BEFORE USING

ACTIVE INGREDIENT:

Pyrethrins.....	0.50%
Piperonyl butoxide	4.80%

PYROCID[®] - Registered trademark of McLaughlin Gormley King Co.

KEEP OUT OF REACH OF CHILDREN

CAUTION



POISON



EXPLOSIVE

Manufactured by:



7325 Aspen Lane North
Minneapolis, MN 55428
U.S.A.

(763) 544-0341

EPA Est. No. 1021-MN-2

NET CONTENTS: _____

VALENT CANADA INC., 201-230 HANLON CREEK BLVD., GUELPH, ONTARIO N1C 0A1, (519) 767-9262

Contains the botanical insecticide, Pyrethrum.

Kills Stable Flies, Horse Flies, Deer Flies, House Flies, Mosquitoes and Biting Midges (No-see-ums)

For use in Livestock Living Quarters, Dairies, Milk Rooms, Milking Parlors, Calving Barns, Dairy and Hog Barns

For use on Beef Cattle, Dairy Cattle, Horses, Hogs, Poultry

PYROCID[®] Pressurized Multi-Purpose Spray 7416 can be used in listed livestock living quarters. This product provides fast, effective kill of the listed pests. PYROCID[®] Pressurized Multi-Purpose Spray 7416 can also be used on listed livestock and horses for quick kill of pests on these animals.

For use in: Livestock Holding Areas, Livestock living Quarters, Attics, Barns, Beef Barns, Structures Housing Livestock or Poultry, Calving Barns, Canneries, Commercial Buildings, Dairy Barns, Dairies, Dried Fruit Storage Areas, Farm Structures, Food Warehouses, Goat Holding Areas, Grain Elevators, Grain Warehouses, Horse Stables, Hog Barns, Kennels, Livestock Holding Areas, Loafing Sheds, Milk Houses, Milk Rooms, Milk Sheds, Milking Rooms, Poultry Houses, Rice Mills, Seed Warehouses, Sheep Holding Areas, Stables, Storage Areas, Swine Houses, Tobacco Warehouses, Trailers, Trucks, Utility Rooms, Veterinaries, Warehouses, Washrooms, Wheat Mills.

DIRECTIONS FOR USE:

READ ALL DIRECTIONS COMPLETELY BEFORE USE. SHAKE BEFORE USE AND AT INTERVALS DURING PROLONGED USE. Hold container upright with nozzle away from face. Press valve down and spray as directed.

PYROCID[®] Pressurized Multi-Purpose Spray 7416 contains the quick killing botanical insecticide, Pyrethrum, which provides effective control of listed insects in listed locations of use and on listed livestock upon direct contact.

Locations of Use

PYROCID[®] Pressurized Multi-Purpose Spray 7416 can be used indoors and outdoors around the following structures*:

STRUCTURES HOUSING LIVESTOCK OR POULTRY: livestock living quarters, dairy barns, beef barns, calving barns, stables, horse stables, hog barns, swine houses, poultry houses, livestock holding areas, goat holding areas, sheep holding areas, loafing sheds;
FOOD/FEED HANDLING ESTABLISHMENTS: milking parlours, milk rooms, milk houses, dairies, canneries, food warehouses, dried fruit storage areas, grain warehouses, grain elevators, rice mills, wheat mills, seed warehouses, tobacco warehouses;
MODES OF TRANSPORT: cargo ships, trailers, trucks;
OTHER STRUCTURES: barns, commercial buildings, farm structures, kennels, storage areas, utility rooms, veterinaries, warehouses and washrooms.
*DO NOT apply in residential areas. Residential areas are defined as any use site where the general public, including children, could be exposed during or after application. For structural uses, in residential sites, this includes homes, schools, restaurants, public buildings, modes of transport in areas where passengers are present, or any other areas where the general public including children may potentially be exposed.
PYROCID® Pressurized Multi-Purpose Spray 7416 can also be used for quick kill of listed pests on beef cattle, dairy cattle (lactating and non-lactating), horses, hogs, sheep, goats and poultry.

Use Limitations for All Uses

- **DO NOT** apply when people or pets are present. Additionally for indoor space applications, **DO NOT** apply when livestock (with exception of poultry) are present.
 - **DO NOT** apply when a food/feed processing facility, milking parlour, milk house or milk room is in operation.
 - **DO NOT** apply to surfaces that may come into contact with food/feed. Cover or remove all food/feed (including bulk tanks that contain milk). Cover or remove all milking equipment and thoroughly wash before use. Cover all food/feed processing surfaces, equipment and utensils or thoroughly wash them following treatment. Any food/feed (including milk) accidentally contaminated with spray solution must be discarded.
 - When applied indoors: **DO NOT** apply to overhead areas or in confined spaces without appropriate respiratory and eye protection. **DO NOT** allow spray to drip or allow drift onto non-target surfaces.
- Ventilate treated areas after application either by opening windows and doors or using fans, where required, to aid in the circulation of air. Air exchange/ventilation systems confirmed to be operational may also be used.
- When applied indoors as a space spray: **DO NOT** remain in treated areas after an indoor space application; **DO NOT** allow people, pets or livestock (with exception of poultry) to enter treated areas until 2 hours after application. The commercial applicator is responsible for informing workers and others of this requirement.
 - When indoors as a surface spray: **DO NOT** allow people or pets to enter treated areas until sprays have dried.
 - When applied outdoors: **DO NOT** apply in areas where plants are grown for food or feed (e.g., vegetables, fruit, herbs). As this product is not registered for the control of pests in aquatic systems, **DO NOT** use to control aquatic pests. **DO NOT** contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

FOR INDOOR USE AT LISTED STRUCTURES

Within the specified use location, to enhance effectiveness of this product, apply directly to the listed pests where possible. For all labelled uses, treatment must only be repeated after {min re-application interval} days if pest problem persists or reoccurs. Good sanitary practices should be maintained to prevent re-infestation.

How to Apply

Apply as a space spray, crack and crevice, perimeter*, spot, broadcast* or furniture (listed storage furniture only) treatment. ***PERIMETER AND BROADCAST TREATMENTS ARE PERMITTED IN POULTRY HOUSES ONLY.** Space application is a suspension of fine droplets in the air within an indoor space. Crack and crevice is an application directly into narrow openings on the surface of the structure. It does not include the treatment of exposed surfaces. Narrow openings typically occur at expansion joints, utility entry points and along baseboards and mouldings. Perimeter application in poultry houses is less than 0.3 m wide along the edges of a room to baseboards, wall-floor and ceiling-wall joints, and around doorways or windows. Spot application is localized to a surface area not more than 0.2 m². Spots are not to be adjoining and the combined area of spots is not to exceed 10% of the total surface area of a room. Broadcast application in poultry houses is to broad expanses of indoor structural surfaces such as walls, floors, ceilings and indoor foundation walls/crawlspaces. Furniture treatment: *broadcast application* covers large areas of the entire surface of items; *crack and crevice treatments* are applications to junction points on items; and *spot application* is up to 10% of the surface of the item.

GENERAL DIRECTIONS PRIOR TO THE TREATMENT OF LISTED STORED PRODUCT PESTS: Good sanitation is necessary for effective treatment of listed stored product pests. Before treatment, remove and dispose of all residues (e.g., deposits and accumulations of food/feed products, waste grain, debris, dust, dirt) and thoroughly vacuum and clean indoor structural surfaces, including walls, floors, ceilings, rafters and door frames; interior and exterior surfaces of conveying, processing and handling equipment; pantries, cabinets, drawers and shelves; and around stored product containers, especially in cracks and crevices (discard vacuum bag after use).

SPACE APPLICATION:

To kill Flies, Mosquitoes, Biting midges (nosee-ums), House fly, Lesser house fly, Stable fly, Horse fly, Deer fly, Horn fly, Black fly, Indian meal moth, Angoumois grain moth, Tobacco moth, Clothes moth, Chocolate moth, Almond moth, Miller moth: Before using as a space spray in structures with livestock, remove all livestock, with exception of poultry, from area to be treated. Close all windows and doors and shut off all ventilating systems and sources of ignition before treatment. While holding can upright at arm's length, direct the spray in sweeping motion to all parts of the room, especially upward to ceiling, upper corners of the room, windows and other light sources, and around pallets, stacks of stored products and stored product containers, and not directly towards poultry. Begin application at far side of the room and work backwards toward exit. **DO NOT** exceed 5-10 seconds per 30 m³. Vacate the treated area and keep windows and doors closed for 15 minutes before ventilating. For ventilation and re-entry requirements, see "Use Limitations for All Uses" (above). After the re-entry interval has passed, clean up dead insects.

SURFACE APPLICATION:

GENERAL DIRECTIONS FOR SURFACE TREATMENTS: Apply spray at the rate of 30 seconds for each 28 square meters of surface area. Keep area closed for 45 minutes after treatment. Ventilate room 20 minutes before re-entry.

To kill cockroaches (brown banded cockroach, German cockroach, American cockroach, Oriental cockroach), crickets, firebrat, silverfish, booklice, spiders, centipedes, sowbugs, pillbugs, earwigs, clover mites, millipedes, ants, pharaoh ants and carpenter ants [Eastern (Western)]: Thoroughly treat indoor structural surfaces, including cracks and crevices in walls, floors and ceilings; along and behind baseboards; around water pipes and tanks; walls behind and floors beneath appliances (e.g., refrigerators and stoves) and furniture (e.g., cabinets, storage units, bookcases); around sinks, drains and pipes; localized areas around garbage cans and dumpsters; and other structural elements where listed pests congregate. Also spray the interior and exterior surfaces of garbage cans and dumpsters. For ants, pharaoh ants, and carpenter ants, also spray trails, nests and areas of entry (e.g., around windows and doors).

For use in canneries, food warehouses and dried fruit storage areas to kill Cadelles, Rice weevils, Granary weevils, Flat grain beetles, Cigarette beetles, Confused flour beetles, Drugstore beetles, Lesser grain borer, Merchant grain beetles, Sawtoothed grain beetles, Warehouse beetles, Red flour beetles, Grain mites, Cheese mites, Yellow mealworm: After sanitation measures have been performed, thoroughly

treat indoor structural surfaces, including walls, floors and ceilings; empty cabinets, drawers and shelves; food processing and handling equipment (not in operation); and other structural elements where listed pests congregate, directing the spray into cracks and crevices. Also spray localized areas around cartons or stored product containers. Treated cabinets, drawers and shelves or equipment must be dry before replacing stored items or before operating.

For use in grain storage sites of grain warehouses, grain elevators, rice mills, wheat mills, seed warehouses, farms and listed modes of transport to kill the stored product pests listed above: This product can be used in warehouse bins, grain or hopper bins, truck beds and cargo ships' holds as a cleanup prior to using them for storage or for transit. This product can also be used to treat conveying equipment.

In mills and elevators, remove all accumulations of infested grain from bins. Thoroughly clean walls, rafters, floors and door frames, and all storage and conveying equipment by sweeping or vacuuming the waste grain, cobwebs and other debris, with special attention to material lodged in cracks and crevices. Remove and burn all of the debris to kill eggs and insects that might be present.

For farms, give particular attention to cleaning up around used feed and grain bags, grain residues from wagons, harvesting equipment and feed troughs. **DO NOT** place newly harvested grain in the same bin with carry-over grain. All carry-over grain stocks that are not treated with grain protectant should be fumigated using a registered product intended for this purpose. Perform these cleaning operations within two or three weeks before harvest. After above sanitation measures have been performed and prior to use for storage or for transit, thoroughly treat conveying equipment (not in operation) and indoor surfaces of empty structures, including walls, floors and ceilings, and along partition boards, paying particular attention to forcing the spray into all cracks and crevices. Treated warehouse bins, equipment and surfaces of listed storage structures must be dry prior to use for storage or for loading prior to transit.

For use in poultry houses and on poultry to kill poultry lice and darkling beetles: Thoroughly spray roosts, walls and nests or cages. **DO NOT** exceed 1 second per m². After treating the structural surfaces of the poultry house, spray product over the poultry. **DO NOT** exceed 5-10 seconds per 30 m³.

For use in poultry houses and on poultry to kill bed bugs and mites: Thoroughly spray crevices of roost poles, cracks in walls and cracks in nests where the bedbugs and mites hide. **DO NOT** exceed 1 second per m². After treating the structural surfaces of the poultry house, spray product over the poultry. **DO NOT** exceed 5-10 seconds per 30 m³.

FOR OUTDOOR USE TO KILL ADULT MOSQUITOES, BITING MIDGES (NO-SEEUMS) AND FLIES

For use in listed outdoor areas prior to occupation to reduce annoyance from listed pests. Apply as a space spray around patios, porches and gazebos of listed structures and around surrounding vegetation, including shrubbery and tall grasses, where listed pests may rest. To enhance effectiveness of this product, apply directly to the listed pest where possible. For best results, treat when the wind velocity does not exceed 8 km per hour. Apply only when the potential for drift beyond the area to be treated is minimal. Take into consideration wind speed, wind direction and temperature inversions. Treatment must only be repeated after 7 days if pest problem persists or reoccurs.

FOR USE ON LISTED LIVESTOCK

Use Limitations

- **DO NOT** exceed 6 seconds per animal per treatment.
- **DO NOT** treat eyes/eyelids and mucous membranes (e.g., nose, mouth, genitals).
- **DO NOT** use on sick animals or animals under drug or pesticide treatment. Before use on pregnant animals, consult a veterinarian.
- Discontinue use if animal develops irritation after treatment with this product.
- When applied to dairy cattle: Treat dairy cattle at least 20 minutes before milking. **DO NOT** spray directly on udders. Wash udders before milking.
- When applied to horses: **DO NOT** cover horses immediately after treating/after treatment. **DO NOT** apply to back of horses prior to saddling.
- **DO NOT** contact treated animals until sprays have dried.

To protect listed livestock from attacks and to kill stable flies, horse flies, deer flies, horn flies, house flies, lesser house flies, mosquitoes and biting midges: Apply a light, uniform spray at the rate of 3 seconds on each side of the animal, taking care to treat back, withers and forelongs.

Thoroughly coat the hair of the animal – **DO NOT** wet the skin. **DO NOT** apply as a spray to the face of the animal. To treat the face, moisten (not to the point of dripping) a clean, soft cloth with product and thoroughly rub the moistened cloth on the face of the animal, being certain to avoid contact with the eyes/eyelids and mucous membranes. **DO NOT** apply more than once per day during peak fly season. **DO NOT** apply more than once a week during non-peak fly season. Treatment must only be repeated if pest problem persists or reoccurs.

To protect listed livestock from face flies: **DO NOT** apply as a spray to the face of the animal. Moisten (not to the point of dripping) a clean, soft cloth with product and thoroughly rub the moistened cloth on the face of the animal, being certain to avoid contact with the eyes/eyelids and mucous membranes. **DO NOT** apply more than once per day during peak fly season. **DO NOT** apply more than once a week during non-peak fly season. Treatment must only be repeated if pest problem persists or reoccurs.

To kill biting and sucking lice and to kill fleas and Ixodid ticks on listed livestock: Apply a light, uniform spray at the rate of 3 seconds per animal to infested areas of the animal (including brush of the tail, if applicable), except the face. Thoroughly coat the hair of the animal – **DO NOT** wet the skin. Use a stiff brush to get the spray down to the base of the hair. To treat the face of the animal, moisten (not to the point of dripping) a clean, soft cloth with product and thoroughly rub the moistened cloth on the face of the animal, being certain to avoid contact with the eyes/eyelids and mucous membranes. Repeat treatment in 10 days to kill newly hatched lice. Treatment must only be repeated if pest problem persists or reoccurs.

To kill sheep "tick" or ked on sheep: Apply a light, uniform spray at the rate of 3 seconds per sheep to cover all portions of the sheep's body, except the face. Thoroughly coat the wool of the sheep – **DO NOT** wet the skin. To treat the face of the sheep, moisten (not to the point of dripping) a clean, soft cloth with product and thoroughly rub the moistened cloth on the face of the sheep, being certain to avoid contact with the eyes/eyelids and mucous membranes. Treatment must only be repeated after 7 days if pest problem persists or reoccurs.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Avoid inhalation of spray. Avoid contact with skin and eyes and clothing. Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. Wash after using. **DO NOT** use this product in or on electrical equipment due to possibility of shock hazard.

ENVIRONMENTAL PRECAUTIONS:

This product contains active ingredients and aromatic petroleum distillates which are toxic to aquatic organisms. Toxic to bees. Avoid application around blooming plants. If applications must be made during the blooming period, restrict applications to evening when most bees are not foraging. Toxic to beneficial arthropods (which may include predatory and parasitic insects, spiders, and mites). Minimize exposure to non-target areas.

Contents under pressure. Do not place in hot water or near radiators, stoves, or other sources of heat. Do not puncture or incinerate container or store at temperatures over 50°C. Do not spray near open flame, spark, or while smoking.

FIRST AID: If swallowed, call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person. If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. If on skin or clothing, take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice. If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: This product contains petroleum distillates. Vomiting may cause aspiration pneumonia. Treat symptomatically.

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, PYROCID® Pressurized Multi-Purpose Spray 7416 contains a Group 3 insecticide. Any insect population may contain individuals naturally resistant to PYROCID® Pressurized Multi-Purpose Spray 7416 and other Group 3 insecticides. The resistant individuals may dominate the insect population if this group of insecticides is used repeatedly in the same locations. Other resistance mechanisms that are not linked to site of action but are specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed. To delay insecticide resistance:

- Where possible, rotate the use of PYROCID® Pressurized Multi-Purpose Spray 7416 or other Group 3 insecticides with different groups that control the same pests in the location to be treated.
- Insecticide use should be based on an IPM program that includes scouting and record keeping, and considers cultural, biological and other chemical control practices.
- Monitor treated pest populations for resistance development.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact MGK at (763) 544-0341.

STORAGE: Contents under pressure. **DO NOT** use or store near heat or open flame. **DO NOT** puncture or incinerate container. Exposure to temperatures above 50°C may cause bursting. Store this product away from food or feed. Store in a cool, dry place. Keep container closed.

DISPOSAL: When container is empty, press valve to release all remaining pressure. Dispose of empty container in accordance with provincial requirements. **DO NOT** puncture or incinerate. Dispose of the container in accordance with provincial requirements. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

EMERGENCY TELEPHONE NUMBER:

In case of Medical emergencies or health and safety inquiries, or in case of fire, leaking or damaged containers, information may be obtained by calling **1-888-740-8712**.