

2016-7138

2016-12-20

(FRONT OF IFC)

ACARICIDE

PREVENTIC®

FOR DOGS

Neck Sizes up to 62 cm

TICK COLLAR FOR

DOGS

DOMESTIC

GUARANTEE:

Amitraz: N'-(2,4 dimethylphenyl)-N-[(2,4 dimethylphenyl)imino]methyl}-N-methylmethanimide  
.....9.0%

KEEP OUT OF REACH OF CHILDREN

WARNING: MAY CONTAIN SOYBEAN OIL

READ THE BOX LABEL BEFORE USING

REG. NO. 24496

PCP ACT

Made in France for: (Virbac Logo)

Virbac AH, Inc. ▪ 3200 Meacham

Blvd.

Fort Worth, Texas 76137 ▪ 1-817-831-5030

NET CONTENTS 27.5 G

Tick Collar for Dogs

Adjustable – fits up to 62 cm neck size.

---

PRECAUTIONS: HUMANS: KEEP OUT OF REACH OF CHILDREN. DO NOT ALLOW CHILDREN TO

HANDLE THIS COLLAR. Dust will form on the collar during storage.

Dust may irritate eyes, nose and throat. Harmful if collar is sucked or chewed, or if dust is inhaled or absorbed through skin. Avoid contact with mouth, eyes, skin and clothing. Avoid breathing dust from collar. After handling or applying, wash hands (or any other skin that came into contact with the product) with soap and water.

PETS: DO NOT APPLY TO CATS. Avoid product contact with food, food utensils or places where food is prepared or stored. Dust may irritate eyes, nose and throat. Harmful if collar is chewed or swallowed. Avoid contact with the dog's mouth, face, eyes and genitals. Avoid letting the pet inhale the dust from the collar. Remove collar if signs of adverse reaction or skin irritation occur and shampoo dog thoroughly to remove residue. There is a risk of collar breakage and ingestion if a collar is worn by unattended dogs that play by mouthing each other in the neck area. If collar is missing from a dog, search for, retrieve and recover the entire collar. Sensitivities may occur after using any pesticide for pets. Smaller animals may be more likely to experience an adverse reaction. Observe carefully. Do not use on nursing animals or on animals under 12 weeks of age. Remove collar if signs of adverse reaction or skin irritation occur and shampoo dog thoroughly to remove residue.

FIRST AID: Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

2016-7138

2016-12-20

**HUMANS:** IF SUCKED OR CHEWED, obtain medical attention or call a poison control centre immediately. IF UNINTENTIONAL SKIN CONTACT OCCURS, wash affected areas with soap and water. IF DUST FROM COLLAR GETS IN EYES, flush eyes with plenty of water. If skin or eye irritation persists, obtain medical attention. IF DUST FROM COLLAR IS INHALED, seek medical attention.

**PETS:** Sensitivities may occur after using ANY pesticide for pets. If signs of sensitivity occur remove collar and bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately. If swallowed or chewed, consult a veterinarian.

**TOXICOLOGICAL INFORMATION:** If a dog ingests a Preventic collar, Yohimbine, 0.1 mg/kg by intravenous injection is antidotal. Repeat as needed.

**DIRECTIONS FOR USE:**

DETACHES, INHIBITS FEEDING AND KILLS AMERICAN AND BROWN DOG TICKS. DO NOT APPLY

MORE THAN ONE COLLAR PER DOG. Replace this solar no more than once every 90 days. A veterinarian should be consulted when pest infestation continues to be a problem on the pet.

Place the Preventic® collar around the dog's neck, buckle and adjust for secure fit ensuring that collar is worn tightly enough that the dog cannot remove it. The collar must be worn loosely so that two fingers may be placed between the collar and the dog's neck. If the collar is worn too tightly, it may produce neck irritation. Cut off and dispose of excess length. Do not allow dog or children to play with the cut-off excess. Activity against ticks starts within 24 hours and continues for up to 90 days. Ticks present in the dog's environment that may reappear on your pet will be prevented from feeding and killed by the action of this collar. Replace collar when effectiveness diminishes. Remove collar before bathing the dog. The Preventic® collar is not effective against fleas. Therefore, use other means for control of fleas. For effective tick control, treatment of the pet should be combined with sanitation of any area used by the pet. Vacuum floors, carpets and furniture (discard vacuum bag after use) and wash the pet's bedding, living quarters and surrounding areas. If pest problems persist, an insecticidal premise treatment may be required.

Do NOT remove the collar from its package until ready to use. Remove the collar from the package, unroll and stretch to activate the pesticidal ingredients. When collar is first worn, observe the pet's neck area every few days for irritation or adverse reaction. Do NOT use on dogs under 12 weeks of age. Consult a veterinarian before use on sick, aged, pregnant or nursing animals or animals receiving drug or other pesticide treatment. Do NOT use this product on dogs at the same time or within 30 days before or after treatment with, or exposure to, pesticides or chemicals.

RETAIN PACKAGING INFORMATION FOR FUTURE REFERENCE.

**STORAGE:** Store in a cool, dry area.

**DISPOSAL:** Do not re-use empty pouch, box or collar. Wrap in a newspaper and put in trash collection.