

Advantage™ 20 Flea and Lice Adulticide
Insert (English)

<Elanco Logo>

Advantage™ 20

Flea and Lice Adulticide

For dogs 8 weeks and older weighing between 4.6 and 11 kg

Once-A-Month Topical Flea and Lice Treatment

LIQUID

READ THE LABEL AND INSERT BEFORE USING

FOR USE ONLY ON DOGS

Do not use on other animals

KEEP OUT OF REACH OF CHILDREN

WARNING



POISON
EYE IRRITANT

DOMESTIC

ACTIVE INGREDIENT: Imidacloprid 9.1%

REG. NO. 25132 P.C.P. ACT

One to Six 1.0 mL Tubes

46 Advantage 20 Flea and Lice Adulticide provides dual protection against fleas and lice.
47 Advantage 20 Flea and Lice Adulticide will control fleas and lice on dogs within one day
48 of treatment and will continue to protect the dog from further flea and lice infestations for
49 at least four weeks. Reinfestation from the emergence of new fleas in the environment
50 may continue to occur for six weeks or longer after treatment is initiated. Reapply every
51 4 weeks for continual pest control in situations where there is an ongoing risk of pest
52 infestation. Treatment must only be repeated if pest problem reoccurs.

53
54 Shampooing of the dog may shorten the duration of flea and lice protection, therefore,
55 re-application following shampooing may be necessary. For effective flea and lice
56 control, treatment of the pet should be combined with the sanitation of any area used by
57 the pet. Vacuum floors, carpets and furniture (discard vacuum bag after use) and wash
58 the pet's bedding, living quarters and surrounding areas. If pest problems persist an
59 insecticidal premise treatment may be required.

60
61 The successive feeding activity of fleas on pets may elicit a hypersensitivity skin
62 disorder known as flea allergy dermatitis (FAD). Treatment of pets with Advantage 20
63 Flea and Lice Adulticide rapidly kills fleas and may reduce the incidence of this
64 condition. Advantage 20 Flea and Lice Adulticide is safe to use concurrently with
65 commonly used veterinary products. Consult your veterinarian before using on animals
66 receiving drug treatment. For use on pregnant/lactating animals consult your
67 veterinarian.

68
69 **PRECAUTIONS:**

70
71 **HUMANS: KEEP OUT OF THE REACH OF CHILDREN.** Harmful if swallowed, or
72 absorbed through the skin. Causes eye irritation. DO NOT get in eyes. Avoid contact
73 with skin and clothing. Wear rubber gloves when applying product. After handling or
74 applying, wash hands (or any other skin that came into contact with the product) with
75 soap and water. Avoid contact with pets until dry. Do not contaminate water, feed or food
76 by storage or disposal.

77
78 **PETS:** Do not use on animals under 8 weeks of age. Care should be taken to prevent
79 exposure of other pets who may be in close physical contact with the treated animal.
80 Individual sensitivities, while rare, may occur after using ANY pesticide product for pets.
81 Smaller animals are more likely to experience an adverse reaction. Observe carefully. If
82 signs persist, or become more severe, consult a veterinarian immediately.

83

| |
|---|
| Side Effects: Monitor your dog after application. Side effects may include: skin irritation such as redness or scratching; changes in behaviour such as agitation or lethargy; or gastrointestinal effects such as vomiting or loss of appetite. If these or other side effects occur, consult your veterinarian or Elanco at 1-800-265-5475. |
|---|

84
85 **FIRST AID:**

86
87 **HUMANS: IF IN EYES,** hold eye open and rinse slowly and gently with water for 15-20

88 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue
89 rinsing eye.

90 Call a poison control center or doctor for treatment advice.

91 IF SWALLOWED, call a poison control center or doctor immediately for treatment
92 advice. Have person sip a glass of water if able to swallow. Do not induce vomiting
93 unless told to do so by a poison control center or doctor. Do not give anything by mouth
94 to an unconscious person.

95 IF ON SKIN OR CLOTHING, take off contaminated clothing. Rinse skin immediately
96 with plenty of water for 15-20 minutes. Call a poison control center or doctor for
97 treatment advice.

98
99 PETS: If signs of sensitivity occur, bathe the pet with mild soap and rinse with large
100 amounts of water. If signs continue, consult a veterinarian immediately. Take container,
101 label or product name and Pest Control Product Registration Number with you when
102 seeking medical or veterinary attention.

103
104 TOXICOLOGICAL INFORMATION: Treat the patient symptomatically.

105
106 STORAGE: Store container at room temperature.

107
108 DISPOSAL: Do not reuse empty container. Wrap empty tube and cap in newspaper and
109 dispose with household garbage.

110
111 DIRECTIONS FOR USE:

112 This product contains imidacloprid. Do not apply another pest control product such as a
113 shampoo, collar, or powder that contains imidacloprid to the treated animal after
114 applying Advantage 20 Flea and Lice Adulticide.

- 115 • Do NOT administer by mouth. Do NOT use on other animals. Consult a
116 veterinarian before using on sick, aged, pregnant or nursing animals or animals
117 receiving drug or other pesticide treatment. A veterinarian should be consulted
118 when pest infestation continues to be a problem on the pet.
- 119 • Treatment must only be repeated if pest problem reoccurs.
- 120 • Do not use on puppies under 8 weeks of age.
- 121 • Use only on dogs weighing between 4.6 to 11 kg.
- 122 • This product is packaged in single dosage tubes. This package contains 1 to 6
123 monthly 1.0 mL dosages. Treat your dog monthly in the manner described below.
124 Do not apply a dosage higher than one 1.0 mL tube at a time.
- 125 • Remove one applicator from the package.
- 126 • Hold applicator tube in upright position. Pull cap off tube.
- 127 • Turn cap around and place other end of cap back on tube.
- 128 • Twist cap to break seal, then remove cap from tube.
- 129 • The dog should be standing for easy application. The entire contents of the
130 Advantage 20 Flea and Lice Adulticide tube should be applied evenly to two or
131 three spots on the top of the back from the shoulder to the base of the tail. At
132 each spot, part the hair until the skin is visible. Place the tip of the tube on the
133 skin and gently squeeze the tube to dispense a portion of the solution on the

2022-0231
2022-01-31

- 134 skin. Do not apply an excessive amount of solution at one spot that could cause
135 some of the solution to run off the side of the dog. Do not get this product in your
136 pet's eyes or mouth.
- 137 • Until Advantage 20 Flea and Lice Adulticide dries, keep pet off furniture and
138 away from hardwood floors.
 - 139 • Discard the empty tube as described in "DISPOSAL".

140

141 RETAIN PACKAGING INFORMATION FOR FUTURE REFERENCE.

142

143 Elanco Canada Limited
144 1919 Minnesota Court, Suite 401
145 Mississauga, Ontario L5N 0C9

146

147 Advantage, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.

148 ©2022 Elanco or its affiliates.

149

150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174

175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194

Advantage™ 20 Flea and Lice Adulticide
Carton (English)

<Elanco Logo>

Advantage™ 20

Flea and Lice Adulticide

For dogs 8 weeks and older weighing between 4.6 and 11 kg

Once-A-Month Topical Flea and Lice Treatment

LIQUID

READ THE LABEL AND INSERT BEFORE USING

FOR USE ONLY ON DOGS

Do not use on other animals

KEEP OUT OF REACH OF CHILDREN

WARNING



POISON
EYE IRRITANT

DOMESTIC

ACTIVE INGREDIENT: Imidacloprid 9.1%

REGISTRATION NO. 25132 P.C.P. ACT

One to Six 1.0 mL Tubes

Elanco Canada Limited
1919 Minnesota Court, Suite 401
Mississauga, Ontario L5N 0C9
1-800-265-5475

195 Advantage 20 Flea and Lice Adulticide provides dual protection against fleas and lice.
196 Advantage 20 Flea and Lice Adulticide will control fleas and lice on dogs within one day
197 of treatment and will continue to protect the dog from further flea and lice infestations for
198 at least four weeks. Reinfestation from the emergence of new fleas in the environment
199 may continue to occur for six weeks or longer after treatment is initiated. Reapply every
200 4 weeks for continual pest control in situations where there is an ongoing risk of pest
201 infestation. Treatment must only be repeated if pest problem reoccurs.

202
203 Shampooing of the dog may shorten the duration of flea and lice protection, therefore,
204 re-application following shampooing may be necessary. For use on pregnant/lactating
205 animals consult your veterinarian.

206
207 **PRECAUTIONS:**

208
209 **HUMANS: KEEP OUT OF THE REACH OF CHILDREN.** Harmful if swallowed, or
210 absorbed through the skin. Causes eye irritation. **DO NOT** get in eyes. Avoid contact
211 with skin and clothing. Wear rubber gloves when applying product. After handling or
212 applying, wash hands (or any other skin that came into contact with the product) with
213 soap and water. Avoid contact with pets until dry. Do not contaminate water, feed or
214 food by storage or disposal.

215
216 **PETS:** Do not use on animals under 8 weeks of age. Care should be taken to prevent
217 exposure of other pets who may be in close physical contact with the treated animal.
218 Individual sensitivities, while rare, may occur after using ANY pesticide product for pets.
219 Smaller animals are more likely to experience an adverse reaction. Observe carefully. If
220 signs persist, or become more severe, consult a veterinarian immediately.

221
222

| |
|---|
| Side Effects: Monitor your dog after application. Side effects may include: skin irritation such as redness or scratching; changes in behaviour such as agitation or lethargy; or gastrointestinal effects such as vomiting or loss of appetite. If these or other side effects occur, consult your veterinarian or Elanco at 1-800-265-5475. |
|---|

223 **FIRST AID:**

224
225 **HUMANS: IF IN EYES,** hold eye open and rinse slowly and gently with water for 15-20
226 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue
227 rinsing eye.

228 Call a poison control center or doctor for treatment advice.

229 **IF SWALLOWED,** call a poison control center or doctor immediately for treatment
230 advice. Have the person sip a glass of water if able to swallow. Do not induce vomiting
231 unless told to do so by a poison control center or doctor. Do not give anything by mouth
232 to an unconscious person.

233 **IF ON SKIN OR CLOTHING,** take off contaminated clothing. Rinse skin immediately
234 with plenty of water for 15-20 minutes. Call a poison control center or doctor for
235 treatment advice.

236

2022-0231
2022-01-31

237 PETS: If signs of sensitivity occur, bathe the pet with mild soap and rinse with large
238 amounts of water. If signs continue, consult a veterinarian immediately. Take container,
239 label or product name and Pest Control Product Registration Number with you when
240 seeking medical or veterinary attention.

241

242 TOXICOLOGICAL INFORMATION: Treat the patient symptomatically.

243

244 STORAGE: Store container at room temperature.

245

246 DISPOSAL: Do not reuse empty container. Wrap empty tube and cap in newspaper and
247 dispose with household garbage.

248

2022-0231
2022-01-31

249
250
251
252
253
254
255
256
257
258

259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276

Advantage™ 20 Flea and Lice Adulticide
Tube (English)

<Elanco Logo>

Advantage™ 20

READ THE LABEL AND INSERT BEFORE USING

WARNING



POISON
EYE IRRITANT

DOMESTIC

EXTERNAL USE ONLY

ACTIVE INGREDIENT: IMIDACLOPRID 9.1%

REG. NO. 25132 P.C.P. ACT

Elanco Canada Limited
1919 Minnesota Court, Suite 401
Mississauga, Ontario L5N 0C9

1.0 mL