**BOTTLE** 

# LATER'S SEVIN® LIQUID INSECTICIDE

For Lawns, Ornamentals, Vegetables and Fruits.

## **DOMESTIC**

REG. NO. 23860 P.C.P. ACT

GUARANTEE: Carbaryl 22.5% Contains 1,2-benzisothiazolin-3-one at 0.0193% as a preservative.

**CAUTION - POISON** 

250 mL

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

Sure-Gro IP Inc. P.O. Box 21001, Brantford, ON N3R 7W9

## READ ENCLOSED BOOKLET BEFORE USING

**PRECAUTIONS:** KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed or inhaled. Do not breathe spray mist. Avoid skin and eye contact. Wear long-sleeved work clothing. Wash hands and face before eating or drinking. Avoid contamination of feed, food or water supplies.

FIRST AID: If swallowed, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person. If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. If on skin or clothing, take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and P.C.P. Reg. No. with you when seeking medical attention.

**TOXICOLOGICAL INFORMATION:** Sevin is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

# LATER'S SEVIN® LIQUID INSECTICIDE

For Lawns, Ornamentals, Vegetables and Fruits.

# **DOMESTIC**

REG. NO. 23860 P.C.P. ACT

GUARANTEE: Carbaryl 22.5% Contains 1,2-benzisothiazolin-3-one at 0.0193% as a preservative.

CAUTION - POISON

250 mL

READ THE LABEL BEFORE USING KEEP OUT OF REACH OF CHILDREN

Sure-Gro IP Inc. P.O. Box 21001, Brantford, ON N3R 7W9

## **DIRECTIONS:**

**Later's Sevin**<sup>®</sup> **Liquid** is effective for the control of many insects on Lawns, Ornamental Plants, Vegetables and Fruits when used as directed. Apply when insects or damage appears. Spray in early morning or late evening to avoid injury to bees and to crops from spraying during high temperature. Spray to point of run-off, monitor insect activity and repeat applications at 7-10 day intervals if necessary.

## **SPRAY PREPARATION:**

**Later's Sevin® Liquid** readily disperses in water to form a spray suspension suitable for home and garden use. Shake container thoroughly before using. Prepare only as much spray mixture as is needed on the day of application. Shake spray mixture frequently before and during use to prevent settling. Do not store spray mixture overnight.

5 mL = 1 teaspoonful 15 mL = 1 tablespoon 1 L = approx. 1 quart  $10 \text{ m}^2 = \text{approx}$ . 100 sq. ft.

#### **HOW TO USE:**

## On Fruit:

Mix 5 mL of **Sevin® Liquid** with 1 L of water (or 5 teaspoons / 1 gal.). Apply for thorough coverage to the point of run-off. Do not spray within 7 days of harvest.

Use on: Apples, Apricots, Cherries, Peaches, Pears, Plums.

To Control: Aphids, Apple Maggots, Coding Moths, Cherry Fruit Flies, Fruitworms,

Leafhoppers, Leafminers, Leafrollers, Mealybugs, Pear Slugs, Scale insect

(crawlers), Tent Caterpillars.

## On Berries & Grapes:

Mix 8 mL of **Sevin Liquid** with 1 L of water to cover 10 m<sup>2</sup> (or 8 teaspoons / 1 gal./500 sq. ft.). Do not spray within 7 days of harvest.

Use on: Blackberries, Blueberries, Boysenberries, Loganberries, Raspberries, Strawberries

and Grapes.

To Control: Climbing Cutworms, Leafhoppers, Leafrollers, Fruitworms, Spittlebugs.

# On Vegetables:

Do not spray within 7 days of harvest or within 14 days of leafy vegetables.

Use on: Asparagus, Beans, Beets, Broccoli, Brussels Sprouts, Cabbage, Carrots, Cauliflower,

Celery, Chinese Cabbage, Corn, Cucumber, Eggplant, Endive, Horseradish, Kale, Kohlrabi, Lettuce, Melons, Parsley, Parsnip, Pepper, Potato, Pumpkin, Radishes,

Rutabagas, Spinach, Squash, Swiss Chard, Tomato, Turnip, Watercress.

To Control: Asparagus, Beetles, Fleabeetles, Leafhoppers, Mexican Bean Beetles, Colorado

Potato Beetles, Corn Earworms, Corn Borers, Cucumber Beetles, Squash Bugs.

Mix 5 mL of **Sevin Liquid** with 1 L of water to cover 10 m<sup>2</sup> (or 5 teaspoons/1 gal./500 sq. ft.)

To Control: Armyworms, Fruitworms, Tomato Hornworms, Stinkbugs, mix 7 mL of Sevin®

**Liquid** with 1 L of water to cover 10 m<sup>2</sup> (or 7 teaspoons/1 gal./500 sq. ft.).

To Control: Climbing Cutworms, mix 10 mL of **Sevin**<sup>®</sup> **Liquid** with 1 L of water to cover 10 m<sup>2</sup>

(or 10 teaspoons/1 gal./500 sq. ft.).

## **On Ornamentals:**

Mix 5 mL of **Sevin® Liquid** with 1 L of water (or 5 teaspoons/1 gal.). Apply for thorough coverage to the point of run-off.

Use on: Arborvitae, Azalea, Birch, Boxwood, Carnation, Chrysanthemum, Dogwood, Elm,

Gladiolus, Holly, Hydrangea, Juniper, Lilac, Maple, Oak, Pines, Rose, Zinnia.

To control: Rose Aphids, Bagworms, Birch Leafminers, Black Vine Weevils, Blister Beetles,

Boxelder Bugs, Boxwood Leafminers, Cankerworms, Elm Leaf Beetles, Flea Beetles, Gypsy Moths, Japanese Beetles, June Beetles, Lace Bugs, Leaf Hoppers, Leaf Rollers, Mealybugs, Pine Saw Flies, Plant Bugs, Psyllids, Rose Chafer, Rose Slugs, Scale Insects (crawlers), Spruce Budworms, Tent Caterpillars, Thrips

(exposed), Willow Leaf Beetles.

Note: Do not use on Boston Ivy, Virginia Creeper and Maidenhair Ferns. During early season, it may also injure Virginia and sand Pines. For Scale Insects, apply when crawlers are present.

# On Lawns:

To Control: Ants, Chinch Bugs, Earwigs, Fall Armyworms, Fleas, Leaf Hoppers, Millipedes, Mosquitoes, Sod Webworms.

Mow lawn and apply at the rate of 50 mL of  $Sevin^{®}$  Liquid with 20 L of water to cover 10 m<sup>2</sup> of area (or 4 tablespoons/4 gal./100 sq. ft.). Do not water for 2 days after treatment. Repeat application in 2-3 weeks if necessary.

**PRECAUTIONS:** KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed or inhaled. Do not breathe spray mist. Avoid skin and eye contact. Wear long-sleeved work clothing. Wash hands and face before eating or drinking. Avoid contamination of feed, food or water supplies.

**ENVIRONMENTAL HAZARDS:** Do not contaminate streams and ponds. Highly toxic to bees. Do not use where bees are foraging. Some damage may occur to tender foliage in the presence of rain or high humidity of several days duration following spraying.

**FIRST AID:** If swallowed, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person. If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. If on skin or clothing, take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice. Take container, label or product name and P.C.P. Reg. No. with you when seeking medical attention.

**TOXICOLOGICAL INFORMATION:** Sevin<sup>®</sup> is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

**STORAGE:** Keep product in original container. Store in a cool, dry place away from food. DO NOT FREEZE.

**DISPOSAL:** Do not reuse empty container. Rinse thoroughly and use rinse-water with regular spray. Wrap container and discard in household garbage.

\*\*\*\*\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.