

2021-0663  
2021-03-22

**VET-KEM® Siphotrol® 1000 Flea & Tick Home Spray**

With PRECOR® (IGR) Insect Growth Regulator

FOR INDOOR USE ONLY

Kills fleas, brown dog and American dog ticks in the home  
7 months residual action against Pre-Adult Fleas

Double Action

Premise Treatment

Kills both Adult and Pre-Adult (larvae) Fleas

Treats surface area of 93 m<sup>2</sup> (1,000 sq. ft.)

KEEP OUT OF REACH OF CHILDREN  
READ THE LABEL BEFORE USING

DOMESTIC

ACTIVE INGREDIENT:

(S)-Methoprene ... 0.078%

Permethrin ... 0.80%

REG. NO. 25739 P.C.P. ACT

Warning contains the allergen soy.

CAUTION

SKIN IRRITANT

POTENTIAL SKIN SENSITIZER

EXPLOSIVE

Net Contents: 227 g

Wellmark International

100 Stone Road West, Suite 111

Guelph, Ontario N1G 5L3

1-800-688-7378

2021-0663

2021-03-22

## **DIRECTIONS FOR USE**

**General Information:** VET-KEM Siphotrol 1000 Flea & Tick Home Spray kills adult and pre-adult fleas and ticks in the home. VET-KEM Siphotrol 1000 Flea & Tick Home Spray contains a unique combination of two insect control ingredients, permethrin and Precor. The permethrin kills adult fleas, brown dog and American dog ticks. Precor, an insect growth regulator, breaks the flea life cycle by preventing the flea's development into the adult biting stage for 7 months.

This product is designed for use on carpeting, however, some natural fibres (such as wool) and synthetics may be adversely affected by any liquid product – always test a hidden area prior to use. Care should be used when treating wood surfaces, as water spotting may occur. Users should test (treat) a small, inconspicuous area first to ensure there are no adverse effects such as staining or discolouration. Good sanitary practices should be maintained to prevent re-infestation. One 227 g can will treat a structural surface area of 93 m<sup>2</sup> (2000 ft<sup>2</sup>).

**Restrictions:** FOR INDOOR USE ONLY. DO NOT contaminate food/feed. DO NOT apply to surfaces that may come into contact with food/feed. DO NOT apply when people or pets are present. DO NOT allow people or pets to enter treated areas until sprays have dried. DO NOT allow spray to drip or allow drift onto non-target surfaces. DO NOT apply to people, pets, or sources of electricity. Remove pets and birds, and cover or remove fish tanks or bowls. DO NOT apply to furniture, mattresses, linens, toys or clothing. DO NOT apply to overhead areas or in confined spaces (for example, attics, crawlspaces). Ventilate treated areas after application by opening windows and doors or using fans, where required, to aid in the circulation of air.

DO NOT apply as a space spray or broadcast application. Apply as a crack and crevice spot or indoor perimeter treatment only. Crack and crevice is an application directly into narrow openings on the surface of the structure. It does not include the treatment of exposed surfaces. Narrow openings typically occur at expansion joints, utility entry points and along baseboards and mouldings. Spot treatment is localized to a surface area not more than 0.2 m<sup>2</sup>. Spots are not to be adjoining and the combined area of spots is not to exceed 10% of the total surface area of a room. Indoor perimeter application is less than 0.3 m wide along the edges of a room to baseboards, wall-floor and ceiling-wall joints, and around doorways or windows.

**Application Instructions for fleas and ticks:** For effective flea and tick control indoors, clean any area used by the pet. Cleaning the areas includes vacuuming floors, carpets, furniture, etc. (discard vacuum bag after use) and washing the pet's bedding, living quarters, and surrounding area. DO NOT TREAT PETS WITH THIS PRODUCT. Only treat pets with products specifically registered for use on that type of pet.

Step One - Prepare the Treatment Area: Vacuum floors, carpets, and furniture (discard vacuum bag after use) and wash the pet's bedding, living quarters and surrounding area prior to treatment with this product. For effective flea and tick control indoors, clean any area used by the pet. Cleaning the areas includes vacuuming floors, carpets, furniture, etc. (discard vacuum bag after use) and washing the pet's bedding, living quarters, and surrounding area.

Step Two - Spray the Area: SHAKE WELL BEFORE USING. This product can only be applied as a spot or indoor perimeter application to cracks and crevices in walls and floors, along and behind baseboards, window and door frames, and localized areas of floor and floor coverings (e.g., rugs, carpets). Pet bedding may be entirely treated with this product or replaced with clean, fresh bedding after other listed areas are treated. When applying this product to these locations, (point can downward 90 cm from the surface to be treated) and use a sweeping motion to apply a light, uniform spray. Within the specified use location, to enhance effectiveness of this product, apply this product directly to the listed pest where possible. If pest problems persists, a second application 2-3 weeks after the initial treatment may be necessary. Treatments must only be repeated after 2 weeks/ 14 days if pest problem persists or reoccurs.

2021-0663  
2021-03-22

### **PRECAUTIONS**

KEEP OUT OF REACH OF CHILDREN. May cause skin irritation and sensitization. Avoid contact with skin, eyes and clothing. Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing mist. After using, wash hands with soap and water.

### **FIRST AID**

Take container, label or product name and PCP Registration Number with you when seeking medical attention.

**If swallowed:** Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to mouth, if possible. Call a poison control centre or doctor for further treatment advice.

### **TOXICOLOGICAL INFORMATION**

Treat symptomatically.

### **STORAGE**

Contents under pressure. Do not puncture container or store near open flame or temperatures over 50°C.

### **DISPOSAL**

When container is empty, press button to release all pressure. Replace cap, wrap empty container and dispose with household garbage.

Hotline: 1-800-688-7378

Vet-Kem, Siphotrol and Precor are registered trademarks of Wellmark International.