

2023-2305
2024-02-16

This label has been updated according to the re-evaluation decision of Pyrethrins RVD2023-06 and Piperonyl Butoxide RVD2023-07. While users are encouraged to follow this updated label immediately, the previously approved label is valid until 03/02/2025 in accordance with the phase out period set out in RVD2023-06 and RVD2023-07. This previously approved label will be provided upon request by emailing hc.pmra.info-arla.sc@canada.ca. In your email please include the product name and Registration number of the label you are requesting.

2023-2305
2024-02-16

Can-Vet Barn and Dairy Fly Shield

CONTAINS PYRETHRINS AND PIPERONYL BUTOXIDE

AGRICULTURAL

For indoor use at dairy barns, milk rooms, beef barns, horse stables and swine housing to (kill/control) House flies, Face flies, Stable flies, Horn flies, Horse flies, Mosquitoes, Wasps, Hornets, Spiders, Ants, Cockroaches, and Crickets. For use on beef and dairy cattle (lactating and non-lactating) and swine to (kill/control) horse flies, horn flies, mosquitoes and stable flies. For use on beef and dairy cattle (lactating and nonlactating) to (kill/control) blood sucking lice.

READ THE LABEL BEFORE USING

ACTIVE INGREDIENTS:

Pyrethrin.....0.1%
Piperonyl Butoxide. 1.0%

Ready to Use

KEEP OUT OF REACH OF CHILDREN

REGISTRATION NO. 24159

PEST CONTROL PRODUCTS ACT

CAUTION POISON
{Skull & Bones Picture}

Net Contents
Can-Vet Animal Health Supplies Ltd.
4-61 Malcolm Rd Guleph, Ontario N1K 1A7

2023-2305
2024-02-16

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the Pest Control Products Act to use this product in a way that is inconsistent with the directions on the label.

GENERAL INFORMATION:

This product may be used in farm buildings and on livestock. Kills on contact: mosquitoes, wasps, hornets, spiders, ants, crickets, house flies, face flies, stable flies, horn flies, horse flies and sucking lice.

DIRECTIONS FOR USE:

For maximum effectiveness, a combination of surface spot treatment and fogging is recommended.

SPACE SPRAY: Space application is a suspension of fine droplets (0.1 to 100 µm) in the air within an indoor space. **Farm Buildings:** (dairy barns, milk rooms, beef barns, horse stable, swine housing). For rapid control of listed flies, mosquitoes, spiders, ants cockroaches, apply using a good space sprayer (Fogger) adjusted to deliver a fine mist such as a Fogmaster or U.L.V. equipment which dispenses a fine mist with droplets 15 - 25 microns in diameter. All animals must be removed from barn/buildings before fogging. Close windows, doors and shut off ventilation system and sources of ignition. Wear mask or respirator and protective clothing. Direct spray upwards to fill the room with a fine mist. Use a rate of 200-300 mL per 100 cubic metres. Leave the room closed for 15 minutes before ventilating. Do not use in dwellings. Ventilate and keep animals out of sprayed areas for at least 30 minutes after treatment. Spray only when temperature is under 32°C. Do not use in milking parlour areas.

Livestock: Spray on Beef, Dairy Cattle and Swine: for horse flies, horn flies, mosquitoes and stable flies, apply up to 30 mL per animal using sprayer which produces a fine mist. Cover the entire animal without making the hair wet. Do not spray in eyes or mouth. Do not treat more than once a day. Treat dairy cattle before milking, but do not spray directly on udder. Wash udder before milking.

To Control Blood Sucking Lice on Cattle: Apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat as necessary but not more than once a day. **CAUTION: If irritation appears, discontinue usage.**

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes and clothing. Wash thoroughly with soap and water after handling and before eating or smoking. DO NOT breathe spray mist. If irritation appears, discontinue usage. Wash contaminated clothing before reuse. DO NOT contaminate feed, food or water. Store in original container away from feed and food. Use only when milk room is NOT in operation. Cover or remove all milk and milking implements before spraying. After spraying, wash all surfaces that may contact milk and rinse thoroughly with potable water before resuming operations.

USE LIMITATIONS

- **DO NOT** apply when people or pets [or livestock] are present.
- **DO NOT** remain in treated areas after application.”
- **DO NOT** allow people or pets [or livestock] to enter treated areas until sprays have settled.
- **DO NOT** enter or allow entry into treated areas for 2 hours following application. The commercial applicator is responsible for informing workers and others of this requirement.
- Ventilate treated indoor areas after application by opening windows and doors or using fans, where required, to aid in the circulation of air.
- **DO NOT** apply when a food/feed processing facility, milking parlour or milk house is in operation.
- **DO NOT** apply to surfaces that may come into contact with food/feed. Cover or remove all

2023-2305
2024-02-16

- food/feed (including bulk tanks that contain milk). Cover all food/feed processing surfaces,³³⁵
- equipment and utensils or thoroughly wash them following treatment. Any food/feed (including milk) accidentally contaminated with spray solution must be discarded. Care should be taken to avoid the pesticide exiting the void. Any residue deposits on non-target surfaces must be removed by the applicator.
- Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair.
- When applying to overhead areas or in confined spaces, wear appropriate respiratory and eye protection.
- If entering treated indoor areas prior to venting, wear chemical-resistant coveralls over long-sleeved shirt, long pants, chemical-resistant hood, chemical-resistant footwear, socks, chemical-resistant gloves, and a respirator with a NIOSH-approved organic-vapour-removing cartridge with a prefilter approved for pesticides OR a NIOSH-approved canister approved for pesticides.
- Avoid contact with treated animals until dried.
- **DO NOT** allow spray to drip or allow drift onto non-target surfaces.

DISPOSAL: Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site: 1. Triple- or pressure-rinse the empty container. Add the rinsing to the spray mixture in the tank. 2. Make the empty, rinsed container unsuitable for further use. If there is no container collection site in your area, dispose of the container in accordance with provincial requirements accordance with provincial requirements. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

FIRST AID: If on skin or clothing, take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice. If in eyes, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. If swallowed, call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Treat symptomatically. This product contains a petroleum distillates.