

2024-02-20
2023-2157

This label has been updated according to the re-evaluation decisions for Pyrethrins RVD2023-06 and Piperonyl Butoxide RVD2023-07. While users are encouraged to follow this updated label immediately, the previously approved label is valid until March 2, 2025 in accordance with the phase out period set out in the re-evaluation decisions for Pyrethrins RVD2023-06 and Piperonyl Butoxide RVD2023-07. This previously approved label will be provided upon request by emailing pmra.info-arla@hc-sc.gc.ca. In your email please include the product name and Registration number of the label you are requesting.

Ovex

Pump Spray for Dogs & Cats with Precor® Insect Growth Regulator (IGR)

FOR USE ONLY ON DOGS AND CATS

Kills dog fleas, cat fleas, brown dog ticks and American dog ticks.

DO NOT USE ON PUPPIES OR KITTENS LESS THAN 12 WEEKS OLD

DO NOT USE ON ANY OTHER ANIMAL

DOMESTIC SOLUTION

ACTIVE INGREDIENTS:

(S)-Methoprene 0.10%

Pyrethrins..... 0.20%

Piperonyl Butoxide Technical..... 0.37%

N-Octyl Bicycloheptene Dicarboximide 0.61%

REG. NO. 23692 P.C.P. ACT

**READ THE LABEL BEFORE USING
KEEP OUT OF REACH OF CHILDREN**

**CAUTION
FLAMMABLE**

**WARNING
EYE IRRITANT**

Wellmark International
100 Stone Road West, Suite 111
Guelph, Ontario N1G 5L3
1-800-688-7378
Net Contents: 175-1000 ml

GENERAL INFORMATION: Kills adult cat fleas, dog fleas and American dog and brown dog ticks. Effectively breaks the flea life cycle and prevents the growth of the pre-adult stages of the flea. On dogs and cats, one treatment kills adult fleas for up to 7 days and keeps flea eggs from hatching for 30 days.

Use Limitations

DO NOT use on puppies or kittens less than 12 weeks old.

DO NOT USE ON ANY OTHER ANIMAL.

Consult a veterinarian before using on sick, aged, pregnant or nursing animals or animals receiving drug or other pesticide treatment.

Use **ONLY** in a well-ventilated area, preferably outdoors.

DO NOT administer by mouth.

Given the residual control afforded by Ovex, **DO NOT** use this spray more often than once every month. Treatment must only be repeated if pest problem persists or reoccurs.

This product contains piperonyl butoxide, N-Octyl bicycloheptene dicarboximide, s-methoprene and pyrethrins. **DO NOT** apply another pest control product such as a shampoo, spot on, collar or powder which contains piperonyl butoxide, N-Octyl bicycloheptene dicarboximide, s-methoprene or pyrethrins to the treated pet when using Ovex.

DIRECTIONS FOR USE:

FOR USE ONLY ON DOGS AND CATS

To Kill dog fleas, cat fleas, brown dog ticks and American dog ticks.

SHAKE WELL BEFORE USING. Retain packaging information for future reference.

Wet the dog's/cat's head with water to drive pests onto the body. Hold the can in an upright position from a distance of 20 cm from the dog/cat. Spray the entire body of the dog/cat, beginning at the head and working towards the tail. Ticks are difficult to control and must be sprayed directly. Pay special attention to neck, hindquarters, areas under legs, around the tail and between toes. **DO NOT** apply to dog's/cat's face, eyes, nose, mouth or genitals. As you spray, use hands to fluff the fur of the dog/cat so that spray will penetrate to the skin. Only spray until fur is damp – not wet. Wipe off any excess product from the dog/cat with a dry cloth and then comb or brush. Using this technique, the application of 2 to 4 squeezes (3 – 6 mls) of the trigger sprayer per kilogram will provide effective flea control. Bathing of the dog/cat may affect product efficacy. For effective flea and tick control, treatment of the pet should be combined with the sanitation of any area used by the pet. Vacuum floors, carpets and furniture (discard vacuum bag after use) and wash the pet's bedding, living quarters, and surrounding areas. If pest problems persist, a premise treatment with an insecticide labelled for that use may be required. The use of an effective product for controlling adult and larva flea populations in the pet's environment (e.g. Vet Kem Siphotrol® Forte, Vet Kem Siphotrol® 1000 or Vet Kem Siphotrol 2000) will reduce the possibility of re-infestation of the pet.

A veterinarian should be consulted when flea infestation continues to be a problem on the pet. **DO NOT** apply to surfaces that may come into contact with food/feed. Cover or

remove all food/feed. Cover all food/feed processing surfaces, equipment and utensils or thoroughly wash following treatment.

PRECAUTIONS:

Use only in well-ventilated areas, preferably outdoors.

DO NOT allow product to contact non-target surfaces.

HUMANS: KEEP OUT OF REACH OF CHILDREN. Avoid contact with treated pets until dried. Harmful if swallowed. Avoid contact with eyes, mouth, skin and clothing. Wear rubber gloves when applying. After handling or applying, wash hands (or any other skin that came into contact with the product) with soap and water.

PETS: Harmful if swallowed. Toxic to fish and other aquatic life. Cover fish aquaria in the vicinity of treatment area. Toxic to birds.

FIRST AID:

HUMANS: IF SWALLOWED, call a poison control centre or doctor immediately for treatment advice. **DO NOT** induce vomiting unless told to do so by a poison control centre or doctor. **DO NOT** give any liquid to the person. Do not give anything by mouth to an unconscious person. IF IN EYES, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. IF ON SKIN OR CLOTHING, take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice. IF INHALED, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

PETS: Sensitivities may occur after using ANY pesticide for pets. Sensitive cats may exhibit temporary and non-harmful salivation when exposed to pyrethrin containing products. In cases of known sensitivity, the use of the lower treatment rates indicated is recommended. If signs of sensitivity occur, bathe the pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

Take container, label or product name and PCP Registration Number with you when seeking medical or veterinary attention.

Side Effects:

Monitor your dog after application. Side effects may include: skin irritation such as redness or scratching; changes in behaviour such as agitation or lethargy; or gastrointestinal effects such as vomiting or loss of appetite. If these or other side effects occur consult your veterinarian or Wellmark International at 1 (800) 688-7378.

Monitor your cat after application. Side effects may include: skin irritation such as scratching or hair loss at the application site, or changes in behaviour such as agitation or lethargy. Gastrointestinal effects such as drooling, vomiting, or loss of appetite may also occur. If these or other side effects occur consult your veterinarian or Wellmark International at 1 (800) 688-7378.

TOXICOLOGICAL INFORMATION: Skin exposure may cause transient sensations (tingling, burning, itching, numbness). Other symptoms of exposure could include respiratory effects (such as cough, respiratory irritation, or shortness of breath), nausea, headache or dizziness. Treat symptomatically.

STORAGE: Store this product away from food or feed. Keep container in a cool location away from heat or open flame.

DISPOSAL: DO NOT re-use the empty container. Dispose in household garbage. Unused or partially used products should be disposed at provincially or municipally designated hazardous waste disposal sites.

PRECOR and SIPHOTROL are registered trademarks of Wellmark International.

HOTLINE 1-800-688-7378