

12-JAN-2010  
2009-3541

**LIBERTY® AT®**  
**HERBICIDE SUSPENSION**



<b>GROUP</b>	<b>5 10</b>	<b>HERBICIDE</b>
--------------	-------------	------------------

**FOR SALE FOR USE IN EASTERN CANADA  
DO NOT USE IN THE PROVINCE OF BRITISH COLUMBIA**

**COMMERCIAL**

**CAUTION - SKIN IRRITANT  
CAUTION - EYE IRRITANT**

**REGISTRATION NUMBER: 26968  
PEST CONTROL PRODUCTS ACT**

**GUARANTEE:**  
glufosinate ammonium 120 g/L  
atrazine 393 g/L  
related triazines 8 g/L

Contains 1,2-benzisothiazolin-3-one at 0.025% as a preservative

**READ THE LABEL AND ATTACHED BOOKLET BEFORE USING**

**Product Information: 1-888-283-6847**

**Bayer CropScience Inc.  
Suite 100, 3131 114<sup>th</sup> Ave. SE  
Calgary, Alberta T2Z 3X2**

**NET CONTENTS:  
10 - 118L**

**PRECAUTIONS:** CAUTION: EYE IRRITANT. CAUTION: POISON. CAUTION: SKIN IRRITANT. Harmful if inhaled. Do not inhale or ingest. May irritate eyes. DO NOT get into eyes. **KEEP OUT OF REACH OF CHILDREN.** Avoid contact with eyes, skin or clothing. Avoid breathing spray mist. Wash thoroughly after using and before eating, drinking or smoking. Mixing and loading is not permitted within 30 m from aquatic habitat. **Protective Clothing and Equipment:** During application, handling, mixing and loading, wear goggles or face shield, chemical resistant gloves, respirator, and coveralls over a long-sleeved shirt and long pants. Wear chemical resistant gloves for clean-up and repair. **Re-entry Restrictions:** Workers should not enter treated Corn fields within 24 hours of treatment. Workers who must enter fields within this time period should wear a long-sleeved shirt, long pants and chemical resistant gloves. If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's website at: [www.croplife.ca](http://www.croplife.ca). **FIRST AID:** Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention. **IF SWALLOWED:** Call a poison control centre or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person. **IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice. **IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice. **IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. **GENERAL INFORMATION:** Seek medical attention or contact a poison control centre immediately if irritation persists or is severe. **TOXICOLOGICAL INFORMATION:** Medical personnel should contact Bayer CropScience's medical information services toll-free 1-800-334-7577. **Note to Physician:** If this product is ingested, endotracheal intubation and gastric lavage should be performed as soon as possible, followed by charcoal and sodium sulfate administration. **Environmental Hazards:** Toxic to non-target terrestrial plants and aquatic organisms. Avoid over-spray or drift to sensitive/non-target aquatic habitats. An appropriate buffer zone is required between the downwind point of direct application and the closest edge of sensitive aquatic habitats including sloughs, coulees, ponds, prairie potholes, lakes, rivers, streams, reservoirs, and wetlands that are situated on the periphery of the treated area. Do not apply during periods of dead calm or when winds are gusty. Observe the buffer zones and precautionary measures specified under Directions for Use ( Section 9). Atrazine is persistent and will carry over. It is recommended that any products containing atrazine not be used in areas treated with this product during the previous season. Use good pesticide practices and apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and parks is minimal. Take into consideration wind speed, wind direction, temperature, application equipment and sprayer settings used for applications. For the protection of non-target habitats, over-spray or drift to any body of water or other

environmentally sensitive habitats must be avoided. The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions. Do not apply this product through any type of irrigation system. **USE ONLY ON LABELLED CROPS.** Avoid spray drift to susceptible plants and **USE ONLY FOR RECOMMENDED PURPOSES AND AT RECOMMENDED RATES.** **DO NOT APPLY BY AIR** Refer to Section 9 for buffer zones.

**Surface Runoff:** To reduce runoff from treated areas into aquatic habitats, consider the characteristics and conditions of the site before treatment. Site characteristics and conditions that may lead to runoff include, but are not limited to, heavy rainfall, moderate to steep slope, bare soil and poorly draining soil (e.g., soils that are compacted, fine textured or low in organic matter such as clay). Avoid applying this product when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip (buffer zone) between the treated area and the edge of the water body. **Leaching:** The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g., sand, loamy sand and sandy loam soils) and/or the depth to the water table is shallow.

**STORAGE:** CANNOT be stored below freezing. If stored for 1 year or longer, shake well before using. Store the tightly closed container away from feeds, seeds, fertilizers, plants and foodstuffs. Do not use or store in or around the home. Keep in original container during storage.

**DISPOSAL: Recyclable Container Disposal:** Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site: 1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank. 2. Make the empty, rinsed container unsuitable for further use. If there is no container collection site in your area, dispose of the container in accordance with provincial requirements. **Returnable Container Disposal:** Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer). **Refillable Container Disposal:** For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

**Disposal of Unused, Unwanted Product:** For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for cleanup of spills. **Disposal of**

**Unused Spray Solution:** If any spray solution remains in the tank after spraying is finished, it should be thoroughly diluted, then sprayed on the perimeter of the area just sprayed, away from water supplies, ditches, and irrigation canals. **NOTICE TO USER:** This pest control product is to be used only in accordance with the directions on label. It is an offence under the *Pest Control Products Act* to use a control product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

Liberty is a trademark of Bayer.

**LIBERTY® AT®**  
**HERBICIDE SUSPENSION**



GROUP	<b>5 10</b>	HERBICIDE
-------	-------------	-----------

**FOR SALE FOR USE IN EASTERN CANADA  
DO NOT USE IN THE PROVINCE OF BRITISH COLUMBIA**

**COMMERCIAL**

CAUTION - SKIN IRRITANT  
CAUTION - EYE IRRITANT

**REGISTRATION NUMBER: 26968**  
**PEST CONTROL PRODUCTS ACT**

**GUARANTEE:**  
glufosinate ammonium 120 g/L  
atrazine 393 g/L  
related triazines 8 g/L

Contains 1,2-benzisothiazolin-3-one at 0.025% as a preservative

**READ THE LABEL AND BOOKLET BEFORE USING**

**Product Information: 1-888-283-6847**

**Bayer CropScience Inc.**  
**Suite 100, 3131 114<sup>th</sup> Ave. SE**  
**Calgary, Alberta T2Z 3X2**

**NET CONTENTS:**  
**10 - 118L**

## LIBERTY AT HERBICIDE

<i>Table of Contents:</i>		<i>Section Number:</i>
<b>GENERAL INFORMATION</b>	The Product	1
<b>SAFETY AND HANDLING</b>	Precautions, Protective Clothing and Equipment, and Re-entry Restriction	2
	First Aid and Toxicological Information	3
	Environmental Hazards	4
	Storage	5
	Disposal	6
<b>DIRECTIONS FOR USE</b>	Crops, Weeds, Rates, Timing	7
	Herbicide Resistance Management	8
	Application Instructions and Cautions	9
	Mixing Instructions	10
	Sprayer Cleanup	11
	Livestock Feeding and Preharvest Intervals	12
	Notices	13

**Product Information:** 1-888-283-6847

**Bayer CropScience Inc.  
Suite 100, 3131 114<sup>th</sup> Ave. SE  
Calgary, Alberta T2Z 3X2**

## GENERAL INFORMATION

### Section 1: THE PRODUCT

LIBERTY AT Herbicide provides control of a broad spectrum of grassy and broadleaf weeds in Corn hybrids that are tolerant to glufosinate ammonium (for example LibertyLink® hybrids).

LIBERTY AT Herbicide may also be applied to glufosinate ammonium tolerant inbred Corn lines grown for seed production.

## SAFETY AND HANDLING

### Section 2: PRECAUTIONS, PROTECTIVE CLOTHING AND EQUIPMENT, AND RE-ENTRY RESTRICTIONS

#### PRECAUTIONS:

CAUTION: EYE IRRITANT

CAUTION: POISON

CAUTION: SKIN IRRITANT

Harmful if inhaled. DO NOT inhale or ingest.

May irritate eyes. DO NOT get into eyes.

**KEEP OUT OF REACH OF CHILDREN.**

Avoid contact with eyes, skin or clothing.

Avoid breathing spray mist.

Wash thoroughly after using and before eating, drinking or smoking.

Mixing and loading is not permitted within 30 m from aquatic habitat.

#### Protective Clothing and Equipment:

During application, handling, mixing and loading, wear goggles or face shield, chemical resistant gloves, respirator, and coveralls over a long-sleeved shirt and long pants.

Wear chemical resistant gloves for cleanup and repair.

#### Re-entry Restriction:

Workers should not enter treated Corn fields within 24 hours of treatment. Workers who must enter fields within this time period should wear long-sleeved shirt, long pants and chemical resistant gloves.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at [www.croplife.ca](http://www.croplife.ca).

### **Section 3: FIRST AID AND TOXICOLOGICAL INFORMATION**

**FIRST AID:** Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**IF SWALLOWED:** Call a poison control centre or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

**GENERAL INFORMATION:** Seek medical attention or contact a poison control centre immediately if irritation persists or is severe.

#### **TOXICOLOGICAL INFORMATION:**

Medical personnel should contact Bayer CropScience's medical information services toll-free 1-800-334-7577.

**Note to Physician:** If this product is ingested, endotracheal intubation and gastric lavage should be performed as soon as possible, followed by charcoal and sodium sulfate administration.

### **Section 4: ENVIRONMENTAL HAZARDS:**

- Toxic to non-target terrestrial plants and aquatic organisms. Observe the buffer zones and precautionary measures specified under Directions for Use (see Section 9 Application Instructions and Cautions).
- Atrazine is persistent and will carry over. It is recommended that any products containing atrazine not be used in areas treated with this product during the previous season.
- Use good pesticide practices and apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and parks is minimal. Take into consideration wind speed, wind direction, temperature, application equipment and sprayer settings used for applications.
- For the protection of non-target habitats, over-spray or drift to any body of water or other environmentally sensitive habitats must be avoided. The interaction of many

equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

- Avoid over-spray or drift to sensitive / non-target aquatic habitats. An appropriate buffer zone is required between the downwind point of direct application and the closest edge of sensitive aquatic habitats including sloughs, coulees, ponds, prairie potholes, lakes, rivers, streams, reservoirs, and wetlands that are situated on the periphery of the treated area. Do not apply during periods of dead calm or when winds are gusty. Do not apply this product through any type of irrigation system. **USE ONLY ON LABELLED CROPS.**
- Avoid spray drift to susceptible plants and **USE ONLY FOR RECOMMENDED PURPOSES AND AT RECOMMENDED RATES.**
- **DO NOT APPLY BY AIR**

## **Section 5: STORAGE**

- CANNOT be stored below freezing.
- If stored for 1 year or longer, shake well before using.
- Store the tightly closed container away from feeds, seeds, fertilizers, plants and foodstuffs.
- Do not use or store in or around the home.
- Keep in original container during storage.

## **Section 6: DISPOSAL**

### **Recyclable Container Disposal:**

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/ dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

### **Returnable Container Disposal:**

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/ dealer).



**Refillable Container Disposal:**

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

**Disposal of Unused, Unwanted Product:**

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for cleanup of spills.

**Disposal of Unused Spray Solution:**

If any spray solution remains in the tank after spraying is finished, it should be thoroughly diluted, then sprayed on the perimeter of the area just sprayed, away from water supplies, ditches, and irrigation canals.

## DIRECTIONS FOR USE

**Section 7: CROPS, WEEDS, RATES, TIMING**

**Crop:** Glufosinate ammonium tolerant corn, i.e. Liberty Link hybrids.

**Weeds:** Crabgrass, Fall Panicum, Giant Foxtail, Green Foxtail, Old Witchgrass, Proso Millet, Yellow Foxtail, Wild Oats, Chickweed, Cocklebur, Eastern Black Nightshade, Lady's-thumb, Lamb's-quarters, Purslane, Ragweed, Redroot Pigweed, Shepherd's-purse, Wild Buckwheat, Wild Mustard, Wormseed Mustard, Velvetleaf.

**Recommended Time (Weed Leaf Stage):** LIBERTY AT Herbicide should be applied when weeds are in the 1 - 4 leaf stage.

**Rate:** Apply LIBERTY AT Herbicide at a rate of 2.9 L/ha

**ADDITIVES:** For enhanced control of velvetleaf under cool and/or dry conditions, add ammonium sulphate to the tank at a rate of 6 L/ha (49% solution) OR 3 kg/ha (99%).

**Note:** Apply LIBERTY AT Herbicide from the 1-8 leaf stage of the Corn plant or 5-6 visible collars (the leaf is counted once the next leaf is visible in the whorl) and before the corn reaches 30 cm in height.

**Note:** Do not tank mix LIBERTY AT Herbicide with pesticides, fertilizers or any other chemical additives unless recommended on this label.

## **Section 8: HERBICIDE RESISTANCE MANAGEMENT**

For resistance management, LIBERTY AT Herbicide is a Group 5 and 10 herbicide. Any weed population may contain or develop plants naturally resistant to LIBERTY AT Herbicide and other Group 5 and 10 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of LIBERTY AT Herbicide or other Group 5 and 10 herbicides with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted.
- Herbicide use should be based on an IPM program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical), cultural, biological and other chemical control practices.
- Monitor treated weed populations for resistance development.
- Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment and planting clean seed.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Bayer CropScience Inc. at 1-888-283-6847.

## **Section 9: APPLICATION INSTRUCTIONS AND CAUTIONS**

### **Application Instructions:**

- Protect sprayer operators from drift or mist.
- When low volumes of spray are applied, complete coverage and thorough application are essential for the most effective results. Schedule applications in accordance with local conditions. Consult your local agricultural authorities for specific use information.
- For best results, apply to emerged, young, actively growing weeds.
- LIBERTY AT Herbicide will have an effect on weeds that are larger than the recommended leaf stage, however speed of activity and control may be reduced.
- Apply LIBERTY AT Herbicide in a minimum of 110 L/ha of water, at a pressure of 175-275 kPa and at a ground speed of 6-10 kph. The use of 80° or 110° flat fan nozzles is highly recommended for optimum spray coverage and

canopy penetration.

- Application of the spray at a 45° angle forward will result in better spray coverage. Follow directions elsewhere on the label for the correct rate and timing of application.
- The speed of action of LIBERTY AT Herbicide is influenced by environmental factors. At cool temperatures (below 10°C), poor moisture and low humidity, speed of action may be reduced. Generally, visual symptoms appear 2-4 days after application.
- IF RAINFALL OCCURS WITHIN 4 HOURS OF APPLICATION, EFFECTIVENESS MAY BE REDUCED.
- Weed control may be reduced when heavy dew, fog or mist are present at the time of application.
- LIBERTY AT Herbicide works primarily as a contact herbicide. Thorough coverage of the weeds to be controlled is essential.

#### **Application Cautions:**

- **GROUND APPLICATION ONLY. DO NOT APPLY BY AIR**
- **DO NOT MAKE MORE THAN ONE APPLICATION PER YEAR.**
- Do not use flood jet nozzles, controlled droplet application equipment or air-assisted spray equipment.
- Uniform, thorough spray coverage is important to achieve consistent weed control.
- **DO NOT** apply if winds exceed 16 kph when using open boom sprayers.
- **DO NOT** apply if winds exceed 25 kph when using hooded sprayers.
- **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) medium classification. Boom height must be 60 cm or less above the crop or ground.
- **DO NOT** overspray non-target terrestrial or aquatic habitats.
- **DO NOT** contaminate aquatic habitats when cleaning and rinsing spray equipment or containers.
- Avoid contact with desirable plants or crops either from direct application or from spray drift as severe damage may occur.
- Application of LIBERTY AT Herbicide to Corn hybrids which are not glufosinate ammonium tolerant or other crops will result in severe crop injury or death of the crop.
- **SUSCEPTIBLE WEEDS:** LIBERTY AT Herbicide has an effect on all weeds and crops except for those crops which are developed to be tolerant to applications of LIBERTY AT Herbicide.

#### **Buffer Zones:**

- The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial

habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, pastures, rangelands and shrublands) or sensitive aquatic habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs, wetlands and estuarine/marine habitats).

Application Method	Buffer Zone for Protection of:	
	Terrestrial Habitat (m)	Aquatic Habitat (m)
Field sprayer*	10	10

\*For field sprayer application, buffer zones can be reduced with the use of drift reducing spray shields. When using a spray boom fitted with a full shield (shroud, curtain) that extends to the canopy or ground, the labelled buffer zone can be reduced by 70%. When using a spray boom where individual nozzles are fitted with cone-shaped shields that are no more than 30 cm above the canopy or ground, the labelled buffer zone can be reduced by 30%.

**Surface Runoff:**

- To reduce runoff from treated areas into aquatic habitats, consider the characteristics and conditions of the site before treatment. Site characteristics and conditions that may lead to runoff include, but are not limited to, heavy rainfall, moderate to steep slope, bare soil and poorly draining soil (e.g., soils that are compacted, fine textured or low in organic matter such as clay).
- Avoid applying this product when heavy rain is forecast.
- Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip (buffer zone) between the treated area and the edge of the water body.

**Leaching:**

- The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g., sand, loamy sand and sandy loam soils) and/or the depth to the water table is shallow.
- Avoid applying this product when heavy rain is forecast.

**Recropping Restrictons:**

- Due to the atrazine component in LIBERTY AT Herbicide, special precautions must be taken with respect to recropping.
- Crop injury may occur if land treated with Liberty AT is planted with any crop other than field corn, including corn hybrids which are not glufosinate ammonium tolerant in the same season. One year after application there is virtually no hazard to most rotational crops when label rates are applied. Sugar beets and tobacco are sensitive to atrazine and should not be planted on land treated with Liberty AT Herbicide the previous year. When extended

periods of dry weather occur during the year of treatment, there can be some injury to succeeding crops such as white beans, onions, peas, tomatoes and turnips. Injury is most likely to occur when the seedling crop is subjected to periods of stress e.g. during periods of abnormally hot, dry weather. Under conditions of heavy rainfall sufficient to cause soil erosion, soil containing atrazine may wash to lower areas of land and injury to existing or subsequent crops other than corn may result. Thorough tillage including moldboard ploughing should precede the planting of crops other than corn. Plough the land flat as opposed to putting the furrows on edge.

- UNEVEN APPLICATION (e.g. spray overlaps) OF LIBERTY AT HERBICIDE WILL NOT INJURE CORN BUT MAY RESULT IN THE INJURY OF SUCCEEDING CROPS OTHER THAN CORN.

## **Section 10: MIXING INSTRUCTIONS**

LIBERTY AT Herbicide must be applied with properly calibrated, clean equipment. LIBERTY AT is specially formulated to mix readily in water.

Prior to adding LIBERTY AT Herbicide to the spray tank, ensure that the spray tank is thoroughly clean (see Section 12 "SPRAYER CLEANUP").

1. Fill tank to three-quarters full with clean water prior to adding LIBERTY AT Herbicide.
2. Thoroughly shake the LIBERTY AT Herbicide container.
3. Add the correct amount of LIBERTY AT Herbicide.
4. Add the remaining amount of water, begin agitation, and spray out immediately.

**Note:** When using ammonium sulphate, ALWAYS add it to the tank first, prior to the addition of LIBERTY AT Herbicide.

**Note:** The addition of an anti-foaming agent may reduce foaming, especially when using soft water.

**Note:** Ensure that all circuits (pipes, booms, etc.) have the correct LIBERTY AT Herbicide/water concentration before application is started.

## **Section 11: SPRAYER CLEANUP**

Before and after using LIBERTY AT Herbicide always complete a thorough cleaning of the spray tank, lines and filter. Spray equipment should be thoroughly rinsed using a strong detergent solution.

## **Section 12: LIVESTOCK FEEDING AND PREHARVEST INTERVALS**

- DO NOT graze or cut for fodder the treated corn before ear emergence has occurred. There are not sufficient data available to support such use.
- DO NOT apply within 86 days of harvesting Corn grain.
- Do NOT harvest within 45 days of application on sweet corn and 60 days for field corn.

## **Section 13: NOTICES**

### **NOTICE TO USER:**

This pest control product is to be used only in accordance with the directions on label. It is an offence under the *Pest Control Products Act* to use a control product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

### **FOR MORE INFORMATION CONTACT:**

**Product Information:** 1-888-283-6847

**Bayer CropScience Inc.  
Suite 100, 3131 114<sup>th</sup> Ave. SE  
Calgary, Alberta T2Z 3X2**

LIBERTY and LibertyLink are trademarks of Bayer.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.