

2023-5931
2024-02-16



container label

GROUP	4A	INSECTICIDE
-------	----	-------------

GAUCHO® 600 FL INSECTICIDE

GAUCHO 600 FL is a systemic insecticide seed treatment for early season protection of canola, mustard (condiment-type only) and rapeseed seed and seedlings against flea beetles. GAUCHO 600 FL Insecticide also provides early season control of wireworms in field corn.

COMMERCIAL

ACTIVE INGREDIENT: Imidacloprid.....600 g/L

Contains 1,2-benzisothiazolin-3-one at 0.035% and 5-chloro-2-methyl-4-isothiazolin-3-one at 0.0009% and 2-methyl-4-isothiazolin-3-one at 0.0003% as preservatives.

REGISTRATION NO. 27170 PEST CONTROL PRODUCTS ACT



WARNING

POISON

READ THE LABEL AND BOOKLET BEFORE USING

Special Use Restrictions: This product contains no colourant. All seed treated with this product must be conspicuously coloured at the time of treatment. An appropriate colourant must be added when this product is applied. Regulations pertaining to the "Seeds Act" must be strictly adhered to when using this product. A blue colourant must be added when this product is applied to oilseed, and an appropriate colour must be added when this product is applied to corn.

Bayer CropScience Inc.
200-160 Quarry Park Blvd. S.E.
Calgary, Alberta T2C 3G3

In case of spills, poisoning or fire, telephone emergency response number
1-800-334-7577 (24 hours a day).
Product Information: 1-888-283-6847

PROTECT FROM FREEZING
NET CONTENTS: 0.25-1000 L
GAUCHO ® is a Registered Trademark of Bayer.

ENVIRONMENTAL PRECAUTIONS:

Toxic to birds, small wild mammals and aquatic organisms. Keep out of lakes, streams, ponds or other aquatic systems. Do not contaminate water when disposing of equipment wash waters.

Apply this product only in accordance with this label.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Toxic to certain beneficial arthropods (which may include predatory and parasitic insects, spiders, and mites). Imidacloprid is toxic to bees. Bees can be exposed to product residues in flowers, leaves, pollen and/or nectar resulting from seed treatment applications. When used according to label directions minimal exposure or risk is expected. To help minimize the dust generated during planting, refer to the complete guidance “Pollinator Protection and Responsible Use of Treated Seed- Best Management Practices” on the Health Canada webpage on pollinator protection at www.canada.ca/pollinators . When using a seed flow lubricant with corn seed treated with GAUCHO 600 FL Insecticide, only a dust-reducing fluency agent is permitted. Talc and graphite are not permitted to be used as a seed flow lubricant for corn seed treated with this insecticide. Carefully follow use directions for the seed flow lubricant. Do not load or clean planting equipment near bee colonies, and avoid places where bees may be foraging, such as flowering crops or weeds. When turning on the planter, avoid engaging the system where emitted dust may contact honey bee colonies.

Spilled or exposed seeds and dust must be incorporated into the soil or cleaned up from the soil surface.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN.

Harmful or fatal if swallowed and harmful if inhaled. Avoid breathing vapour or spray mist. For use in commercial seed treatment facilities (and by mobile treaters) with closed-transfer, including closed mixing, loading, calibrating, and closed-treatment equipment only. No open transfer is permitted. During mixing, loading, application, bagging, sewing and stacking, wear coveralls over long-sleeved shirt, long pants, chemical-resistant gloves, socks and chemical-resistant footwear. When cleaning seed treatment equipment, wear chemical-resistant coveralls over long-sleeved shirt, long pants, chemical-resistant gloves, socks, chemical-resistant footwear and a respirator with a NIOSH-approved organic-vapour-removing cartridge with a prefilter approved for pesticides OR a NIOSH-approved canister approved for pesticides. When handling and planting treated seeds, wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and chemical-resistant footwear. Protective eye wear should be worn when handling this product. Use good personal hygiene, washing hands and exposed skin before eating, drinking or smoking. No food, drink or tobacco should be allowed in areas of chemical storage or use.

If this pest control product is to be used on a commodity that may be exported and you require information regarding Maximum Residue Limits for an importing country, please contact Bayer CropScience Canada Inc. at 1-888-283-6847 or www.cropscience.bayer.ca.

FIRST AID:

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label, or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

NOTE TO PHYSICIANS: Treat symptomatically. There is no specific antidote.

STORAGE AND DISPOSAL:

STORAGE: Store product in original container only, away from other pesticides, fertilizer, food or feed. Keep container closed. Store in a cool place. Do not store GAUCHO 600 FL Insecticide in direct sunlight. Protect from freezing temperatures.

DISPOSAL (FOR NON-RETURNABLE CONTAINERS):

1. Triple- or pressure-rinse the empty container. Add the rinsings to the mixture in the tank.
2. Follow provincial instruction for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of spill and for clean-up of spills

DISPOSAL (FOR RECYCLABLE CONTAINERS):

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.



DISPOSAL (FOR RETURNABLE CONTAINERS):

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill and for clean-up of spills.

TRIPLE RINSING INSTRUCTIONS FOR CONTAINER DISPOSAL (FOR NONRETURNABLE AND RECYCLABLE CONTAINERS ONLY):

This container should be triple rinsed prior to its disposal. The rinsate can be added to the seed treatment provided the volume of water does not exceed 6 % of the container size.

Add 1/3 of the rinse water to the container and mix the contents thoroughly. Empty the rinsate into the seed treater holding tank and repeat this process two more times making sure the total volume of rinsate does not exceed 6 % of the container size. Make sure rinsate is thoroughly mixed with seed treatment before treating. Close the container.

CAUTION: Do not dilute beyond 6 % or excess seed wetness may result in seed handling difficulties. Using excess water can also cause GAUCHO 600 FL Insecticide to thin resulting in settling of solids in the product.

SEED QUALITY: Treatment of highly mechanically damaged seed, or seed of known low vigour and poor quality may result in reduced germination and/or reduction of seed and seedling vigour. If seed lot quality is unknown, conduct a germination test on a small portion of seed before committing the total seed lot to a selected chemical treatment.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Product Act* to use this product in a way that is inconsistent with the directions on the label.

booklet label

GROUP	4A	INSECTICIDE
-------	----	-------------

GAUCHO® 600 FL INSECTICIDE

GAUCHO 600 FL is a systemic insecticide seed treatment for early season protection of canola, mustard (condiment-type only) and rapeseed seed and seedlings against flea beetles. GAUCHO 600 FL

Insecticide also provides early season control of wireworms in field corn.

COMMERCIAL

ACTIVE INGREDIENT: Imidacloprid.....600 g/L

Contains 1,2-benzisothiazolin-3-one at 0.035% and 5-chloro-2-methyl-4-isothiazolin-3-one at 0.0009% and 2-methyl-4-isothiazolin-3-one at 0.0003% as preservatives.

REGISTRATION NO. 27170 PEST CONTROL PRODUCTS ACT



WARNING

POISON

READ THE LABEL AND BOOKLET BEFORE USING

Special Use Restrictions: This product contains no colourant. All seed treated with this product must be conspicuously coloured at the time of treatment. An appropriate colourant must be added when this product is applied. Regulations pertaining to the "Seeds Act" must be strictly adhered to when using this product. A blue colourant must be added when this product is applied to oilseed, and an appropriate colour must be added when this product is applied to corn.

Bayer CropScience Inc.
200-160 Quarry Park Blvd. S.E.
Calgary, Alberta T2C 3G3

In case of spills, poisoning or fire, telephone emergency response number
1-800-334-7577 (24 hours a day).
Product Information: 1-888-283-6847

PROTECT FROM FREEZING
NET CONTENTS: 0.25-1000 L

GAUCHO ® is a Registered Trademark of Bayer.

GENERAL INFORMATION:

GAUCHO 600 FL is a systemic insecticide seed treatment for early season protection of canola, mustard (condiment-type only) and rapeseed seed and seedlings from flea beetles. In areas where flea beetle populations are often high, the higher application rate is recommended.

GAUCHO 600 FL Insecticide also provides early season control of wireworms in field corn.

DIRECTIONS FOR USE:

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests. DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Dispose of all excess treated seed. Leftover treated seed may be buried away from water sources in accordance with local requirements. Do not dispose of seed by double-sowing in headlands. Broadcast seeding is prohibited.

For use in commercial seed treaters only. Not for use on agricultural establishments in hopper-box, slurry-box, or other seed treatment applications at, or immediately before, planting. This product is to be used in liquid or slurry treaters. Mix thoroughly before use, or use entire container at one time.

Equipment used in seed treating plants can vary thus it is advised that a Bayer Technical Representative be contacted to discuss the use of GAUCHO 600 FL Insecticide in the intended seed treater prior to use. GAUCHO 600 FL Insecticide treated seed may reduce seed flow in the seed drill. Recalibration of the seed drill may be required to obtain correct seeding rate before planting. GAUCHO 600 FL Insecticide can be used as an over-treatment.

CANOLA, MUSTARD (CONDIMENT-TYPE ONLY), AND RAPESEED

For early season protection of seed and seedlings against flea beetles.

Mixing Instructions per 100 kg seed.

GAUCHO 600 FL Application Rate	Component		Total Mixture
	GAUCHO 600 FL	Water*	
667 mL/100kg seed	670 mL	67 mL	737 mL
1333 mL/100kg seed	1330 mL	133 mL	1463 mL

*See triple rinsing instructions.

NOTE: A colourant MUST be added to this product to colour seed in accordance with the PCP Act and the Seeds Act Regulations. A blue colourant must be added when this product is applied to oilseed. A suitable seed colourant such as Magna-Coat Colourant must be applied to the seed during the pelleting or the coating process. Magna-Coat Polymer and Magna-Coat Colourant are also available from Bayer. For assistance please contact a Bayer representative.

CORN

For early season control of wireworms.

Crop	Pest	Application Rate	Remarks
Field Corn	Wireworms	21.3 mL product per 80,000 seeds (0.16 mg ai per Kernel)	Dilute in sufficient water to achieve uniform distribution on the seed. Ensure that seed is adequately coloured. Other polymers and coating materials may be required.
Field Corn for Seed Production	Wireworms	21.3 mL product per 80,000 seeds (0.16 mg ai per kernel)	Contact your Bayer representative for dilution instructions specific to the equipment in your treatment facility

Pre-test the germination of a small sample of each seed lot with GAUCHO 600 FL Insecticide prior to commercial application to the whole lot. Carry-over of GAUCHO 600 FL Insecticide treated seed is not recommended. GAUCHO 600 FL Insecticide can be used as an overtreatment. NOTE: A colourant MUST be added to this product to colour seed in accordance with the PCP Act and the Seeds Act Regulations. An appropriate colour for corn must be added when this product is applied to corn seed. A suitable seed colourant such as PRO-IZED® Liquid Red Seed Colourant must be applied to the seed during the pelleting or the coating process. Polymers and coating materials are available from Bayer. For assistance please contact a Bayer representative.

METHOD OF APPLICATION:

Storage of GAUCHO 600 FL Insecticide at low temperatures is not recommended. Prior to and during application, GAUCHO 600 FL Insecticide must be THOROUGHLY AGITATED to ensure uniform mixing of the product. Due to the viscosity of the material, it should be kept above 10o C prior to and during application. Do not apply direct heat to container.

TANK MIXES:

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Bayer CropScience Inc. at 1-888-283-6847 or www.cropscience.bayer.ca for information before applying any tank mix that is not specifically recommended on this label.

Bayer CropScience Inc. has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) of all potential tank mixes under all environmental conditions or for all crop varieties. Conduct a germination test on a small portion of seed before committing the total seed lot to tank mixes that are not specifically listed on this label.

RESISTANCE MANAGEMENT RECOMMENDATIONS:

For resistance management, please note that GAUCHO 600 FL Insecticide contains a Group 4A insecticide. Any insect population may contain individuals naturally resistant to GAUCHO 600 FL Insecticide and other Group 4 insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay insecticide resistance:

- Where possible, rotate the use of GAUCHO 600 FL Insecticide or other Group 4 insecticides with different groups that control the same pests in a field.
- Use tank mixtures with insecticides from a different group that is effective on the target pest when such use is permitted.
- Insecticide use should be based on an IPM program that includes scouting, record keeping, and considers cultural, biological and other chemical control practices.
- Monitor treated pest populations for resistance development.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact Bayer CropScience via internet at www.cropscience.bayer.ca or phone at 1-888-283-6847.

USE RESTRICTIONS:

Do not use treated seed for food, feed or oil processing.

Do not graze or feed livestock on treated areas for four weeks after planting.

Mustard greens grown or harvested from GAUCHO 600 FL Insecticide treated seed must not be used for human consumption.

Treated canola, rapeseed or mustard (condiment-type only) seed stored for periods in excess of 6 months may decrease in germination at a faster rate than untreated seed. Treated seed stored for more than 6 months should be tested for germination before planting. Do not store treated seed above 25°C or in direct sunlight.

LABELLING TREATED SEED:

All bags containing treated seed must be labelled or tagged as follows:

- This seed has been treated with GAUCHO 600 FL Insecticide seed protectant, which contains imidacloprid.
- Do not use for feed, food or oil processing. Store away from feeds and other foodstuffs.
- Toxic to birds, small wild mammals and aquatic organisms. Any spilled or exposed seeds must be incorporated into the soil or otherwise cleaned-up from the soil surface.
- Dispose of all excess treated seed. Leftover treated seed may be buried away from water sources in accordance with local requirements. Do not dispose of seed by double-sowing in headlands.
- Broadcast seeding is prohibited.
- Closed cabs must be used for planting commercially treated and bagged seeds.
- Do not plant treated seeds by hand.

Additionally, all treated corn seed for sale or use in Canada must be labelled with the following information:

- Imidacloprid is toxic to bees. Dust generated during planting of treated seed may be harmful to bees and other pollinators.
- To help minimize the dust generated during planting, refer to the complete guidance “Pollinator Protection and Responsible Use of Treated Seed- Best Management Practices” on the Health Canada webpage on pollinator protection at www.canada.ca/pollinators .
- When using a seed flow lubricant with this treated seed, only a dust-reducing fluency agent is permitted. Talc and graphite are not permitted to be used as a seed flow lubricant for corn seed treated with this insecticide. Carefully follow use directions for the seed flow lubricant.
- Do not load or clean planting equipment near bee colonies, and avoid places where bees may be foraging, such as flowering crops or weeds.
- When turning on the planter, avoid engaging the system where emitted dust may contact honey bee colonies.
- Spilled or exposed seeds and dust must be incorporated into the soil or cleaned up from the soil surface.

ENVIRONMENTAL PRECAUTIONS:

Toxic to birds, small wild mammals and aquatic organisms. Keep out of lakes, streams, ponds or other aquatic systems. Do not contaminate water when disposing of equipment wash waters. Apply this product only in accordance with this label.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Toxic to certain beneficial arthropods (which may include predatory and parasitic insects, spiders, and mites). Imidacloprid is toxic to bees. Bees can be exposed to product residues in flowers, leaves, pollen and/or nectar resulting from seed treatment applications. When used according to label directions minimal exposure or risk is expected. Dust generated during planting of treated seed may be harmful to bees and other pollinators. To help minimize the dust generated during planting, refer to the complete guidance “Pollinator Protection and Responsible Use of Treated Seed- Best Management Practices” on the Health Canada webpage on pollinator protection at www.canada.ca/pollinators . When using a seed flow lubricant with corn seed treated with GAUCHO 600 FL Insecticide, only a dust-reducing fluency agent is permitted. Talc and graphite are not permitted to be used as a seed flow lubricant for corn seed treated with this insecticide. Carefully follow use directions for the seed flow lubricant. Do not load or clean planting equipment near bee colonies, and avoid places where bees may be foraging, such as flowering crops or weeds. When turning on the planter, avoid engaging the system where emitted dust may contact honey bee colonies.

Spilled or exposed seeds and dust must be incorporated into the soil or cleaned up from the soil surface.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN.

Harmful or fatal if swallowed and harmful if inhaled. Avoid breathing vapour or spray mist. For use in commercial seed treatment facilities (and by mobile treaters) with closed-transfer, including closed mixing, loading, calibrating, and closed-treatment equipment only. No open transfer is permitted. During mixing, loading, application, bagging, sewing and stacking, wear coveralls over long-sleeved shirt, long pants, chemical-resistant gloves, socks and chemical-resistant footwear. When cleaning seed treatment equipment, wear chemical-resistant coveralls over long-sleeved shirt, long pants, chemical-resistant gloves, socks, chemical-resistant footwear and a respirator with a NIOSH-approved organic-vapour-removing cartridge with a prefilter approved for pesticides OR a NIOSH-approved canister approved for pesticides. When handling and planting treated seeds, wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and chemical-resistant footwear. Protective eye wear should be worn when handling this product.

Use good personal hygiene, washing hands and exposed skin before eating, drinking or smoking. No food, drink or tobacco should be allowed in areas of chemical storage or use.

If this pest control product is to be used on a commodity that may be exported and you require information regarding Maximum Residue Limits for an importing country, please contact Bayer CropScience Canada Inc. at 1-888-283-6847 or www.cropscience.bayer.ca.

FIRST AID:

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice.

Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label, or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

NOTE TO PHYSICIANS: Treat symptomatically. There is no specific antidote.

STORAGE AND DISPOSAL:

STORAGE: Store product in original container only, away from other pesticides, fertilizer, food or feed. Keep container closed. Store in a cool place. Do not store GAUCHO 600 FL Insecticide in direct sunlight. Protect from freezing temperatures.

DISPOSAL (FOR NON-RETURNABLE CONTAINERS):

1. Triple- or pressure-rinse the empty container. Add the rinsings to the mixture in the tank.
2. Follow provincial instruction for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of spill and for clean-up of spills

DISPOSAL (FOR RECYCLABLE CONTAINERS):

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

DISPOSAL (FOR RETURNABLE CONTAINERS):

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill and for clean-up of spills.

TRIPLE RINSING INSTRUCTIONS FOR CONTAINER DISPOSAL (FOR NONRETURNABLE AND RECYCLABLE CONTAINERS ONLY):

This container should be triple rinsed prior to its disposal. The rinsate can be added to the seed treatment provided the volume of water does not exceed 6 % of the container size.

Add 1/3 of the rinse water to the container and mix the contents thoroughly. Empty the rinsate into the seed treater holding tank and repeat this process two more times making sure the total volume of rinsate does not exceed 6 % of the container size. Make sure rinsate is thoroughly mixed with seed treatment before treating. Close the container.

CAUTION: Do not dilute beyond 6 % or excess seed wetness may result in seed handling difficulties. Using excess water can also cause GAUCHO 600 FL Insecticide to thin resulting in settling of solids in the product.

SEED QUALITY: Treatment of highly mechanically damaged seed, or seed of known low vigour and poor quality may result in reduced germination and/or reduction of seed and seedling

vigour. If seed lot quality is unknown, conduct a germination test on a small portion of seed before committing the total seed lot to a selected chemical treatment.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Product Act* to use this product in a way that is inconsistent with the directions on the label.