

2024-2968
2024-07-18

Brenntag Canada Inc.

Sprout Nip 980 Potato Sprout Inhibitor

Sprout Inhibitor for Potatoes in Storage

For Use Only by Brenntag Canada Inc. Trained Operators

COMMERCIAL

Solid

Fog Application

GUARANTEE: chlorpropham 98%

REGISTRATION NUMBER 27213 PEST CONTROL PRODUCTS ACT



CAUTION POISON

READ THE LABEL AND CARTON BEFORE USING

NET CONTENTS: 200 kg

4.5 Kg x 4 (18 kg)

5.0 Kg x 4 (20 Kg)

4.54 Kg x 5 (22.68 Kg)

4.50 Kg x 5 (22 Kg)

Brenntag Canada Inc.
5025 Creekbank Road
Building 'A', Suite 201
Mississauga, Ontario
L4W 0B6 Canada
416 259 8231

KEEP OUT OF THE REACH OF CHILDREN

PRECAUTIONS

Warning - May be irritating to eyes, skin and respiratory tract. May be harmful if inhaled. Use only in well ventilated areas. Avoid breathing spray mist or vapour. Avoid contact with skin, eyes and clothing.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's website at www.CropLife.ca.

PROTECTIVE CLOTHING AND EQUIPMENT:

PERSONAL PROTECTIVE EQUIPMENT

For mixing and loading wear cotton coveralls and chemical-resistant gloves.

Eye Protection: Use full face shield or chemical safety goggles when handling.

Skin Protection: Wear cotton coveralls, chemical-resistant gloves, shoes and socks when handling.

Respiratory Tract Protection: Wear a respirator with a NIOSH/MSHA/BHSA approved organic-vapour-removing cartridge with a prefilter approved for pesticides when handling or if there is potential for inhalation of the vapour or spray mist.

For all re-entry tasks (including cleaning) prior to the completion of ventilation, wear chemical-resistant coveralls over a long-sleeved shirt and pants, chemical resistant gloves and self-contained breathing apparatus. For cleaning following ventilation, wear chemical-resistant coveralls over a long-sleeved shirt and pants, chemical-resistant gloves and a suitable respirator.

For tasks other than cleaning following ventilation, wear cotton coveralls and chemical-resistant gloves.

RESTRICTED-ENTRY INTERVAL (REI) : Do not allow entry into treated areas (to attend to fans, etc.) until after two (2) hours of mechanical ventilation (fans should circulate outside air) or four (4) hours of passive ventilation (windows, vents and doors should be opened). If visible aerosol remains after these periods, do not enter until the aerosol has settled.

FIRST AID

If on skin or clothing take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If in eyes hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If swallowed call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice.

Take container, label or product name and Pest Control Product registration number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION Note to physician: Treat symptomatically. Overexposure to chlorpropham may result in methemoglobinemia.

DIRECTIONS FOR USE

Seal the potato storage building to prevent or reduce leakage. Adjust the ventilation system to ensure even distribution of moving air. Adjust the application equipment to deliver droplets in the range of 1 to 10 microns. Ensure that air blower screens are clear and no restrictions in air flow are present on the thermal generator. Apply sufficient SPROUT NIP 980 POTATO SPROUT INHIBITOR (1.67 - 3.33 kg per 100 metric tonnes of potatoes) to achieve a deposit on the potatoes in the range of 6 to 12 ppm (whole weight basis). The rate applied will depend on storage conditions, e.g., temperature, free moisture and length of storage period. A second application may be required in 90-150 days depending on storage conditions. Following treatment with Sprout Nip 980 Potato Sprout Inhibitor plant growth regulator continue air circulation until fog clears (2-4 hours). Standard aeration/ventilation for temperature and humidity control can then be resumed.

Use Precautions

1) SPROUT NIP 980 POTATO SPROUT INHIBITOR plant growth regulator is applied as a mist or fog to inhibit sprouting of potatoes in storage. It should only be used with an approved thermal generator for safe distribution. The concentration of SPROUT NIP 980 POTATO SPROUT INHIBITOR plant growth regulator must be kept below 11.5 kilograms per 283 m³ (10,000 standard cubic feet) of air per hour. Explosion may occur if the lower limit for combustion is exceeded.

2) Potatoes to be treated must be firm and free from excessive dirt or other debris. Mechanically damaged potatoes cannot be treated until suberization is complete, a process which normally takes 2-3 weeks or more.

3) Prolonged application with thermal generators may cause fires. Remove all flammable material from the vicinity of the application point before operations commence. Keep at least two operational fire extinguishers suitable for use in confined areas at the point of application. Do not leave equipment unattended during operation.

4) The fog from this product is irritating to lungs and skin.

5) This product will inhibit sprouting of potatoes used for seed. Do not apply on or near potatoes to be used for this purpose. Do not store potatoes to be used for seed in treated buildings.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's website at www.CropLife.ca.

CONTAINER STORAGE AND DISPOSAL

Store away from food, feed and seed potatoes. Store in a cool dry environment. Do not contaminate water by cleaning of equipment or disposal of wastes.

1. Empty contents of container thoroughly into the thermal generator.
2. Follow provincial instruction for any required additional cleaning on the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *PEST CONTROL PRODUCTS ACT* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.