

2014-1102  
2014-04-02

[FRONT PANEL - ENGLISH]

**ORTHO® BUG-B-GON® MAX™ ANT & CHINCH BUG ELIMINATOR  
READY-TO-SPRAY**

DOMESTIC  
INSECTICIDE

For control of insect pests on fruits,  
vegetables, ornamentals and lawns

NET CONTENTS: 1 L

GUARANTEE:

Carbaryl ..... 22.5% by wt.

Contains 1,2-benzisothiazolin-3-one at 0.019% as a preservative.

REG. NO.: 27207 P.C.P. ACT

READ THE LABEL BEFORE USING  
KEEP OUT OF REACH OF CHILDREN  
SHAKE BEFORE USING

Scotts Canada Ltd.  
2000 Argentia Road  
Plaza #2, Suite 300  
Mississauga, ON L5N 1V8

CAUTION



POISON

This product contains the SEVIN® brand of carbaryl.  
SEVIN® is a registered trademark of The Aventis Group.

[BACK PANEL - ENGLISH]

OPEN  
Resealable Label  
for Additional Product  
Information

**ORTHO® BUG-B-GON® MAX™ ANT & CHINCH BUG ELIMINATOR  
READY-TO-SPRAY**

**PRODUCT FACTS**

<b>CONTROLS INSECTS</b> (as listed on the label)	<ul style="list-style-type: none"><li>• Aphids</li><li>• Birch leafminer</li><li>• Colorado potato beetle</li><li>• Cutworms</li><li>• European chafer</li><li>• Japanese beetle</li><li>• Leafhoppers</li></ul>	<ul style="list-style-type: none"><li>• Leafrollers</li><li>• Mealybugs</li><li>• Tarnished plant bugs</li><li>• Tent caterpillars</li><li>• Scale insects</li></ul>
<b>WHERE TO USE</b>	<ul style="list-style-type: none"><li>• Vegetables</li><li>• Fruit Trees</li><li>• Small fruit crops</li></ul>	<ul style="list-style-type: none"><li>• Ornamentals</li><li>• Trees</li><li>• Turf/Lawns</li><li>• Recreational areas</li></ul>

[IMAGE OF TELEPHONE]

[ICON OF QUESTION MARK]

**Questions, Comments or Medical Information**  
Call 1-800-668-5669 or visit our Web Site at [www.scottscanada.ca](http://www.scottscanada.ca)

**APPLICATOR DIRECTIONS:**

[ILLUSTRATION - ATTACH HOSE]	[ILLUSTRATION - OPEN SYSTEM ]	[ILLUSTRATION - TURN ON AND SPRAY]
<ul style="list-style-type: none"> <li>• Ensure that the large Water Control Knob is turned to the "OFF" position.</li> <li>• Connect sprayer to garden hose.</li> </ul>	<ul style="list-style-type: none"> <li>• Turn the small Product Control Button so that the flat side aligns with the tab on the right side of the button, and push the button all the way in.</li> <li>• Turn on water at faucet.</li> </ul>	<ul style="list-style-type: none"> <li>• To spray, point nozzle where you want to spray and turn the large knob all the way to the left.</li> <li>• To stop spraying, push small Product Control Button in from opposite side. Turn large Water Control Knob to the right to the "OFF" position and turn off faucet.</li> <li>• NOTE: BEFORE DISCONNECTING HOSE (to relieve water pressure): Turn large Water Control Knob to the left; then turn "OFF" again.</li> <li>• Disconnect garden hose from sprayer.</li> <li>• To prevent accidental discharge when not in use, turn the small Product Control Button so that it is <u>NOT</u> aligned with the tab and cannot be pushed in.</li> </ul>

**DIRECTIONS FOR USE:**

TO ASSURE A UNIFORM PRODUCT, ALWAYS SHAKE BEFORE USING.

Be sure the plants, areas of use and pests you wish to treat are listed on the label before you spray. Be sure to follow important limitations for the specific edible plant you are spraying, including the amount of spray which can be applied, the interval between sprays, and the days to wait before harvest. Do not exceed maximum application rate recommended for area of coverage.

Apply when insects and/or their damage appear. To maintain control, repeat at 7 to 10 day intervals or as necessary. When a dosage range is indicated, use the lower rate on young plants and immature insects, and higher rate on mature plants and insects. Thorough and uniform spray coverage is essential for effective control. Avoid application just before rainfall.

Do not treat fruits or vegetables grown in greenhouses. Do not use on humans or pets. Do not use inside the home.

**VEGETABLES**

<b>CROP - Asparagus seedlings, spears</b>		
<b>INSECT</b>	<b>PRE-HARVEST INTERVAL (DAYS)</b>	<b>CONTAINER COVERS</b>
Asparagus beetle Climbing cutworms	1	800 - 1,923 sq. metres (8,611 - 20,699 sq. feet)
<b>Timing/Application:</b> Apply when insects and/or their damage appear. Repeat at intervals of 7 to 10 days or as necessary. Use the lower rate on young plants. Do not treat more than once every 3 days.		
<b>CROP - Beans</b>		
<b>INSECT</b>	<b>PRE-HARVEST INTERVAL (DAYS)</b>	<b>CONTAINER COVERS</b>
Mexican bean beetle	3	3,333 - 4,000 sq. metres (35,876 - 43,056 sq. feet)
Flea beetles Leafhoppers		1,923 sq. metres (20,699 sq. feet)
Lygus bugs Stink bugs		800 - 962 sq. metres (8,611 - 10,355 sq. feet)
<b>Timing/Application:</b> Use lower rate on young plants. Apply for full coverage when insects and/or damage appear. Repeat at intervals of 7 to 10 days or as necessary.		
<b>CROP - Beets, broccoli, Brussels sprouts, cabbage, carrots, cauliflower, celery</b>		
<b>INSECT</b>	<b>PRE-HARVEST INTERVAL (DAYS)</b>	<b>CONTAINER COVERS</b>
Flea beetles Leafhoppers	1 - carrots 3 - broccoli, Brussels sprouts, cabbage, cauliflower, celery 7 - beet roots 14 - beet tops	1,923 - 4,000 sq. metres (20,699 - 43,056 sq. feet)
<b>Timing/Application:</b> Use the lower rate on young plants. Apply when insects and/or damage appear. Repeat at intervals of 7 to 10 days or as necessary.		

**CROP - Chinese cabbage, dandelion, endive, horseradish, kale, kohlrabi, lettuce, mustard greens, parsley, parsnips, radishes, rutabagas, salsify, spinach, Swiss chard, turnip, watercress**

INSECT	PRE-HARVEST INTERVAL (DAYS)	CONTAINER COVERS
Armyworms Corn earworm Diamondback moth (larvae) Imported cabbage worm Lygus bugs Meadow spittlebug Stink bugs	3 - kohlrabi 7 - horseradish, radish, rutabaga roots, salsify & turnip root 14 - Chinese cabbage, dandelion, endive, kale, leaf lettuce, mustard greens, parsley, parsnips, salsify tops, spinach, Swiss chard, turnip top, watercress	962 - 1,923 sq. metres (10,355 - 20,699 sq. feet)

**Timing/Application:** To avoid possible injury on lettuce, do not apply when foliage is wet or when rain or excessive humidity is expected during the next 2 days. Apply for thorough coverage when insects and/or damage appear. Repeat at intervals of 7 to 10 days as necessary.

**CROP - Corn (field & sweet)**

INSECT	PRE-HARVEST INTERVAL (DAYS)	CONTAINER COVERS
Corn earworm European corn borer Fall armyworm Northern corn rootworm adults	1	1,205 - 1,923 sq. metres (12,971 - 20,699 sq. feet)

**Timing/Application:** For larvae in whorls and for foliage feeders, treat the entire plant. Repeat as necessary. For insects attacking silks and ears, apply at intervals of 2 to 4 days starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on the severity of the infestation. Timing and good coverage are essential for effective control.

**CROP - Cucumber, melons, pumpkins, squash**

<b>INSECT</b>	<b>PRE-HARVEST INTERVAL (DAYS)</b>	<b>CONTAINER COVERS</b>
Cucumber beetles      Leafhoppers Cutworms (climbing)      Squash bugs Flea beetles	3	1,923 sq. metres (20,699 sq. feet)
<b>Timing/Application:</b> Apply for thorough coverage when insects and/or damage appear. Repeat at intervals of 7 to 10 days or as necessary. To avoid possible injury on tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the next 2 days.		

**CROP - Eggplant, pepper, potato, tomato**

<b>INSECT</b>	<b>PRE-HARVEST INTERVAL (DAYS)</b>	<b>CONTAINER COVERS</b>
Colorado potato beetle	1 - eggplant, pepper & tomato	4,000 sq. metres (43,056 sq. feet)
European corn borer Fall armyworm Tomato hornworm Tomato fruitworm	7 - potato	962 - 1,923 sq. metres (10,355 - 20,699 sq. feet)
Stink bugs Tarnished plant bug		800 - 962 sq. metres (8,611 - 10,355 sq. feet)
Cutworms (climbing)		962 - 1,205 sq. metres (10,355 - 12,971 sq. feet)
<b>Timing/Application:</b> Apply when insects and/or damage appear. Repeat at intervals of 7 to 10 days or as necessary. For cutworms, spray in a 25 to 30 cm band (10 to 12 inches) over the row.		

**FRUIT TREES**

<b>CROP - Apples, pears</b>		
<b>INSECT</b>	<b>PRE-HARVEST INTERVAL (DAYS)</b>	<b>CONTAINER COVERS</b>
Apple leafhopper Apple leafroller Codling moth Eyespotted bud moth	Mealybug Pistol casebearer Redbanded leafroller (1 <sup>st</sup> brood)	7  2,174 sq. metres (23,400 sq. feet)
Apple maggot Eastern tent caterpillar Fruittree leafroller Green fruitworm Pearleaf blister mite Pear psylla Pear slug	Plum curculio Redbanded leafroller (2 <sup>nd</sup> brood) Rust mites Tarnished plant bug Tentiform leafminer Wooly apple aphid	7  725 sq. metres (7,804 sq. feet)
<b>Timing/Application:</b> To avoid undesirable apple thinning, delay use until at least 30 days after full bloom. Apply for full coverage. For psylla, apply when eggs hatch or when young nymphs are present. For optimum maggot control, apply at 10 day intervals.		
<b>CROP - Apricots, peaches</b>		
<b>INSECT</b>	<b>PRE-HARVEST INTERVAL (DAYS)</b>	<b>CONTAINER COVERS</b>
Cat-facing insects Codling moth European earwig European fruit lecanium Fruittree leafroller Lesser peachtree borer	Oriental fruit moth Peach silver mite Peach twig borer Plum curculio Redbanded leafroller Scale insects	1 - peaches 3 - apricots  725 sq. metres (7,804 sq. feet)
<b>Timing/Application:</b> Apply for full coverage in normal spray schedules. For lesser peachtree borer, spray limbs and trunk thoroughly. For scale insects apply when crawlers are present.		

<b>CROP - Plums, cherries</b>			
<b>INSECT</b>	<b>PRE-HARVEST INTERVAL (DAYS)</b>	<b>CONTAINER COVERS</b>	
Codling moth Eastern tent caterpillar	Oak leafhopper Prune leafhopper	1	1,087 sq. metres (11,700 sq. feet)
Apple maggot Black cherry aphid Cherry fruit fly Cherry fruitworm Eye-spotted bud moth Fruittree leafroller	Lesser peachtree borer Mealy plum aphid Peach twig borer Plum curculio Redbanded leafroller Scale insects	1	725 sq. metres (7,804 sq. feet)
<b>Timing/Application:</b> Apply for full coverage in normal spray schedule. For scale insects, apply when crawlers are present. For lesser peachtree borer, thoroughly spray limbs and trunk area.			

**SMALL FRUITS**

<b>CROP - Blackberries, boysenberries, dewberries, loganberries, raspberries</b>			
<b>INSECT</b>	<b>PRE-HARVEST INTERVAL (DAYS)</b>	<b>CONTAINER COVERS</b>	
Blackberry leafminer Japanese beetle Leafhoppers	Leafroller, Aphid Rose stem girdler Spotted winged raspberry aphid	7	962 sq. metres (10,355 sq. feet)
<b>Timing/Application:</b> Apply when insects and/or damage appear. Repeat at intervals or 7 to 10 days or as necessary.			
<b>CROP - Blueberries</b>			
<b>INSECT</b>	<b>PRE-HARVEST INTERVAL (DAYS)</b>	<b>CONTAINER COVERS</b>	
Blueberry maggot Cranberry fruitworm Lecanium scale	1	1,205 sq. metres (12,971 sq. feet)	
<b>Timing/Application:</b> Apply 3 weeks before harvest and repeat in 10 days or as necessary.			



<b>CROP - Cranberries</b>		
<b>INSECT</b>	<b>PRE-HARVEST INTERVAL (DAYS)</b>	<b>CONTAINER COVERS</b>
Bluntnosed cranberry leafhopper Cranberry fruitworm Cutworms (climbing) Fireworms	1	714 - 800 sq. metres (7,685 - 8,611 sq. feet)
<b>Timing/Application:</b> Apply during late bloom. Repeat as necessary at intervals of 7 to 10 days.		
<b>CROP - Grapes</b>		
<b>INSECT</b>	<b>PRE-HARVEST INTERVAL (DAYS)</b>	<b>CONTAINER COVERS</b>
Grape berry moth Leafhoppers	3	800 - 962 sq. metres (8,611 - 10,355 sq. feet)
<b>Timing/Application:</b> Apply for full coverage when insects and/or damage appear. Repeat at intervals of 7 to 10 days or as necessary.		
<b>CROP - Strawberries</b>		
<b>INSECT</b>	<b>PRE-HARVEST INTERVAL (DAYS)</b>	<b>CONTAINER COVERS</b>
Meadow spittlebug Strawberry leafroller	1	800 - 1,923 sq. metres (8,611 - 20,699 sq. feet)
<b>Timing/Application:</b> Apply when insects and/or damage appear. Repeat at intervals of 7 to 10 days or as necessary.		

TREES AND ORNAMENTALS

<p style="text-align: center;"><b>CROP - Arborvitae, azalea, birch, boxwood, carnation, chrysanthemum, dogwood, elm, gladiolus, holly, hydrangea, juniper, lilac, maple, oak, pines, rose, zinnia</b></p>		
<b>INSECT</b>		<b>CONTAINER COVERS</b>
Bagworm Birch leafminer Blackvine weevil (taxus weevil) Blister beetles Boxelder bug Boxwood leafminers Cankerworms Flea beetle Elm leaf aphid Elm leaf beetle Japanese beetle June beetles Lace bugs	Leafhoppers Leafrollers Mealybugs Pine sawflies Plant bugs Psyllids Rose aphid Rose chafer Rose slug Scale insects Tent caterpillars Thrip (exposed) Willow leaf beetle	1,600 - 1,923 sq. metres (17,222 - 20,699 sq. feet)
<p><b>Timing/Application:</b> Spray in the early morning or evening. Repeat at intervals of 7 to 14 days. For scale insects, apply when crawlers are present. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. Do not use on Boston ivy, maidenhair fern or Virginia creeper.</p>		

**LAWN AND RECREATIONAL AREAS**

<b>SITE - Turf (ornamental and sports)</b>		
<b>INSECTS</b>		<b>CONTAINER COVERS</b>
Ants	Fleas	192 - 267 sq. metres (2,067 - 2,874 sq. feet)
Chinch bugs	Leafhoppers	
Climbing cutworms	Millipedes	
Earwigs	Mosquitoes	
Fall armyworms	Sod webworms	
<b>Timing/Application:</b> For best results mow lawn and apply after watering or rain. Ensure good penetration of turf. Repeat 2 to 3 weeks later if necessary.		
<b>SITE - Turf</b>		
<b>INSECTS</b>		<b>CONTAINER COVERS</b>
White Grubs - European chafer* Japanese beetle* (suppression of population)		172 - 238 sq. metres (1,851 - 2,562 sq. feet)
<b>Timing/Application:</b> Remove excessive thatch and water 24 hours before treatment if soil is dry. Repeat as necessary.		
*To optimize control of white grubs (European chafer and Japanese beetle) apply in late July/early August.		

**PRECAUTIONS:**

KEEP OUT OF REACH OF CHILDREN. HARMFUL IF INHALED OR SWALLOWED. Avoid breathing of spray. Do not take internally. Avoid contact with skin and eyes. Wear regular long-sleeved work clothing. Remove and wash contaminated clothing separately after pesticide use. Change to clean clothing daily. Wash hands and face before eating. Wash thoroughly after handling. AVOID CONTAMINATION OF FOOD, FEED, WATER SUPPLIES, STREAMS AND PONDS.

**BEE CAUTION:** MAY KILL HONEY BEES IN SUBSTANTIAL NUMBERS. This product is highly toxic to bees exposed to direct treatment or residues on crops. Do not spray when bees are active. Avoid application on flowers and fruit blossoms.

**PLANT RESPONSE AND PRECAUTIONS:** To avoid possible injury to tender foliage, do not apply to wet foliage or when rain or high humidity is expected during the next two days. Carbaryl injures Boston ivy, Virginia creeper and maidenhair fern. During early season, it may also injure Virginia and sand pines.

**FIRST AID INSTRUCTIONS:**

If contact with eyes, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. If contact with skin occurs, take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If swallowed, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**Note to Physicians: Emergency Information call 1-800-668-5669.**

**TOXICOLOGICAL INFORMATION:**

Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal. DO NOT use 2-PAM, opiates, or cholinesterase inhibiting drugs.

**STORAGE AND DISPOSAL:**

Store unused product in original container only, in a cool, dry area out of reach of children and animals. Do not store in areas where temperatures frequently exceed 38°C. Protect from freezing. Do not contaminate water, food or feed by storage or disposal. Do not reuse empty pesticide containers. Wrap empty container in newspaper and dispose of with household garbage.

Product Use Information: 1-800-668-5669  
Web Site: [www.scottscanada.ca](http://www.scottscanada.ca)

**Scotts No-Quibble Guarantee®**

If for any reason you, the consumer, are not satisfied after using this product, you are entitled to get your money back. Simply send us proof of purchase and we will mail you a refund cheque promptly.

® used under license by Scotts Canada Ltd.

™ used under license by Scotts Canada Ltd.

© 2006, Scotts Canada Ltd. World rights reserved.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.