



Reliance™ STS™ Toss-N-Go™ Herbicide

GROUP	2	HERBICIDE
--------------	----------	------------------

Dry Flowable

FOR SALE FOR USE ON SULFONYLUREA TOLERANT SOYBEANS IN EASTERN CANADA ONLY

COMMERCIAL

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENT: Chlorimuron ethyl 21.0%
Thifensulfuron methyl 12.0%

REGISTRATION NO. 25784 PEST CONTROL PRODUCTS ACT

Do not sell soluble bags separately.

NET CONTENTS: 171.2 grams
(4 x 42.8 grams water soluble bags)

Production Agriscience Canada Company
P. O. Box 730
7398 Queen's Line
Chatham, Ontario
N7M 5L1
519-352-6350

PRECAUTIONS

- KEEP OUT OF REACH OF CHILDREN.
- Avoid breathing spray mist.
- Avoid contact with skin, eyes and clothing.
- Keep outer bag and soluble bags away from water.
- Do not contaminate any body of water.
Wear long-sleeved shirt, long pants, socks, and chemical-resistant gloves and footwear during mixing, loading, application, clean-up and repair. Chemical-resistant gloves are not required while operating groundboom sprayers.
- DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.
- Apply only to agricultural crops when the potential for drift to areas of human habitation and human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, applications equipment, and sprayer settings.
- Gloves are not required during application within a closed cab.

FIRST AID

IN CASE OF CONTACT WITH EYES: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IN CASE OF CONTACT WITH SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

For medical emergencies call 1-800-667-3852 (24 hours).

TOXICOLOGICAL INFORMATION: Treat symptomatically.

ENVIRONMENTAL PRECAUTIONS

TOXIC to non-target terrestrial plants. Observe buffer zones specified under DIRECTIONS FOR USE.

TOXIC to freshwater plants and algae. Observe buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats:

Avoid application to areas with a moderate to steep slope, compacted soil or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

STORAGE

Store product in original container only, away from other pesticides, fertilizer, food, seed or feed. Not for use or storage in or around the home. Keep container closed. Keep water soluble bags in original outer bag away from moisture.

DISPOSAL

1. Make the empty container unsuitable for further use.
2. Dispose of the container in accordance with provincial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

The seller warrants that the purchase by the buyer and the use of this product, as such, will not infringe on any Canadian patent.

[[DIRECTIONS FOR USE LABEL]]



Reliance™ STS™ Toss-N-Go™ Herbicide

GROUP	2	HERBICIDE
--------------	----------	------------------

Dry Flowable

FOR SALE FOR USE ON SULFONYLUREA TOLERANT SOYBEANS IN EASTERN CANADA ONLY

COMMERCIAL

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENT: Chlorimuron ethyl 21.0%
Thifensulfuron methyl 12.0%

REGISTRATION NO. 25784 PEST CONTROL PRODUCTS ACT

Do not sell soluble bags separately.

NET CONTENTS: 171.2 grams
(4 x 42.8 grams water soluble bags)

Production Agriscience Canada Company
P. O. Box 730
7398 Queen's Line
Chatham, Ontario
N7M 5L1
519-352-6350

PRECAUTIONS

- KEEP OUT OF REACH OF CHILDREN.
- Avoid breathing spray mist.
- Avoid contact with skin, eyes and clothing.
- Keep outer bag and soluble bags away from water.
- Do not contaminate any body of water.
- Wear long-sleeved shirt, long pants, socks, and chemical-resistant gloves and footwear during mixing, loading, application, clean-up and repair. Chemical-resistant gloves are not required while operating groundboom sprayers.

FIRST AID

IN CASE OF CONTACT WITH EYES: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IN CASE OF CONTACT WITH SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF INGESTED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

For medical emergencies call 1-800-667-3852 (24 hours).

TOXICOLOGICAL INFORMATION: Treat symptomatically.

IMPORTANT

Injury to or loss of desirable trees or vegetation may result from failure to observe the following:

Do not apply, drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not use on lawns, walks, driveways, tennis courts or similar areas. Prevent drift of spray to desirable plants. Do not contaminate any body of water, including irrigation water that may be used on other crops. Carefully observe sprayer clean-up instructions, as spray tank residue may damage crops other than soybeans.

DO NOT apply using aerial application equipment.

APPLY RELIANCE™ STS™ Toss-N-Go™ Herbicide WITH A RECOMMENDED SURFACTANT.

Do not handle pesticides with bare hands. Chemical resistant gloves significantly reduce hand exposure. Wear chemical resistant gloves for mixing/loading/clean-up operations and when making sprayer and nozzle repairs and adjustments. Do not use leather or cloth gloves. Wear coveralls or long-sleeve shirt and long pants for mixing/loading/applying and during clean-up and repair activities. Gloves are not required during application within a closed cab.

GENERAL INFORMATION

Reliance STS Toss-N-Go Herbicide is recommended for selective postemergence control of broadleaf weeds ONLY in sulfonyleurea tolerant soybean varieties (e.g. varieties designated as "STS"). USE OF RELIANCE STS TOSS-N-GO ON SOYBEAN VARIETIES NOT DESIGNATED AS SULFONYLUREA TOLERANT MAY CAUSE SEVERE CROP INJURY. Due to risk of crop injury, do not use Reliance STS Toss-N-Go Herbicide on crop grown from seed saved from previous year's production (bin run seed). Information on sulfonyleurea tolerant soybean varieties may be obtained from your seed supplier or Production Agriscience Canada Company representative.

Reliance STS Toss-N-Go Herbicide is formulated as a dry flowable granule to be mixed in water and applied as a uniform broadcast spray. It is non-corrosive, non-flammable, non-volatile, and does not freeze.

Reliance STS Toss-N-Go Herbicide must be applied with a recommended non-ionic surfactant early postemergence to the main flush of actively growing weeds. Warm, moist growing conditions promote active weed growth and enhance the activity of Reliance STS Toss-N-Go Herbicide by allowing maximum foliar uptake and contact activity. Weeds hardened off by environmental stress may not be adequately controlled and regrowth may occur. Reliance STS Toss-N-Go Herbicide may only be applied using ground equipment. For best results ensure thorough spray coverage of target weeds.

Reliance STS Toss-N-Go Herbicide rapidly stops growth of susceptible weeds. However, typical symptoms (discolouration) of dying weeds may not be noticeable for 1 to 3 weeks after application, depending on growing conditions and weed susceptibility. Favourable growing conditions following treatment promote the activity of Reliance STS Toss-N-Go Herbicide while cold, dry conditions delay the activity.

Degree of control and duration of effect depend on weed sensitivity, weed size, crop competition, growing conditions and spray coverage.

DIRECTIONS FOR USE

- Thifensulfuron-methyl is formulated as a wettable granule or a soluble granule sealed within a water-soluble package (WSP). Drop intact WSP directly into spray tank. The WSP and pesticide will dissolve readily in water.
- **DO NOT** allow the WSP to become wet prior to use. Do not handle individual WSP with wet hands or wet gloves as this may cause breakage. Do not remove WSP from overwrap container except for immediate use.
- **DO NOT** open WSP or attempt to remove the contents from them. The contents are not designed to be re-measured or subdivided.
- **DO NOT** use opened or punctured WSP.
- If broken WSP is found when the over-wrap bag is opened, avoid contact with, and inhalation of the product. Handle with chemical-resistant gloves and dispose of broken WSP according to the DISPOSAL section.
- **DO NOT** enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.
- Apply only to agricultural crops when the potential for drift to areas of human habitation and human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, applications equipment, and sprayer settings.

Field sprayer application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Boom height must be 60 cm or less above the crop or ground.

BUFFER ZONES

Spot treatments using hand-held equipment DO NOT require a buffer zone.

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands) and sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands).

Method of application	Crop	Buffer Zones (metres) Required for the Protection of:		
		Freshwater Habitat of Depths:		Terrestrial Habitat
		Less than 1 m	Greater than 1 m	
Field Sprayer	Soybean	1	1	20

For tank mixes, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The buffer zones for this product, with the exception of the freshwater habitat, can be modified based on weather conditions and spray equipment configuration by accessing the Buffer Zone Calculator on the Pest Management Regulatory Agency web site.

ENVIRONMENTAL PRECAUTIONS

- Toxic to non-target terrestrial plants. Observe buffer zones specified under DIRECTIONS FOR USE.
- Toxic to freshwater plants and algae. Observe buffer zones specified under DIRECTIONS FOR USE.
- To reduce runoff from treated areas into aquatic habitats:
 - Avoid application to areas with a moderate to steep slope, compacted soil or clay.
 - Avoid application when heavy rain is forecast.
 - Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

RATES AND TIMING

Optimum timing of Reliance STS Toss-N-Go Herbicide for annual broadleaf weed control is when soybeans are in the 1 to 3 trifoliolate stage of growth. Do not apply Reliance STS Toss-N-Go before the first trifoliolate has fully expanded or after the initiation of flowering. Reliance STS Toss-N-Go can be applied up to 60 days before harvest.

Use 27.6 or 42.8 grams of Reliance STS Toss-N-Go per hectare. A non-ionic surfactant such as Agral 90, Citowett Plus, or Ag-Surf must also be used at 0.2% v/v. The optional addition of liquid fertilizer containing 28% nitrogen to the spray mixture at the rate of 2 litres per hectare may improve control of velvetleaf and should be added after the surfactant.

For best results apply to young actively growing weeds before the soybean canopy closes. Weeds that emerge after treatment will not be controlled.

If rain occurs after application, control may be reduced. Two to four hours of dry weather are needed to allow Reliance STS Toss-N-Go Herbicide to be absorbed by weed foliage.

WEEDS CONTROLLED BY 27.6 GRAMS PER HECTARE RELIANCE STS TOSS-N-GO

Weeds	Leaf Stage	Height (cm)
Redroot Pigweed	2-8	10 cm tall or across
Lady's Thumb	2-8	"
Wild Mustard	2-8	"

One 171.2 gram container of Reliance STS Toss-N-Go Herbicide treats 6.2 ha at 27.6 grams/ha.

WEEDS CONTROLLED BY 42.8 GRAMS PER HECTARE RELIANCE STS TOSS-N-GO

Weeds	Leaf Stage	Height (cm)
Common Ragweed	2-6	10 cm tall or across
Velvetleaf	2-6	"
Lamb's-Quarters	2-6	"

Plus those weeds controlled by Reliance STS Toss-N-Go at 27.6 grams per hectare.

One 171.2 gram container of Reliance STS Toss-N-Go Herbicide treats 4 ha at 42.8 grams/ha.

WEEDS CONTROLLED BY RELIANCE STS TOSS-N-GO HERBICIDE + ASSURE II Herbicide

For the control of Annual Grasses, Redroot Pigweed, Lady's-Thumb, Wild Mustard, Common Ragweed, Velvetleaf and Lamb's-Quarters in sulfonylurea tolerant soybean varieties (varieties designated "STS"), apply a tank mixture of Assure II Herbicide plus Reliance STS Toss-N-Go Herbicide. Use Assure II Herbicide at 0.38 to 0.5 litres per hectare plus Reliance STS Toss-N-Go Herbicide at 42.8 grams per hectare. Add Sure-Mix at 0.5% v/v (0.5 litre per 100 litres of spray solution).

Use the 0.5 litre per hectare rate of Assure II Herbicide when quackgrass and yellow foxtail are present in the field. The tank-mix of Reliance STS Toss-N-Go Herbicide and Assure II Herbicide may result in reduced control of these two grassy weeds.

Consult the Assure II Herbicide label for specified rates and optimum leaf stage timing for grassy weeds. If leaf stages of the grass and broadleaf weeds do not coincide, a sequential application of the grass and broadleaf herbicides is required to ensure satisfactory control.

Consult the Assure II Herbicide label for general information and use precautions.

MIXING INSTRUCTIONS

1. Fill clean tank about 1/4 full with fresh water.
2. Turn on full agitation.
3. Add the proper amount of Reliance STS Toss-N-Go Herbicide to the water in the spray tank with the agitator running. Maintain full agitation until water soluble bags are completely dissolved and product is fully dispersed. Continuous agitation is required to keep the product in suspension.
4. After Reliance STS Toss-N-Go Herbicide has been well mixed and is in suspension add the proper amount (if applicable) of Assure II Herbicide.
5. Add a recommended non-ionic surfactant such as Citowett Plus, Agral 90, or Ag-Surf at 2 L per 1000 L spray solution (0.2% v/v). If tank mixing with Assure II Herbicide, add Sure-Mix at 0.5% v/v (0.5 litre per 100 litres of spray solution).
6. Fill the remainder of the spray tank with clean water.

On repeat tank loads, ensure that the amount of spray solution left in the tank from the previous load is less than 10% of the volume about to be mixed.

Agitation is required for uniform mixing and application. The optimum water volume for Reliance STS Toss-N-Go Herbicide application is 140-190 litres of water per hectare (minimum of 100 litres of water per hectare). Flat fan nozzles are recommended. Use 50 mesh filter screens or larger. Use spray preparation of Reliance STS Toss-N-Go Herbicide within 24 hours or product degradation may occur resulting in a loss of weed control. Use vigorous agitation to thoroughly disperse spray mixtures that have been allowed to stand in the tank.

NOTE: Reliance STS Toss-N-Go Herbicide will degrade in acidic or highly alkaline water. Mix no more than can be used in one day. If spraying is interrupted, thoroughly re-agitate the spray mixture before resuming spraying.

NOTE: Do not mix the soluble bags with liquid fertilizers, substances that contain boron or substances that release free chlorine. Mixing the water soluble bags with any of these compounds will result in an insoluble substance in the tank.

Do not mix or load where there is potential to contaminate wells or aquatic systems.

SPRAYER CLEAN-UP

To avoid subsequent injury to crops other than soybeans immediately after spraying and prior to spraying other crops thoroughly remove all traces of Reliance STS Toss-N-Go Herbicide from mixing and spray equipment as follows:

1. Drain tank; then flush tank, boom and hoses with clean water for a minimum of ten minutes. Visually inspect tank to assure removal of all visible residues of RELIANCE™ STS™ Toss-N-Go® Herbicide. If necessary, repeat step 1. **DO NOT CLEAN SPRAY EQUIPMENT NEAR WELLS, AQUATIC SYSTEMS OR OTHER WATER SOURCES OR NEAR TO DESIRABLE VEGETATION.**
2. Fill tank with clean water while adding 1 litre household ammonia (containing a minimum 3% ammonia) per 100 litres of water. Flush solution through boom and hoses, and then add more water to completely fill tank. Allow to sit for 15 minutes with agitation. Again flush the hoses, boom and nozzles with the cleaning solution and drain tank.
3. Remove the nozzles and screens and clean separately in a bucket containing cleaning agent and water.
4. Repeat Step 2.
5. Thoroughly rinse the tank with clean water for a minimum of 5 minutes, flushing water through the hoses and boom.

REPLANTING TO OTHER CROPS

Reliance STS Toss-N-Go is degraded by natural soil processes, and field tests have shown that the following crops may safely be planted at the prescribed interval following soybeans that have been treated with Reliance STS Toss-N-Go. The recropping intervals are dependent on the pH of the soil, as well as the sensitivity of the specific rotational crop.

Replanting Interval - Months

Soil Ph	Field Corn	Winter Wheat	White Beans	Alfalfa	Soybeans*
≤ 7.4	10	3	10	10	10
≥ 7.5	-----	Do Not Apply-	-----		

*(includes sulfonylurea tolerant soybean varieties)

Soil pH may vary significantly within a single field. It may be affected by soil amendments (i.e. lime, fertilizers, etc.) or soil conditions (i.e. eroded knolls, etc.)

For other crops, a field bioassay is recommended before planting. A successful field bioassay means growing to maturity a test strip of the crop(s) intended for production the following year.

FIELD BIOASSAY

Select a representative area or areas of the field previously treated with Reliance STS Toss-N-Go Herbicide to plant your bioassay crop(s). Be sure to consider factors such as size of field, soil texture, drainage and turn-around areas when selecting the site(s) that are most representative of the soil conditions in the field. On large fields, more than one site may be needed in order to obtain reliable results.

Plant the test strips perpendicular to the direction in which the field was sprayed. The strips should be long enough to cross the width of several spray swaths. Large test strip areas are more reliable than small ones.

Use standard tillage and seeding equipment to plant the bioassay. Prepare a seed bed and plant the crops and varieties you want the option of growing the following year. It is important to use the same planting time, conditions, techniques and cultural practices you normally use to plant and grow the bioassay crop(s). Also plant into an adjacent area not treated with Reliance STS Toss-N-Go Herbicide to use as a comparison.

As the crop(s) emerges and grows, examine these key points in Reliance STS Toss-N-Go Herbicide treated and non-treated areas:

-crop stand	-root development	-rate of growth
-plant colour and vigour	-yield	

Allow the bioassay crop(s) to grow to maturity while making your observations. Do not overspray the test strips with herbicides that may damage the bioassay crop(s). If the bioassay indicates that Reliance STS Toss-N-Go Herbicide residues are still present, continue cropping only to those crops listed on the label and do not rotate to other crops until bioassay results indicate that susceptible crops are growing normally.

USE PRECAUTIONS

Do not apply Reliance STS Toss-N-Go Herbicide during periods of intense rainfall or to soils saturated with water. Do not apply directly to standing or running water. Do not apply in areas where surface water from the treatment site can run off to adjacent cropland, either planted or to be planted, or into bodies of water (such as streams, ponds, rivers, lakes, and irrigation water), wetlands, or wells. Applications should only be made when there is no hazard of spray drift contaminating non-target land areas since very small quantities of the Reliance STS Toss-N-Go Herbicide spray solution may severely injure susceptible crops and other sensitive terrestrial habitats during both growing and dormant periods.

Only apply to sulfonylurea tolerant soybean varieties (e.g. varieties designated as "STS") or crop injury may result. Do not apply to soybeans that have been stressed by severe weather conditions, frost, low fertility, drought, water saturated soil, disease or insect damage as crop injury may result.

Under extreme weather conditions, such as hot, dry weather, excessive moisture, or frost, weed control may be reduced.

Rainfall within 2-4 hours after application of Reliance STS Toss-N-Go Herbicide may reduce weed control.

Apply Reliance STS Toss-N-Go Herbicide when the minimum temperature in the 24 hours before and after application is above 5 degrees C. If soybeans have been injured by frost, wait at least 48 hours before applying Reliance STS Toss-N-Go Herbicide.

Reliance STS Toss-N-Go Herbicide must not be applied within 60 days of harvest.

Make only one application of Reliance STS Toss-N-Go Herbicide per year.

Do not graze the treated crop or cut for hay; sufficient data is not available to support such use.

Do not apply Reliance STS Toss-N-Go Herbicide to sulfonylurea tolerant soybeans grown on soil of pH ≥ 7.5 due to risk of rotational crop injury.

As this product is not registered for the control of pests in aquatic systems, **DO NOT** use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

RESISTANCE-MANAGEMENT RECOMMENDATIONS

For resistance management, Reliance STS Toss-N-Go Herbicide is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to Reliance STS Toss-N-Go Herbicide and other Group 2 herbicides. The resistance biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- ◆ Where possible rotate the use of Reliance STS Toss-N-Go Herbicide or other Group 2 herbicides within a growing season or among growing seasons with different herbicide groups that control the same weeds in a field.
- ◆ Use tank mixtures with herbicides from different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- ◆ Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- ◆ Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group.
- ◆ Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- ◆ Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.

- ◆ For further information or to report suspected resistance, contact your local Production Agriscience Canada Company representative or the Production Agriscience Canada Company hotline at 1-800-667-3852 for further information

STORAGE

Store product in original container only, away from other pesticides, fertilizer, food, seed or feed. Not for use or storage in or around the home. Keep container closed. Keep water soluble bags in original outer bag away from moisture.

DISPOSAL

1. Make the empty container unsuitable for further use.
2. Dispose of the container in accordance with provincial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

The seller warrants that the purchase by the buyer and the use of this product, as such, will not infringe on any Canadian patent.

™Trademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners.

All other products mentioned are trademarks of their respective companies.

[[WATER SOLUBLE BAG]]



Reliance™ STS™ Toss-N-Go™ Herbicide

DRY FLOWABLE

KEEP OUT OF REACH OF CHILDREN

SEE DIRECTIONS FOR USE BROCHURE
FOR COMPLETE PRECAUTIONARY STATEMENTS

ACTIVE INGREDIENT:	Chlorimuron ethyl	21.0%
	Thifensulfuron methyl	12.0%

NET CONTENTS: 42.8 grams