

2024-1378
2024-06-04

SELECT[®]

EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE

GROUP	1	HERBICIDE
-------	---	-----------

CONTAINS CLETHODIM

POST-EMERGENCE HERBICIDE FOR CONTROL OF GRASSES IN SEEDLING ALFALFA, LEGUME VEGETABLES (SUCCULENT OR DRIED), EDAMAME,, HIGHBUSH BLUEBERRY, CANOLA, BASIL, CORIANDER, DILL, CARAWAY, FENUGREEK, CRANBERRY, FLAX (including low linolenic acid varieties), ORIENTAL (BROWN) MUSTARD (condiment and oilseed types), *BRASSICA CARINATA*, YELLOW MUSTARD, DRY BULB ONIONS, DRY BULB SHALLOTS AND GARLIC, GREEN ONION, LEEK, POTATOES, PRAIRIE CARNATIONS, SAFFLOWER, SOYBEANS, SPINACH, SUNFLOWERS, GARDEN (TABLE) BEET, PARSNIP, CARROT, RADISH, CELERIAC, RUTABAGA, CHERRY (tart and sweet), CERTAIN HERBS, SPICES, HOPS, CHINESE CABBAGE (napa), CELERY AND BUCKWHEAT.

Commented [NGRaD1]: Per OMAFRA request: grower contacted them who grows red and gold table beets. Gold beets are the same species/crop, but they technically are not 'red' beets.

COMMERCIAL (AGRICULTURAL)

ACTIVE INGREDIENT: CLETHODIM 240 g/L

REGISTRATION NO. 22625 PEST CONTROL PRODUCTS ACT

WARNING: EYE AND SKIN IRRITANT

**KEEP OUT OF REACH OF CHILDREN
READ THE LABEL AND BOOKLET BEFORE USING**

NET CONTENTS: (250 mL - 9 LITRES)

For Product Use Information Call 1-800-438-6071

**UPL AgroSolutions Canada Inc.
PO Box 12219**

Research Triangle Park, NC 27709

**FOR CHEMICAL EMERGENCY: spill, leak, fire, exposure, or accident call
CHEMTREC 1-800-424-9300**

2024-1378

2024-06-04

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. CAUSES EYE AND SKIN IRRITATION. Avoid contact with skin, eyes and clothing. Wash concentrate from skin or eyes immediately. Do not inhale fumes. Avoid breathing vapours or spray mist. After use, wash hands and other exposed skin. When using, do not eat, drink or smoke. Remove and launder contaminated clothing separately from household laundry before reuse. Store the container tightly closed and away from seeds, feeds, fertilizer, plants and foodstuffs. Observe appropriate provincial buffer zones around bodies of water and wetland areas. Wear goggles or face shield, rubber apron, socks and rubber boots, long-sleeved shirt, long pants, and chemical-resistant gloves when handling the concentrate, mixing, loading, or during clean-up and repair. Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during application. Gloves are not required during application within a closed cab or cockpit.

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

ENVIRONMENTAL PRECAUTIONS:

TOXIC to aquatic organisms and non-target terrestrial plants. Observe buffer zones specified under DIRECTIONS FOR USE.

Toxic to certain beneficial insects. Minimize spray drift to reduce harmful effects on beneficial insects in habitats next to the application site such as hedgerows and woodland.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

This product contains aromatic petroleum distillates that are toxic to aquatic organisms.

The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (for example, sandy soil) and/or the depth to the water table is shallow.

FIRST AID:

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do **not** give any liquid to the person. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

2024-1378

2024-06-04

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Treat symptomatically. This product contains petroleum distillates. Vomiting may cause aspiration pneumonia. Danger from lung aspiration of petroleum based solvents must be weighed against toxicity when considering emptying stomach.

STORAGE AND DISPOSAL

STORAGE: To prevent contamination, store this product away from food or feed.

DISPOSAL:

RECYCLABLE CONTAINER DISPOSAL: Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

DISPOSAL OF UNUSED, UNWANTED PRODUCT: For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

UPL, the UPL logo, SELECT, AMIGO and INTERLINE are registered trademarks of a UPL Corporation Limited Group Company.

©2023 UPL Corporation Limited Group Company.

2024-1378
2024-06-04

SELECT®

EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE

GROUP	1	HERBICIDE
-------	---	-----------

CONTAINS CLETHODIM

POST-EMERGENCE HERBICIDE FOR CONTROL OF GRASSES IN SEEDLING ALFALFA, LEGUME VEGETABLES (SUCCULENT OR DRIED), EDAMAME, HIGHBUSH BLUEBERRY, CANOLA, BASIL, CORIANDER, DILL, CARAWAY, FENUGREEK, CRANBERRY, FLAX (including low linolenic acid varieties), ORIENTAL (BROWN) MUSTARD (condiment and oilseed types), *BRASSICA CARINATA*, YELLOW MUSTARD, DRY BULB ONIONS, DRY BULB SHALLOTS, GARLIC, GREEN ONION, LEEK, POTATOES, PRAIRIE CARNATIONS, SAFFLOWER, SOYBEANS, SPINACH, SUNFLOWERS, GARDEN (TABLE) BEET, PARSNIP, CARROT, RADISH, CELERIAC, RUTABAGA, CHERRY (tart and sweet), CERTAIN HERBS, SPICES, HOPS, CHINESE CABBAGE (NAPA), CELERY and BUCKWHEAT.

COMMERCIAL (AGRICULTURAL)

ACTIVE INGREDIENT: CLETHODIM 240 g/L

REGISTRATION NO. 22625 PEST CONTROL PRODUCTS ACT

WARNING: EYE AND SKIN IRRITANT

**KEEP OUT OF REACH OF CHILDREN
READ THE LABEL AND BOOKLET BEFORE USING**

NET CONTENTS: (250 mL - 9 LITRES)

For Product Use Information Call 1-800-438-6071

UPL AgroSolutions Canada Inc.
630 Freedom Business Center, Suite 402
King of Prussia, PA 19406

**FOR CHEMICAL EMERGENCY: spill, leak, fire, exposure, or accident call
CHEMTREC 1-800-424-9300**

2024-1378

2024-06-04

GENERAL INFORMATION:

SELECT® EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE is a selective post-emergence herbicide for control of a broad range of grasses in seedling alfalfa, legume vegetables (succulent or dried), edamame, highbush blueberry, canola, basil, coriander, dill, caraway, fenugreek, cranberry, flax (including low linolenic acid varieties), oriental (brown) mustard (condiment and oilseed types), *Brassica carinata*, safflower, yellow mustard, dry bulb onions, dry bulb shallot, garlic, green onion, leek, potatoes, prairie carnation, soybeans, spinach, sunflowers, garden (table) beet, parsnip, carrot, radish, celeriac, rutabaga, cherry (tart and sweet), certain herbs, spices, hops, Chinese cabbage (napa), celery and buckwheat.

SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE needs to be applied with an adjuvant (see Application Instructions under the Directions For Use).

SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE is a systemic herbicide which is translocated from the treated foliage to the growing points of the leaves, shoots and roots. Uptake into the plant is primarily through its leaves. Thorough coverage of the foliage is important for consistent grass control. Susceptible grasses that are actively growing will undergo a burn-back and colour change after treatment. Leaf foliage will first change from green to yellowish, then purplish and finally brown colour. The time required for complete control is normally 7 to 21 days following treatment, depending on growing conditions and crop competition.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. CAUSES EYE AND SKIN IRRITATION. Avoid contact with skin, eyes and clothing. Wash concentrate from skin or eyes immediately. Do not inhale fumes. Avoid breathing vapours or spray mist. After use, wash hands and other exposed skin. When using, do not eat, drink or smoke. Remove and launder contaminated clothing separately from household laundry before reuse. Store the container tightly closed and away from seeds, feeds, fertilizer, plants and foodstuffs. Observe appropriate provincial buffer zones around bodies of water and wetland areas. When handling the concentrate, mixing, loading or during cleanup and repair activities, wear goggles or face shield, rubber apron, chemically resistant gloves, rubber boots, long sleeved shirt and long legged pants. During application, wear long sleeved shirt, long legged pants and chemical-resistant gloves. Gloves are not required during application within a closed cab or cockpit.

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

ENVIRONMENTAL PRECAUTIONS:

2024-1378

2024-06-04

TOXIC to aquatic organisms and non-target terrestrial plants. Observe buffer zones specified under DIRECTIONS FOR USE.

Toxic to certain beneficial insects. Minimize spray drift to reduce harmful effects on beneficial insects in habitats next to the application site such as hedgerows and woodland.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

This product contains aromatic petroleum distillates that are toxic to aquatic organisms.

The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (for example, sandy soil) and/or the depth to the water table is shallow.

FIRST AID:

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do **not** give any liquid to the person. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Treat symptomatically. This product contains petroleum distillates. Vomiting may cause aspiration pneumonia. Danger from lung aspiration of petroleum based solvents must be weighed against toxicity when considering emptying stomach.

STORAGE AND DISPOSAL

STORAGE: To prevent contamination, store this product away from food or feed.

DISPOSAL:

RECYCLABLE CONTAINER DISPOSAL: Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site.

2024-1378

2024-06-04

Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

DISPOSAL OF UNUSED, UNWANTED PRODUCT: For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

DIRECTIONS FOR USE:

SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE is to be used only on seedling alfalfa, legume vegetables (succulent or dried), edamame, highbush blueberry, canola, basil, coriander, dill, caraway, fenugreek, cranberry, flax (including low linolenic acid varieties), oriental (brown) mustard (condiment and oilseed types), *Brassica carinata*, safflower, yellow mustard, dry bulb onions, dry bulb shallots, garlic, green onions, leeks, potatoes, prairie carnations, soybeans, spinach, sunflowers, garden (table) beet, parsnip, carrot, radish, celeriac, rutabaga, cherry (tart and sweet), certain herbs, spices, hops, Chinese cabbage (napa), celery and buckwheat.

Not for use in greenhouses.

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

A 30-day plant-back interval should be observed for all unlabelled crops.

APPLICATION INSTRUCTIONS

ADJUVANTS

Apply **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** with either AMIGO Adjuvant at 0.5 to 1% v/v or Journey HSOC at 0.5% v/v. Consult the adjuvant labels for complete directions for use and application instructions such as rate and registered crops.

TIME OF APPLICATION:

Apply **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** when the annual grasses and volunteer cereals are in the 2 to 6 leaf stage as specified under Application Rates of **SELECT EMULSIFIABLE CONCENTRATE POST-**

2024-1378

2024-06-04

EMERGENCE HERBICIDE. Most effective control is achieved when application is made prior to tillering when annual grasses are small and actively growing.

For suppression or control of quackgrass, apply **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** when the quackgrass is in the 2 to 6 leaf stage and 6 to 15 cm in height. Most effective results are achieved when application is made at the 3 to 5 leaf stage, when the canopy is uniform and actively growing.

SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE will be less effective when plants are stressed by lack of moisture, excessive moisture, low-temperature and/or very low relative humidity. Regrowth by tillering may occur if application is made under any of the above stress conditions. In wide row crops, where the canopy may be slow to close, cultivation may be necessary to control grasses that emerge after treatment.

For chickpeas, apply **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** at a maximum of one application per season before the crop reaches the 9 node stage (18 cm height maximum) and when the annual grasses and volunteer cereals are in the 2 to 6 leaf stage.

For dill and fenugreek, apply **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** post-emergence, when dill or fenugreek is in the 3 to 5 leaf stage at a maximum of one application per season.

For coriander, or prairie carnation, apply **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** post-emergence, when coriander or prairie carnation is in the 2 to 5 leaf stage. A maximum of one application may be made per year.

For safflower, apply **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** post-emergence, when safflower is in the 6 to 8 leaf stage at a maximum of one application per season.

Cranberry is tolerant to **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** at all stages of growth, however, do not apply **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** between the hook stage and full fruit set.

For dry bulb onions, dry bulb shallots and garlic, apply **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** post-emergence, when crops are in the 1-4 leaf stage.

For green onion, apply **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** post emergence, when crop is in the 2 – 3 leaf stage.

For leek, apply **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** post-emergent at least 2 weeks after transplanting when crop is well established and has produced one new fully expanded leaf.

2024-1378

2024-06-04

Apply **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** only to established sweet and tart cherry trees. The pre-harvest interval outlined in the Crop Recommendations table must be followed to avoid excessive crop residues.

All other labelled crops are tolerant to **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** at all stages of growth. However, the pre-harvest interval outlined in the Crop Recommendations table must be followed to avoid excessive crop residues.

APPLICATION RATES OF SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE:

Rates of use are given below to control various grass species. Apply **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** with either AMIGO Adjuvant or JOURNEY HSOC. Refer to crop section for the maximum use rate on each crop.

Grass Species	Leaf Stage	Application Rate of SELECT® EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE	Application Rate ¹ of	
			AMIGO® Adjuvant	JOURNEY HSOC
*Foxtail (green, yellow)	2 to 4	0.125 L/ha	0.5 % v/v	0.5 % v/v
Wild oats				
Volunteer cereals (wheat, barley, oats)				
Barnyard Grass	2 to 6	0.125 L/ha	0.5 % v/v	0.5 % v/v
Witchgrass				
Fall panicum				
Proso millet				
Volunteer Corn				
Volunteer canary grass				
Wild Oats	2 to 6	0.19 L/ha	0.5 % v/v	0.5 % v/v
Volunteer cereals (wheat, barley, oats)				
Foxtail (green, yellow)				
Persian darnel				
Crabgrass (smooth and large)				
Proso millet				
Witchgrass				
Fall panicum				
Barnyard grass				
Volunteer corn				
Volunteer canary grass				

2024-1378

2024-06-04

Japanese brome Foxtail barley Quackgrass suppression				
Quackgrass control**	2 to 6	0.38 L/ha	1.0 % v/v	n/a
Annual bluegrass suppression	2 to 6	0.38 L/ha	0.5 % v/v	0.5 % v/v
Japanese brome** Foxtail barley**	2 to 6 leaf maximum 3 tillers	0.38 L/ha	1.0 % v/v	n/a

¹ Choose either AMIGO ADJUVANT or JOURNEY HSOC

***SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** applied at 0.125 L/ha for the control of weeds listed in this section of the table should only be applied under the following conditions:

- Good crop stand
- Early application (as in above table, prior to tillering)
- Do not tank-mix with other pesticides
- Do not apply to volunteer winter cereals
- Light to moderate weed infestation
- Adequate moisture and fertility
- Absence of stress, good growing conditions

If any one of the above conditions are not present at the time of application, use 0.19 L/ha rate of **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE**.

** For adequate control, use a minimum spray volume of 100L / ha.

2024-1378

2024-06-04

CROP RECOMMENDATIONS:

Crops	Maximum Application Rate of SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE	Application Method ⁷	Pre-Harvest Interval
Seedling Alfalfa	0.38 L/ha	Ground application only	30 days
Pigeon Pea ^{1,2}	0.19 L/ha	Ground and aerial application	75 days
<p>Edamame and Edible-podded legume vegetables (Crop subgroup 6A): Bean (<i>Phaseolus</i> spp.) (includes runner bean, snap bean, wax bean), Bean (<i>Vigna</i> spp.) (includes asparagus bean, Chinese longbean, moth bean, yardlong bean), Jackbean (<i>Canavalia ensiformis</i>), Pea (<i>Pisum</i> spp.) (includes dwarf pea, edible-pod pea, snow pea, sugar snap pea), Sword bean (<i>Canavalia gladiata</i>).</p> <p>Succulent shelled peas and beans (Crop subgroup 6B): Bean (<i>Phaseolus</i> spp.) (includes lima bean, green bean), Bean (<i>Vigna</i> spp.) (includes blackeyed pea, cowpea, southern pea), Broad bean (fava bean) (<i>Vicia faba</i>), Pea (<i>Pisum</i> spp.) (includes English pea, garden pea, green pea), Pigeon pea (<i>Cajanus cajan</i>)^{1,2,5}</p>	0.19 L/ha	Ground application only	21 days

2024-1378

2024-06-04

<p>Dried Shelled Pea and Bean (Crop subgroup 6C) (except soybean, field pea, lentils, pigeon pea) including: Bean (<i>Lupinus</i> spp.) (includes grain lupin, sweet lupin, white lupin and white sweet lupin); bean (<i>Phaseolus</i> spp.) (includes field bean, kidney bean, lima bean (dry), navy bean, pinto bean, tepary bean); bean (<i>vigna</i> spp.) (includes adzuki bean, blackeyed pea, catjang, cowpea, crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean); broad bean (fava bean) (<i>Vicia faba</i>); chickpea (garbanzo bean) (<i>Cicer arietinum</i>); guar (<i>Cyamopsis tetragonoloba</i>); lablab bean (hyacinth bean) (<i>Lablab purpureus</i>); ^{1,2}</p>	<p>0.19 L/ha</p>	<p>Ground and aerial application</p>	<p>60 days</p>
<p>Caraway, Coriander (cilantro)^{1,5}</p>	<p>0.38 L/ha</p>	<p>Ground application only</p>	<p>60 days for seed 30 days for leaves</p>
<p>Angelica; balm (lemon balm); basil; borage; burnet; chamomile; burnet; chamomile; catnip; chervil; clary; costmary; culantro (leaf); horehound; hyssop; lavender; lovage; marigold; marjoram (includes sweet or annual marjoram, wild marjoram or oregano and pot marjoram); nasturtium; parsley (dried); pennyroyal; rosemary; rue; sage;</p>	<p>0.19 L/ha</p>	<p>Ground application only</p>	<p>30 days</p>

2024-1378

2024-06-04

summer and winter savory; sweet bay (bay leaf); tansy; tarragon; thyme; wintergreen; woodruff; and wormwood 1,2,4,5			
Anise (anise seed); caraway (black); celery seed; culantro (seed); cumin; fennel [common and Florence (seed)]; juniper berry; lovage (seed); mustard (seed); poppy (seed); and Saffron. 1,2,4,5	0.19 L/ha	Ground application only	60 days
Cherry (sweet and tart) ^{1, 5, 8} (established trees)	0.38 L/ha	Ground application only	14 days
Buckwheat ^{1,5,7}	0.19 L/ha	Ground and aerial application	60 days
Highbush Blueberry ^{1,4,5}	0.38 L/ha	Ground application only	14 days
Canola	0.38 L/ha	Ground and aerial application	60 days
Carrot, Celeriac, garden (table) beet, Parsnip Radish, Rutabaga ^{3,5}	0.38 L/ha	Ground application only	30 days
Cranberry ^{1,2,5}	0.38 L/ha	Ground application only	30 days
Chive, Celery, Chinese Cabbage (napa) ^{1,5}	0.38 L/ha	Ground application only	30 days
Dill ^{1,5}	0.38 L/ha	Ground application only	30 days (dillweed leaves) 40 days (dill seed)
Flax (including low linolenic acid varieties)	0.38 L/ha	Ground and aerial application	60 days
Fenugreek ^{1,3,5}	0.38 L/ha	Ground application only	30 days (leaf and seed)
Field Peas	0.38 L/ha	Ground and aerial application	75 days
Hops ^{1,4,5}	0.38 L/ha	Ground application only	21 days
Lentils	0.38 L/ha	Ground and aerial application	60 days

2024-1378

2024-06-04

Oriental (Brown) Mustard (condiment and oilseed types)	0.38 L/ha	Ground and aerial application	60 days
Yellow Mustard	0.38 L/ha	Ground and aerial application	60 days
<i>Brassica carinata</i>	0.38 L/ha	Ground and aerial application	60 days
Dry Bulb Onions, Dry Bulb Shallot and Garlic ^{1,5}	0.38 L/ha	Ground application only	45 days
Potatoes	0.38 L/ha	Ground and aerial application	60 days
Green Onion ^{1,5}	0.38 L/ha	Ground application only	14 days
Leek ^{1,5}	0.38 L/ha	Ground application only	14 days
Prairie Carnation ^{1,5,6}	0.19 L/ha	Ground application only	N/A
Safflower ^{1,5}	0.38 L/ha	Ground application only	70 days
Soybeans	0.38 L/ha	Ground and aerial application	75 days
Spinach ^{3,5}	0.19 L/ha	Ground application only	14 days
Sunflowers	0.38 L/ha	Ground and aerial application	72 days

1. Apply a maximum of one application per year.
2. NOTE: Varieties of this crop or crops in this crop group may vary in their tolerance to herbicides, including **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE**. Since not all crops or crop varieties have been tested for tolerance to **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE**, first use of **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** should be limited to a small area of each crop or variety to confirm tolerance prior to adoption as a general field practice. Additionally, consult your supplier for information on the tolerance of specific varieties to **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE**.
3. Apply a maximum of two applications per year. If repeat application is required, allow at least 14 days between first and second application.
4. Apply **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** as a broadcast spray directed to the ground.
5. Refer to section "MINOR USES" in this label for additional use instructions.
6. Do not harvest treated prairie carnations for human consumption.
7. SPRAY VOLUME FOR AERIAL APPLICATION: Apply in 28 to 55 L/ha. Increase spray volumes up to 100L/ha as grass or crop foliage becomes dense.

2024-1378

2024-06-04

8. Do not apply **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** directly over the top of the sweet and tart cherry trees. Direct the spray at the base of the tree where grassy weeds are growing. If **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** is improperly applied, crop injury can occur.

TANK-MIXES:

When tank-mixes are permitted, read and observe all label directions, including rates and restrictions for each product used in the tank-mix. Follow the more stringent label precautionary measures for mixing, loading and applying stated on both product labels.

SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE AND A HERBICIDE CONTAINING ONLY BROMOXYNIL AND MCPA (E.G., BUCTRIL® M):

Flax (including low linolenic acid varieties): **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** may be tank-mixed with a herbicide containing only bromoxynil and MCPA for control of certain broadleaf and grassy weeds in flax (including low linolenic acid varieties) in one spray operation. Apply in 55 to 110 litres of water per hectare at a pressure of 275 kPa. DO NOT spray this tank-mix on plants under stress. DO NOT spray during periods of hot, humid weather. Follow all precautions, limitations and timing recommendations on the bromoxynil + MCPA herbicide product label.

Time of Application: Flax (including low linolenic acid varieties) may be treated from 5 cm high to the early bud stage. For best results, apply when flax (including low linolenic acid varieties) is 5 to 10 cm high and weeds are in the seedling stage. If broadleaf weeds and grassy weeds are not in the recommended treatment stage at the same time, then separate applications should be made.

Application Rate: **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** at 0.19 to 0.38 L/ha tank-mixed with a bromoxynil + MCPA herbicide product at 280 g ae/ha + 280 g ae/ha and AMIGO Adjuvant at 0.5 to 1.0 % v/v or JOURNEY HSOC at 0.5% v/v.

Weeds Controlled: As listed for **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** used alone, plus certain seedling broadleaf weeds. Refer to the bromoxynil + MCPA herbicide product label for the broadleaf weeds controlled.

MIXING INSTRUCTIONS:

1. Thoroughly clean the sprayer by flushing the system with water containing detergent.
2. Fill clean spray tank half full with clean water. Start agitation system.
3. If tank-mixing **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** and a bromoxynil + MCPA herbicide product for once-over grass and broadleaf weed control in flax (including low linolenic acid-varieties), add the correct amount of the bromoxynil + MCPA herbicide product. Continue to agitate.
4. Add the correct amount of **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE**. Continue to agitate.

2024-1378

2024-06-04

5. Add the correct amount of AMIGO ADJUVANT or JOURNEY HSOC along with the remaining amount of water necessary to fill the spray tank.
6. Continue to agitate or run the by-pass system.
7. After any break in the spraying operation, agitate thoroughly before spraying again. Check inside the tank to ensure that sprayer agitation is sufficient to re-mix the spray materials. Do not allow the mixture to sit overnight.
8. Immediately after use, thoroughly clean the sprayer by flushing the system with clean water containing detergent.

For disposal of cleaning solution, see section on DISPOSAL.

SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE AND CURTAIL® M HERBICIDE:

Flax (including low linolenic acid varieties): SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE may be tank-mixed with CURTAIL M HERBICIDE for control of certain broadleaf and grass weeds in flax in one spray operation. Apply in a minimum of 100 L/ha spray volume at a pressure of 275 kPa.

Follow all precautions, limitations, restrictions and timing recommendations on the CURTAIL M HERBICIDE label.

Time of Application: Flax may be treated from 5 to 15 cm in height. Apply when the grassy weeds are in the 2 to 6 leaf stage and the broadleaf weeds are actively growing. Application to Canada thistle should be made when Canada thistle is between 10 cm and up to the pre-bud stage. If broadleaf weeds and grass weeds are not in recommended treatment stage at the same time, then separate applications should be made.

Application Rate: **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** at 0.19 L/ha tank-mixed with CURTAIL M HERBICIDE at 1.5 to 2.0 L/ha and with AMIGO ADJUVANT at 0.5 % v/v or JOURNEY HSOC at 0.5% v/v.

Weeds Controlled: Weed claims listed on the **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** label plus weed claims listed on Curtail M HERBICIDE label. Refer to the CURTAIL M HERBICIDE label for the broadleaf weeds controlled.

Grazing Interval

Do not cut treated crops for feed or graze until 60 days after application.

MIXING INSTRUCTIONS:

1. Thoroughly clean the sprayer by flushing the system with water containing detergent.
2. Fill clean spray tank half full with clean water. Start agitation system.
3. Add the required amount of CURTAIL M HERBICIDE to the tank. Continue to agitate.
4. Add the correct amount of **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE**. Continue to agitate.
5. Add the correct amount of AMIGO ADJUVANT or JOURNEY HSOC along with the remaining amount of water necessary to fill the spray tank.

2024-1378

2024-06-04

6. Continue to agitate or run the by-pass system.
7. After any break in the spraying operation, agitate thoroughly before spraying again. Check inside the tank to ensure that sprayer agitation is sufficient to re-mix the spray materials. Do not allow the mixture to sit overnight. In the case of tank mixtures with broadleaf herbicides, settling will occur if agitation is not continuous.
8. If an oil film starts to build up in the tank, drain it and clean tank with strong detergent solution.
9. Immediately after use, thoroughly clean the sprayer by flushing the system with clean water containing detergent.

SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE AND MCPA ESTER:

Flax: SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE may be tank-mixed with MCPA ESTER for control of certain broadleaf and grassy weeds in flax in one spray operation. For ground applications, apply in a minimum of 100 L/ha spray volume at a pressure of 275 kPa. For aerial application apply this tank-mix in a minimum of 30 L/ha spray volume at a pressure of 235 kPa.

NOTE: THIS TANK-MIX MAY NOT BE APPLIED TO LOW LINOLENIC ACID VARIETIES OF FLAX.

Time of application: Flax may be treated from 5 cm tall to just before the bud stage. Do not apply after the early bud stage. Apply when the grassy weeds are in the 2 to 6 leaf stage and broad leaf weeds are in the seedling stage and actively growing. If broadleaf weeds and grassy weeds are not in the recommended treatment stage at the same time, then separate applications should be made.

Application Rate: SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE at 0.19 to 0.38 L/ha tank-mixed with MCPA ESTER and

AMIGO ADJUVANT at 0.5 to 1.0 % v/v or JOURNEY HSOC at 0.5% v/v.

Weeds Controlled: As listed for **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** used alone, plus certain seedling broadleaf weeds. Refer to the MCPA ESTER label for the rates and broadleaf weeds controlled and specific information on flax.

MIXING INSTRUCTIONS:

1. Thoroughly clean the sprayer by flushing the system with water containing detergent.
2. Fill clean spray tank half full with clean water. Start agitation system.
3. Add the correct amount of MCPA ESTER. Continue to agitate.
4. Add the correct amount of **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE**. Continue to agitate.
5. Add the correct amount of AMIGO ADJUVANT or JOURNEY HSOC along with the remaining amount of water necessary to fill the spray tank.
6. Continue to agitate or run the by-pass system.

2024-1378

2024-06-04

7. After any break in the spraying operation, agitate thoroughly before spraying again. Check inside the tank to ensure that sprayer agitation is sufficient to re-mix the spray materials. Do not allow the mixture to sit overnight.
8. Immediately after use, thoroughly clean the sprayer by flushing the system with clean water containing detergent.

SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE AND LONTREL®:

Flax (including low linolenic acid varieties): SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE may be tank-mixed with LONTREL 360 HERBICIDE, LONTREL XC HERBICIDE, or LONTREL 72 HERBICIDE for control of certain broadleaf and grass weeds in flax in one spray operation. Apply in a minimum of 100 L/ha spray volume at a pressure of 275 kPa.

Follow all precautions, limitations and timing recommendations on the LONTREL 360 HERBICIDE, LONTREL XC HERBICIDE or LONTREL 72 HERBICIDE label.

Time of Application: Flax may be treated from 5 to 10 cm in height. Apply when the grassy weeds are in the 2 to 6 leaf stage and the broadleaf weeds are actively growing. Application to Canada thistle should be made when Canada thistle is in the rosette to pre-bud stage of growth. If broadleaf weeds and grass weeds are not in recommended treatment stage at the same time, then separate applications should be made.

Application Rate: SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE at 0.19 L/ha tank-mixed with LONTREL 360 HERBICIDE at 0.56 to 0.83 L/ha, or LONTREL XC HERBICIDE at 0.34 L/ha to 0.5 L/ha or LONTREL 72 HERBICIDE at 280 g/ha to 415 g/ha and with AMIGO ADJUVANT at 0.5 % v/v or JOURNEY HSOC at 0.5% v/v.

Weeds Controlled: Wild oats, green foxtail, volunteer barley, volunteer wheat, volunteer oats, wild buckwheat.

Canada thistle	200 g ae/ha clopyralid	Season-long control
	300 g ae/ha clopyralid	Control extended into following year.

MIXING INSTRUCTIONS: For mixing instructions, refer to **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** and LONTREL 360 HERBICIDE, LONTREL XC HERBICIDE or LONTREL 72 HERBICIDE tank-mix on canola.

Canola: SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE may be tank-mixed with LONTREL 360 HERBICIDE, LONTREL XC HERBICIDE or LONTREL 72 HERBICIDE for control of certain broadleaf and grass weeds in canola in one spray operation. Apply in a minimum of 100 L/ha spray volume at a pressure of 275 kPa. Follow all precautions, limitations and timing recommendations on the LONTREL 360 HERBICIDE, LONTREL XC HERBICIDE or LONTREL 72 HERBICIDE label.

Time of Application: Canola may be treated from the 2 to 6 leaf stage. Apply when the grassy weeds are in the 2 to 6 leaf stage and the broadleaf weeds are actively growing. Application to Canada thistle should be made when Canada thistle is in the rosette to pre-bud stage of growth. If broadleaf weeds and grass weeds are not in recommended treatment stage at the same time, then separate applications should be made.

2024-1378

2024-06-04

Application Rate: SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE at 0.19 L/ha tank-mixed with LONTREL 360 HERBICIDE at 0.42 to 0.83 L/ha or LONTREL XC HERBICIDE at 0.25 L/ha to 0.5 L/ha or LONTREL 72 HERBICIDE at 210 g/ha to 415 g/ha and with AMIGO ADJUVANT at 0.5 % v/v or JOURNEY HSOC at 0.5% v/v.

2024-1378

2024-06-04

Weeds Controlled: Wild oats, green foxtail, volunteer barley, volunteer wheat, volunteer oats.

Wild Buckwheat	200 g ae/ha clopyralid	Season-long control
Canada thistle	150 g ae/ha clopyralid	Top growth control to 6 to 8 weeks
	200 g ae/ha clopyralid	Season-long control
	300 g ae/ha clopyralid	Control into following year.

MIXING INSTRUCTIONS:

1. Thoroughly clean the sprayer by flushing the system with water containing detergent.
2. Fill clean spray tank half full with clean water. Start agitation system.
3. Add the required amount of LONTREL 360 HERBICIDE, LONTREL XC HERBICIDE or LONTREL 72 HERBICIDE to the tank. Continue to agitate.
4. Add the correct amount of **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE**. Continue to agitate.
5. Add the correct amount of AMIGO ADJUVANT or JOURNEY HSOC along with the remaining amount of water necessary to fill the spray tank.
6. Continue to agitate or run the by-pass system.
7. After any break in the spraying operation, agitate thoroughly before spraying again. Check inside the tank to ensure that sprayer agitation is sufficient to remix the spray materials. Do not allow the mixture to sit overnight. In the case of tank mixtures with broadleaf herbicides, settling will occur if agitation is not continuous.
8. If an oil film starts to build up in the tank, drain it and clean tank with strong detergent solution.
9. Immediately after use, thoroughly clean the sprayer by flushing the system with clean water containing detergent.

SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE AND A HERBICIDE CONTAINING ONLY IMAZETHAPYR (E.G., PURSUIT 240):

Field Peas: **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** may be tank-mixed with a herbicide containing only imazethapyr for control of certain broadleaf and grassy weeds in field peas in one spray operation. Apply in a minimum of 100 litres of water per hectare at a spray pressure of 275 kPa. Follow all precautions, limitations and timing recommendations on the imazethapyr herbicide product label.

Time of Application: Field peas may be treated up to the sixth (6th) trifoliate leaf stage. Apply when volunteer wheat, volunteer barley, wild oats and green foxtail are in the 2 to 6 leaf stage. Apply up to the 4 leaf stage for control of broadleaf weeds. If broadleaf weeds and grassy weeds are not in the recommended treatment stage at the same time, then separate applications should be made.

Application Rate: **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE at 0.19 L/ha tank-mixed with an imazethapyr herbicide product at 50.4 g ai/ha and AMIGO Adjuvant at 0.5 % v/v or JOURNEY HSOC at 0.5% v/v.

2024-1378

2024-06-04

Weeds Controlled: Wild oats, green foxtail, volunteer barley, volunteer wheat plus the weeds as listed on the imazethapyr herbicide product label.

MIXING INSTRUCTIONS:

1. Thoroughly clean the sprayer by flushing the system with water containing detergent.
2. Fill clean spray tank half full with clean water. Start agitation system.
3. If tank-mixing **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** and an imazethapyr herbicide product for once over grass and broadleaf weed control in field peas, add the correct amount of the imazethapyr herbicide product. Continue to agitate.
4. Add the required amount of **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** to the tank. Continue to agitate.
5. Add the correct amount of AMIGO ADJUVANT or JOURNEY HSOC along with the remaining amount of water necessary to fill the spray tank.
6. Continue to agitate or run the by-pass system.
7. After any break in the spraying operation, agitate thoroughly before spraying again. Check inside the tank to ensure that sprayer agitation is sufficient to re-mix the spray materials. Do not allow the mixture to sit overnight.
8. Immediately after use, thoroughly clean the sprayer by flushing the system with clean water containing detergent.

SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE AND MUSTER®:

Canola: **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** may be tank-mixed with MUSTER for control of certain broadleaf and grassy weeds in canola in one spray operation. Apply in a minimum of 100 litres of water per hectare at a spray pressure of 275 kPa. Follow all precautions, limitations and timing recommendations on the MUSTER label.

Time of Application: Canola may be treated from the 2 leaf stage to the beginning of bolting. Apply when the grassy weeds are in the 2 to 6 leaf stage and when the broadleaf weeds are in the cotyledon to 6 leaf stage. If broadleaf weeds and grassy weeds are not in the recommended treatment stage at the same time, then separate applications should be made.

Application Rate: **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** at 0.19 L/ha tank-mixed with MUSTER at 20 or 30 g/ha and AMIGO ADJUVANT at 0.5 % v/v or JOURNEY HSOC at 0.5% v/v.

Weeds Controlled:

Muster at 20 or 30 g/ha: Wild oats, green foxtail, volunteer barley, volunteer oats, volunteer wheat, hempnettle, wild mustard, green smartweed, flixweed (spring seedlings), stinkweed*, and redroot pigweed.

* *Stinkweed suppression at the lower rate. Apply to stinkweed in the 1 to 4 leaf stage.*

MIXING INSTRUCTIONS:

1. Thoroughly clean the sprayer by flushing the system with water containing

2024-1378

2024-06-04

detergent.

2. Fill clean spray tank half full with clean water. Start agitation system.
3. Add the required amount of MUSTER to the tank. Continue to agitate.
4. Once MUSTER is in suspension, add the required amount of **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE**. Continue to agitate.
5. Add the correct amount of AMIGO ADJUVANT or JOURNEY HSOC along with the remaining amount of water necessary to fill the spray tank.
6. Continue to agitate or run the by-pass system.
7. On repeat tankloads, prepare a MUSTER/water slurry and add to required water volume in the spray tank prior to adding **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** and AMIGO ADJUVANT or JOURNEY HSOC.
8. After any break in the spraying operation, agitate thoroughly before spraying again. Check inside the tank to ensure that sprayer agitation is sufficient to re-mix the spray materials. Do not allow the mixture to sit overnight.
9. Immediately after use, refer to MUSTER label for appropriate tank cleanout instructions.

SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE AND LIBERTY® 150 SN HERBICIDE or INTERLINE® HERBICIDE – PRAIRIE PROVINCES AND INTERIOR OF BRITISH COLUMBIA (INCLUDING PEACE RIVER REGION) ONLY:

Canola: For use on Glufosinate Ammonium tolerant canola varieties only (e.g. varieties or hybrids labelled LIBERTYLINK).

SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE may be tank-mixed with LIBERTY 150 SN Herbicide or INTERLINE Herbicide for control of certain broadleaf and grass weeds in glufosinate ammonium tolerant canola varieties in one spray operation. For ground applications, apply in a minimum of 110 L/ha spray volume at a pressure of 275 kPa. For aerial applications apply this tank-mix in a minimum of 55 L/ha spray volume at a pressure of 300 kPa.

Follow all precautions, limitations and timing recommendations on the LIBERTY 150 SN Herbicide or INTERLINE Herbicide label.

2024-1378
2024-06-04

RECOMMENDED APPLICATIONS:

Products	Rate / ha	Weeds Controlled	Time of Application
SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE + LIBERTY 150 SN Herbicide or INTERLINE Herbicide + AMIGO Adjuvant or JOURNEY HSOC	63 mL/ha + 2.67 or 4.0 L/ha + 0.5% v/v	Wild oat, green foxtail, volunteer barley, volunteer wheat, volunteer oat plus all remaining grass weeds and broadleaf weeds listed on the LIBERTY 150 SN Herbicide or INTERLINE Herbicide label at the application rates of 2.67 and 4.0 L/ha.	Glufosinate ammonium tolerant canola varieties may be treated from the cotyledon stage up to the early bolting stage. Apply when the wild oat, volunteer wheat, volunteer oat and green foxtail are in the 1- to 5-leaf stage. Refer to the LIBERTY 150 SN Herbicide or INTERLINE Herbicide label for application timing on all remaining grass and broadleaf weeds for 2.67 or 4.0 L/ha.
SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE + LIBERTY 150 SN Herbicide or INTERLINE Herbicide + AMIGO Adjuvant or JOURNEY HSOC	190 mL/ha + 3.33 or 4.0 L/ha + 0.5 % v/v	Weed species listed on the SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE and LIBERTY 150 SN Herbicide or INTERLINE Herbicide labels plus foxtail barley, spring-germinated Japanese brome, spring-germinated downy brome.	Glufosinate ammonium tolerant canola varieties may be treated from the cotyledon stage up to the early bolting stage. Apply when the weeds are at growth stages listed on the SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE and LIBERTY 150 SN Herbicide or INTERLINE Herbicide labels, foxtail barley is at the 1- to 4-leaf stage with a maximum of 2 tillers and spring germinated Japanese and downy brome are at the 1- to 6-leaf stage up to the emergence of the 1 st tiller stage.

2024-1378

2024-06-04

Facet L + SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE + LIBERTY 150 SN Herbicide or INTERLINE Herbicide + AMIGO Adjuvant or JOURNEY HSOC	0.28 L/ha + 63, 125, or 190 mL/ha + 3.33 L/ha + 0.5 % v/v	For enhanced and more consistent control of cleavers and annual grasses	Apply in 100 L/ha of water. Application should be made from the 2 to 6 true leaf stage of the canola crop.
--	---	---	--

MIXING INSTRUCTIONS:

1. Thoroughly clean the sprayer by flushing the system with water containing detergent.
2. Fill clean spray tank half full with clean water. Start agitation system.
3. Add the required amount of FACET L and then AMIGO ADJUVANT or JOURNEY HSOC to the tank. Continue to agitate until thoroughly mixed.
4. STOP agitation. Add the correct amount of LIBERTY 150 SN Herbicide or INTERLINE HERBICIDE to the spray tank. Start agitation system.
5. Add the correct amount of **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** along with the remaining amount of water necessary to fill the spray tank.
6. Continue to agitate or run the by-pass system and spray out immediately.
7. After any break in the spraying operation, agitate thoroughly before spraying again. Check inside the tank to ensure that sprayer agitation is sufficient to re-mix the spray materials. Do not allow the mixture to sit overnight. In the case of tank mixtures with broadleaf herbicides, settling will occur if agitation is not continuous.
8. If an oil film starts to build up in the tank, drain it and clean tank with strong detergent solution.
9. Immediately after use, thoroughly clean the sprayer by flushing the system with clean water containing detergent.

SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE AND LIBERTY® 200 SN HERBICIDE– EASTERN CANADA AND BRITISH COLUMBIA ONLY

Canola: For use on glufosinate ammonium tolerant canola varieties only (e.g. varieties or hybrids labelled LIBERTYLINK).

2024-1378

2024-06-04

SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE may be tank-mixed with LIBERTY 200 SN Herbicide for control of certain broadleaf and grass weeds in glufosinate ammonium tolerant canola varieties in one spray operation. Apply in a minimum of 110 L/ha spray volume at a pressure of 275 kPa. Follow all precautions, limitations and timing recommendations on the LIBERTY 200 SN Herbicide label.

Time of Application: Glufosinate ammonium tolerant canola varieties may be treated from the cotyledon stage up to the early bolting stage. Apply when the wild oats, volunteer wheat, volunteer oats and green foxtail are in the 1 – 5 leaf stage. Refer to the LIBERTY 200 SN Herbicide label for application timing on all remaining grass and broadleaf weeds for 2.0 or 2.5 L/ha.

Application Rate: **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** at 63 mL/ha tank-mixed with LIBERTY 200 SN Herbicide at 2.0 to 2.5 L/ha and with AMIGO ADJUVANT at 0.5% v/v or JOURNEY HSOC at 0.5% v/v.

Weeds Controlled: Wild oats, green foxtail, volunteer barley, volunteer wheat, volunteer oats plus all remaining grass weeds and broadleaf weeds listed on the LIBERTY 200 SN Herbicide label at the application rates of 2.0 and 2.5 L/ha.

MIXING INSTRUCTIONS:

1. Thoroughly clean the sprayer by flushing the system with water containing detergent.
2. Fill clean spray tank half full with clean water. Start agitation system.
3. Add the required amount of AMIGO ADJUVANT or JOURNEY HSOC to the tank. Continue to agitate until thoroughly mixed.
4. STOP agitation. Add the correct amount of **LIBERTY 200 SN Herbicide** to the spray tank. Start agitation system.
5. Add the correct amount of **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** along with the remaining amount of water necessary to fill the spray tank.
6. Continue to agitate or run the by-pass system and spray out immediately.
7. After any break in the spraying operation, agitate thoroughly before spraying again. Check inside the tank to ensure that sprayer agitation is sufficient to re-mix the spray materials. Do not allow the mixture to sit overnight. In the case of tank mixtures with broadleaf herbicides, settling will occur if agitation is not continuous.
8. If an oil film starts to build up in the tank, drain it and clean tank with strong detergent solution.
9. Immediately after use, thoroughly clean the sprayer by flushing the system with clean water containing detergent.

GLYPHOSATE TOLERANT SOYBEANS

SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE may be tank-mixed with Roundup-Weathermax (or other liquid glyphosate products registered for use on glyphosate tolerant soybeans) for control of volunteer glyphosate tolerant corn and other weeds listed on the product labels.

All weeds listed on each label can be controlled in one spray operation.

2024-1378

2024-06-04

Apply in a minimum of 100 L/ha spray volume at a pressure of 275 kPa. Follow all precautions, limitations, restrictions and timing recommendations on the glyphosate brand labels.

Time of application: Glyphosate tolerant soybeans may be treated from 1st trifoliolate through to flowering, observing the 75-day PHI. Apply when grass weeds are in the 2- to 6-leaf stage and the broadleaf weeds are actively growing.

Application Rate: SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE at 0.19 L/ha tank-mixed with 900 to 1800 g ae/ha of a liquid glyphosate product that is registered for use on glyphosate tolerant soybeans (equivalent to 1.67 - 3.33 L/ha of Roundup Weathermax, a 540 g/L product, plus AMIGO Adjuvant at 0.5% v/v or JOURNEY HSOC at 0.5% v/v.

Weeds Controlled: Weed claims listed on the **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** label plus weed claims listed on the glyphosate product label. Refer to the glyphosate product label for the broadleaf weeds controlled.

Grazing Interval: Do not cut treated crops for feed or graze until 60 days after application.

MIXING INSTRUCTIONS:

1. Thoroughly clean the sprayer by flushing the system with water containing detergent.
2. Fill clean spray tank half full with clean water. Start agitation system.
3. Add the correct amount of **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE**. Continue to agitate.
4. Add the correct amount of AMIGO Adjuvant or JOURNEY HSOC. Continue to agitate.
5. Add the required amount of glyphosate brand to the tank along with the remaining amount of water necessary to fill the spray tank.
6. Continue to agitate or run the by-pass system.
7. After any break in the spraying operation, agitate thoroughly before spraying again. Check inside the tank to ensure that sprayer agitation is sufficient to re-mix the spray materials. Do not allow the mixture to sit overnight. In the case of tank-mixtures with broadleaf herbicides, settling will occur if agitation is not continuous.
8. If an oil film starts to build up in the tank, drain it and clean tank with strong detergent solution.
9. Immediately after use, thoroughly clean the sprayer by flushing the system with clean water containing detergent.

MINOR USES:

NOTE TO USER:

READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR THE INDICATED SPECIAL USE APPLICATIONS (LISTED BELOW):

The DIRECTIONS FOR USE for the uses described in this section of the label were developed by persons other than UPL AgroSolutions Canada, Inc. (UPL) under the User Requested Minor Use

2024-1378

2024-06-04

<p>Label Expansion program. For these uses, UPL AgroSolutions Canada, Inc. (UPL) has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.</p>					
CROP	Weeds Controlled	Application Rate of: ¹			Remarks
		SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE	AMIGO ADJUVANT	JOURNEY HSOC	
PRAIRIE CARNATION	Annual weeds listed on SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE label. Refer to weed/rate chart in this label.	0.19 L/ha	0.5 % v/v	0.5 % v/v	<p>TIME OF APPLICATION: Apply SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE post-emergence of weeds and crop.</p> <p>Apply a maximum of one application per year using ground equipment.</p> <p>Apply in a minimum spray volume of 110 L/ha.</p> <p>Do not harvest treated prairie carnations for human consumption.</p> <p>Ground application only.</p>
Edamame, Edible-podded legume vegetables (Crop subgroup 6A): Bean (<i>Phaseolus</i> spp.) (includes runner bean, snap bean, wax bean), Bean (<i>Vigna</i> spp.) (includes asparagus	Labelled grassy weeds. Refer to weed/rate chart in this label.	0.19 L/ha	0.5 % v/v	0.5 % v/v	<p>TIME OF APPLICATION: Apply SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE postemergence of weeds and crop.</p> <p>Apply a maximum of one application per year using ground equipment.</p> <p>Apply in a minimum spray volume of 110 L/ha.</p>

2024-1378

2024-06-04

<p>bean, Chinese longbean, moth bean, yardlong bean), Jackbean (<i>Canavalia ensiformis</i>), Pea (<i>Pisum</i> spp.) (includes dwarf pea, edible-pod pea, snow pea, sugar snap pea), Pigeon pea (<i>Cajanus cajan</i>), Soybean (immature seed) (<i>Glycine max</i>), Sword bean (<i>Canavalia gladiata</i>).</p> <p>Succulent shelled peas and beans (Crop subgroup 6B): Bean (<i>Phaseolus</i> spp.) (includes lima bean, green bean), Bean (<i>Vigna</i> spp.) (includes blackeyed pea, cowpea, southern pea), Broad bean (fava bean) (<i>Vicia faba</i>), Pea (<i>Pisum</i> spp.) (includes English pea, garden pea, green pea), Pigeon pea (<i>Cajanus cajan</i>).</p>					<p>Observe a PHI of 21 days.</p> <p>Ground application only.</p>
<p>BUCKWHEAT¹</p>	<p>Labelled grassy weeds. Refer to weed/rate chart in this label.</p>	<p>0.19 L/ha</p>	<p>0.5% v/v</p>	<p>0.5% v/v</p>	<p>TIME OF APPLICATION: Apply SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE</p>

2024-1378
 2024-06-04

					<p>postemergence of weeds and crop.</p> <p>Apply a maximum of one application per year using ground or aerial equipment.</p> <p>Apply in a minimum spray volume of 100 L/ha for ground application. Apply in a minimum spray volume of 28-55 for aerial application. Increase to 100 L/ha if grass or crop foliage becomes dense.</p> <p>Observe a PHI of 60 days.</p>
<p>CHERRY (SWEET AND TART)</p> <p>(established trees)</p>	<p>Labelled grassy weeds. Refer to weed/rate chart in this label.</p>	<p>0.19 – 0.38 L/ha</p>	<p>0.5 – 1.0 % v/v</p>	<p>0.5 % v/v</p>	<p>TIME OF APPLICATION: Apply SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE to established sweet and tart cherry trees, post-emergence of weeds. Do not apply SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE directly over the top of the trees. Direct the spray at the base of the tree where grassy weeds are growing. If SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE is improperly applied, crop injury can occur.</p> <p>Apply a maximum of one application per year using ground equipment.</p> <p>Apply in a minimum spray volume of 100 litres of water per hectare.</p> <p>Observe a PHI of 14 days.</p> <p>Ground application only.</p>

2024-1378
2024-06-04

<p>HIGHBUSH BLUEBERRY</p>	<p>SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE labelled grassy weeds. Refer to weed/rate chart in this label.</p>	<p>0.19 - 0.38 L/ha</p>	<p>0.5 - 1.0 % v/v</p>	<p>0.5 % v/v</p>	<p>TIME OF APPLICATION: Apply SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE post-emergence of weeds and crop.</p> <p>Apply a maximum of one application per year, post-emergent by ground equipment.</p> <p>Apply as a broadcast spray directed to the ground.</p> <p>Apply in a minimum spray volume of 100 L/ha.</p> <p>Observe a PHI of 14 days.</p> <p>Ground application only.</p>
<p>CORIANDER, (Cilantro), CARAWAY</p>	<p>Labelled grassy weeds. Refer to weed/rate chart in this label</p>	<p>0.19 - 0.38 L/ha Refer to SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE + AMIGO ADJUVANT or JOURNEY HSOC weed/rate chart in this label.</p>	<p>0.5 - 1.0% v/v Refer to SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE + AMIGO ADJUVANT weed/rate chart in this label.</p>	<p>0.5 % v/v Refer to SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE + JOURNEY HSOC weed/rate chart in this label.</p>	<p>TIME OF APPLICATION: Apply SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE post-emergent when coriander is in the 2 to 5 leaf stage.</p> <p>Apply a maximum of one application per year, using ground equipment.</p> <p>Observe a PHI of 30 days for leaves and 60 days for seed.</p> <p>Ground application only.</p>
<p>DILL</p>	<p>Labelled grassy weeds. Refer to weed/rate chart on this label.</p>	<p>0.19 - 0.38 L/ha</p>	<p>0.5 - 1.0% v/v</p>	<p>0.5 % v/v</p>	<p>TIME OF APPLICATION: Apply SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE post-emergence of weeds and crop when dill is in the 3-5 leaf stage. Apply a maximum of one application per year using</p>

2024-1378
2024-06-04

					ground equipment. Apply a minimum spray volume of 100 L/ha. Observe a PHI of 30 days for dillweed leaves and a PHI of 40 days for dill seed. Ground application only.
CHIVE	Labelled grassy weeds. Refer to weed/rate chart in this label	0.19 – 0.38 L/ha Refer to SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE + AMIGO ADJUVANT or JOURNEY HSOC weed/rate chart in this label.	0.5 – 1.0 % v/v Refer to SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE + AMIGO ADJUVANT weed/rate chart in this label.	0.5 % v/v Refer to SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE + JOURNEY HSOC weed/rate chart in this label.	TIME OF APPLICATION: Apply a maximum of one application per year, post-emergent using ground equipment. Apply in a minimum spray volume of 110 L/ha. Observe a PHI of 30 days. Ground application only.
Angelica; balm (lemon balm); basil; borage; burnet; chamomile; catnip; chervil; clary; costmary; culantro (leaf); horehound; hyssop; lavender; lovage; marigold; marjoram (includes sweet or annual marjoram, wild marjoram or oregano and pot marjoram); nasturtium; parsley (dried); pennyroyal; rosemary; rue; sage; savory, summer and	Labelled grassy weeds. Refer to weed/rate chart in this label.	0.19 L/ha	0.5 % v/v	0.5 % v/v	TIME OF APPLICATION: Apply SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE post-emergence of weeds and crop. Apply a maximum of one application per year using ground equipment. Apply as a broadcast spray directed to the ground. Apply in a minimum spray volume of 110 L/ha. Observe a PHI of 30 days. Ground application only.

2024-1378

2024-06-04

winter; sweet bay (bay leaf); tansy; tarragon; thyme; wintergreen; woodruff; and wormwood					
Anise (anise seed), caraway (black); celery seed; culantro (seed); cumin; fennel [common and Florence (seed)]; juniper berry; lovage (seed); mustard (seed); poppy (seed); and saffron	Labelled grassy weeds. Refer to weed/rate chart in this label.	0.19 L/ha	0.5 % v/v	0.5 % v/v	<p>TIME OF APPLICATION: Apply SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE post-emergence of weeds and crop. Apply a maximum of one application per year using ground equipment. Apply as a broadcast spray directed to the ground. Apply in a minimum spray volume of 110 L/ha.</p> <p>Observe a PHI of 60 days.</p> <p>Ground application only.</p>
HOPS	Labelled grassy weeds. Refer to weed/rate chart in this label.	0.19 - 0.38 L/ha	0.5 -1.0% v/v	0.5 % v/v	<p>TIME OF APPLICATION: Apply SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE post-emergence of weeds and crop. Apply a maximum of one application per season, using ground equipment. Apply as a broadcast spray directed to the ground. Apply in a minimum spray volume of 110 litres of water per hectare. Observe a PHI of 21 days.</p> <p>Ground application only.</p>
FENUGREEK	Annual weeds listed on SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE label	Refer to SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE + AMIGO	Refer to SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE +	Refer to SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE + JOURNEY HSOC	<p>TIME OF APPLICATION: Apply SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE post-emergent when fenugreek is in the 3 to 5 leaf stage. Apply a maximum of one application per season, using ground equipment.</p>

2024-1378
2024-06-04

		ADJUVANT or JOURNEY HSOC weed/rate chart in this label.	AMIGO ADJUVA NT weed/rate chart in this label.	weed/rate chart in this label.	Apply in a minimum spray volume of 100L/ha. Observe a PHI of 30 days for leaves and seed. Ground application only.
DRY BULB ONIONS, DRY BULB SHALLOT and GARLIC	Labelled grassy weeds. Refer to weed/rate chart in this label.	0.19-0.38 L/ha	0.5 % v/v	0.5 % v/v	TIME OF APPLICATION: Apply SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE post-emergent when the crop is in the 1 to 4 leaf stage. Apply a maximum of one application per season, using ground equipment. Observe a PHI of 45 days. Ground application only.
GREEN ONION	Labelled weeds. Refer to weed/rate chart in this label.	0.19-0.38 L/ha	0.5-1.0 % v/v		TIME OF APPLICATION: Apply SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE post-emergent when the crop is in the 2 to 3 leaf stage. Apply a maximum of one application per season, using ground equipment. Apply in a minimum spray volume of 110 L/ha. Observe a PHI of 14 days. Ground application only.
LEEK	Labelled weeds. Refer to weed/rate chart in this label	0.19-0.38 L/ha	0.5-1.0 % v/v		TIME OF APPLICATION: Apply SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE post-emergent at least 2 weeks after transplanting when crop is well established and has produced one new fully expanded leaf. Apply a maximum of one application per season,

2024-1378

2024-06-04


					<p>using ground equipment. Apply in a minimum spray volume of 110 L/ha. Observe a PHI of 14 days.</p> <p>Ground application only.</p>
SAFFLOWER	Annual weeds listed on SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE label	0.19 - 0.38 L/ha.	0.5 - 1.0% v/v.	0.5 % v/v	<p>TIME OF APPLICATION: Apply SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE post-emergent when safflower is in the 6-8 leaf stage. Apply a maximum of one application per season using ground equipment.</p> <p>Apply in a minimum spray volume of 100 litres of water per hectare.</p> <p>Observe a PHI of 70 days.</p> <p>Ground application only.</p>
SPINACH	Annual Weeds listed on SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE label	Refer to SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE + AMIGO ADJUVANT or JOURNEY HSOC weed/rate chart in this label.	Refer to SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE + AMIGO ADJUVANT weed/rate chart in this label.	Refer to SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE + JOURNEY HSOC weed/rate chart in this label.	<p>TIME OF APPLICATION: Apply a maximum of two applications per year with 14 days between applications, post emergent by ground application when grassy weeds are in the 2 to 6 leaf stage.</p> <p>Observe a PHI of 14 days.</p> <p>Ground application only.</p>
CRANBERRY	Labelled grassy weeds. Refer to weed/rate chart in this label.	0.19 - 0.38 L/ha	0.5 - 1.0% v/v	0.5 % v/v	<p>TIME OF APPLICATION: Apply SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE post-emergence of weeds and</p>

2024-1378
2024-06-04

					<p>crop. Do not apply between the hook stage and full fruit set. Apply a maximum of one application per year using ground equipment. Apply in a minimum spray volume of 110 L of water /ha.</p> <p>Observe a PHI of 30 days.</p>
<p>GARDEN (TABLE) BEET, PARSNIP, CARROT, RADISH, RUTABAGA, CELERIAC</p>	<p>Labelled grassy weeds. Refer to weed/rate chart in this label.</p>	<p>0.19 - 0.38 L/ha</p>	<p>0.5 - 1.0% v/v</p>	<p>0.5 % v/v</p>	<p>TIME OF APPLICATION: Apply SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE post-emergence of weeds and crop using ground equipment. Apply a maximum of two applications per year. If repeat application is required, allow at least 14 days between first and second application.</p> <p>Do not apply more than 0.38 L/ha (90 grams ai/ha) per crop season.</p> <p>Apply in a minimum spray volume of 110 L/ha.</p> <p>Observe a PHI of 30 days.</p> <p>Ground application only.</p>
<p>CELERY, CHINESE CABBAGE (NAPA) ¹</p>	<p>Labelled grassy weeds. Refer to weed/rate chart in this label.</p>	<p>0.19-0.38 L/ha</p>	<p>0.5-1.0% v/v</p>	<p>0.5 % v/v</p>	<p>TIME OF APPLICATION: Apply SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE post-emergence of weeds and crop. Apply a maximum of one application per year using ground equipment. Apply in a minimum spray volume of 110 L/ha.</p> <p>Observe a PHI of 30 days.</p> <p>Ground application only.</p>

2024-1378
2024-06-04

¹ Choose either AMIGO ADJUVANT or JOURNEY HSOC



2024-1378

2024-06-04

APPLICATION INSTRUCTIONS:

GROUND APPLICATION

Water volume and spray pressure:

For ground application, use sprayers equipped with standard flat fan nozzles. For optimum spray coverage, use nozzles tilted 45° forward.

The use of flood jet or hollow cone nozzles is not recommended because of uneven and inadequate spray coverage. Thorough and uniform spray coverage over the entire leaf area of the target weeds is necessary for optimum weed control.

Thoroughly clean all screens to prevent nozzle clogging, especially when spray volumes of 55 to 110 L/ha are used.

For ground applications, use spray volumes of 55 to 225 L/ha and a minimum of 275 kPa pressure. For applications to dense weed infestations or to dense crop canopies, use a minimum of 110 L/ha spray volume at pressures of 275 to 425 kPa. In any situation, ensure the proper rate of AMIGO ADJUVANT or JOURNEY HSOC is maintained as listed on the label under APPLICATION RATES OF **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE**.

For legume vegetables (succulent or dried) (except soybeans) and edamame, a minimum spray volume of 110 L/ha is recommended.

AERIAL APPLICATION

Water volume and spray pressure:

Apply **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** alone or in tank mixes (ONLY with the recommended tank mix partners that are registered for aerial use) in no less than 28 L/ha spray volume at a pressure no less than 300 kPa.

Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended on this label.

When applying **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** by aircraft, uniform coverage is essential. To avoid streaked, uneven or overlapped application, use appropriate marking devices.

Aerial Use Precautions

- Read and understand the entire label before opening this product. If you have questions, call UPL AgroSolutions Canada, Inc. 1-800-438-6071 or obtain technical advice from the distributor or your provincial agricultural representative.
- Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the National Aerial Pesticide Application Manual, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

2024-1378

2024-06-04

- Do not apply to terrain where there is a potential for surface run-off to enter aquatic systems.
- Do not apply to any body of water. Avoid drifting of spray onto any body of water or other non-target areas. Specified buffer zones should be observed.
- Do not apply during periods of dead calm or when wind velocity and direction pose a risk of spray drift.
- Do not apply when wind speed is greater than 16 km/h at a flying height at the site of application.
- Do not apply with spray droplets smaller than ASAE coarse classification.

Aerial Operator Precautions:

- Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.
- It is desirable that the pilots have communication capabilities at each treatment site at the time of application.
- The field crew and the mixer/loaders must wear goggles or face shield, rubber apron, chemical-resistant gloves, rubber boots, long sleeved shirt and long-legged pants.
- All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

Product Specific Precautions:

- Read and understand the entire label before opening this product. If you have questions, call the manufacturer at 1-800-438-6071 or obtain technical advice from the distributor or your provincial agricultural representative.

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty.

DO NOT apply when wind speed is greater than 16 km/h at flying height at the site of application.

DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) medium classification. To reduce drift caused by turbulent wingtip vortices, the nozzle distribution along the spray boom length **MUST NOT** exceed 65% of the wing- or rotorspan.

Buffer zones:

Field sprayer application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Boom height must be 60 cm or less above the crop or ground.

Aerial application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply when wind speed is greater than 16 km/h at flying height at the site of application. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. To

2024-1378
 2024-06-04

reduce drift caused by turbulent wingtip vortices, the nozzle distribution along the spray boom length MUST NOT exceed 65% of the wing- or rotorspan.

Spot treatments using hand-held equipment DO NOT require a buffer zone.

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands) and sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands).

Method of application	Crop	Buffer Zones (metres) Required for the Protection of:		
		Freshwater Habitat of Depths:		Terrestrial Habitat
		Less than 1 m	Greater than 1 m	
Field sprayer	Seedling alfalfa, highbush blueberry, cherry, cranberry, caraway, chive, coriander, dill, fenugreek, dry bulb onion, dry bulb shallot, safflower, canola, carrot, flax (including low linolenic varieties), field pea, hops, lentils, oriental (brown) mustard (condiment and oilseed types), yellow mustard, <i>Brassica carinata</i> , potato, parsnip, radish, garden (table) beet, soybean, sunflower, celery, Chinese cabbage (napa), celeriac	1	1	2
	Spinach, Edamame, Edible-podded Legume (crop subgroup 6A) vegetables, Succulent shelled peas and beans (crop subgroup 6B),	1	1	1

2024-1378

2024-06-04

	bean (<i>lupines</i> , <i>phaseolus</i> , and <i>vigna</i> spp.), broad bean, chickpea, guar, lablab bean, Prairie carnation, pigeon pea, various herbs and spices (except chive, coriander, and dill), buckwheat				
Aerial	Canola, flax (including low linolenic acid varieties), field peas, lentils, potatoes, Oriental (brown) mustard (condiment and oilseed types), yellow mustard, soybeans, sunflowers, <i>Brassica carinata</i> , dry bulb onions, garlic	Fixed wing	10	1	60
		Rotary wing	10	1	50
	Desi and Kabuli chickpeas, Bean (<i>lupines</i> , <i>phaseolus</i> , and <i>vigna</i> spp.), broad bean, chickpea, guar, lablab bean, pigeon pea, buckwheat	Fixed wing	4	1	30
		Rotary wing	1	1	30

For Cherry (sweet and tart):

Use of the following spray methods or equipment DO NOT require a buffer zone: hand-held or backpack sprayer and spot treatment, *inter-row hooded sprayer*, *low-clearance hooded or shielded sprayers that ensure spray drift does not come in contact with orchard crop fruit or foliage*.

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Buffer Zone Calculator the Pesticides portion of the Canada.ca web site.

RESTRICTIONS AND LIMITATIONS:

2024-1378

2024-06-04

1. Use only for crops listed on the label.
2. Thorough preplant tillage operations are required to fields where sod or forage grass crops may have grown in the previous year.
3. Take necessary precautions to avoid sprayer overlaps.
4. Rainfall within one hour of application may reduce the effectiveness of the spray.
5. Do not mix or apply **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** with any other additive, pesticide or with fertilizer except as specifically recommended on the label.
6. Allow 4 days between application of **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** and any other chemical not recommended as a tank-mix combination on the label.
7. Do not apply more than 0.38 L/ha (90 grams ai/ha) per season.
8. Do not use crops for grazing of livestock or green feed following application of **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** until after the appropriate interval as specified in the crop recommendations table has been observed. Do not cut treated crops for forage until the appropriate interval as specified in the crop recommendations table has been observed.
9. **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE may be applied by air for certain crops specified in this label.**
10. **DO NOT** enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

HERBICIDE RESISTANCE MANAGEMENT RECOMMENDATIONS:

For resistance management, **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** is a Group 1 herbicide. Any weed population may contain or develop plants naturally resistant to **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** and other Group 1 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of **SELECT EMULSIFIABLE CONCENTRATE POST-EMERGENCE HERBICIDE** or other Group 1 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.

2024-1378

2024-06-04

- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact UPL AgroSolutions Canada, Inc. 1-800-438-6071.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

UPL, the UPL logo, SELECT, AMIGO and INTERLINE are registered trademarks of a UPL Corporation Limited Group Company.

©2023 UPL Corporation Limited Group Company.

BUCTRIL is a registered trademark of Bayer.

LONTREL is a trademark and CURTAIL is a registered trademark of Corteva AgriScience.

MUSTER is a registered trademark of FMC Corporation.

PURSUIT and LIBERTY are registered trademarks of BASF.

FACET is a trademark of BASF SE, used with permission by BASF Canada Inc.