

2012-4280
2012-11-22

PRESCRIPTION TREATMENT brand
AVERT AF
Cockroach Gel Bait
COMMERCIAL

GUARANTEE:

Abamectin 0.05%

Contains diazolidinyl urea at 0.30%, propyl paraben at 0.03%, and methyl paraben at 0.11% as preservatives

REGISTRATION NO. 28978 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND BROCHURE BEFORE USING

IN CASE OF EMERGENCY ENDANGERING LIFE OR PROPERTY, INVOLVING THIS PRODUCT, CALL DAY
OR NIGHT 1-800-454-2673.

NET CONTENTS: 30 grams

BASF Canada Inc.
100 Milverton Drive, 5th Floor
Mississauga, ON L5R 4H1
Tel. No. 1-877-371-2273

PRESCRIPTION TREATMENT brand

AVERT AF

Cockroach Gel Bait
COMMERCIAL

GUARANTEE:

Abamectin..... 0.05%

Contains diazolidinyl urea at 0.30%, propyl paraben at 0.03%, and methyl paraben at 0.11% as preservatives

REGISTRATION NO. 28978 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND BROCHURE BEFORE USING

IN CASE OF EMERGENCY ENDANGERING LIFE OR PROPERTY, INVOLVING THIS PRODUCT, CALL DAY OR NIGHT 1-800-454-2673

NET CONTENTS: 30 grams

BASF Canada Inc.
100 Milverton Drive, 5th Floor
Mississauga, ON L5R 4H1
Tel. No. 1-877-371-2273

SEE BROCHURE INSIDE FOR DIRECTIONS FOR USE.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN. CAUTION: Wash hands and exposed skin before eating, drinking or smoking and after handling. Wash all contaminated clothing thoroughly before re-use. Contents under pressure. Do not puncture or incinerate container. Do not use or store near heat or open flame. Exposure to temperatures above 54° C may cause bursting. Do not treat directly into any electronic equipment such as radios, televisions, computers, etc. Do not treat where electrical short circuits might result. Do not use on or contaminate fruit, vegetables or other food or feed crops.

Do not apply to humans, animals, clothing or bedding. Do not apply to areas accessible to children. Do not contaminate feed or food products or food preparation surfaces, dishes, kitchen utensils and food containers. To prevent staining, exposed bait should be cleaned up immediately with a damp cloth or sponge.

FIRST AID

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 – 20 min. Remove contact lenses, if present, after the first 5 min, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 – 20 min. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Treat symptomatically

STORAGE: Store in a cool dry area away from heat or open flame and inaccessible to children.

DISPOSAL: Wastes resulting from the use of this product may be disposed of at an approved waste disposal facility. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill and for clean-up of spills.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on this label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

PRESCRIPTION TREATMENT brand

AVERT AF

Cockroach Gel Bait
COMMERCIAL

GUARANTEE:

Abamectin 0.05%

Contains diazolidinyl urea at 0.30%, propyl paraben at 0.03%, and methyl paraben at 0.11% as preservatives

REGISTRATION NO. 28978 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND BROCHURE BEFORE USING

IN CASE OF EMERGENCY ENDANGERING LIFE OR PROPERTY, INVOLVING THIS PRODUCT,
CALL DAY OR NIGHT 1-800-454-2673.

NET CONTENTS: 30 grams
BASF Canada Inc.
100 Milverton Drive, 5th Floor
Mississauga, ON L5R 4H1
Tel. No. 1-877-371-2273

DIRECTIONS FOR USE

CONTROLS:

German cockroaches (including carbamate, organophosphate, and pyrethroid resistant strains).

FOR USE IN AND AROUND:

Commercial, Industrial and Residential Buildings including Apartments, Garages, Food Storage Areas, Homes, Hospitals and Nursing Homes, Hotels, Meat Packing and Food Processing Plants, Motels, Restaurants and other Food Handling Establishments, Schools, Supermarkets, Transportation Equipment (Buses, Boats, Ships, Trains and Planes) and Warehouses.

GENERAL USE DIRECTIONS

Prescription Treatment brand Avert AF Cockroach Gel Bait is for use in controlling German cockroaches indoors. Use suitable equipment capable of application into cracks and crevices, void spaces and other locations where insects may be harbouring, living and breeding. Bait points may be made every 10 - 20 centimetres. Avoid placing baits on surfaces that are openly exposed such as the junction between wall and floor.

DO NOT APPLY TO AREAS ACCESSIBLE TO CHILDREN.

APPLICATION RATES - ALL USES

For light infestations of Cockroaches, 4 - 6 bait points are recommended per 10 m² (100 sq. ft.) of treatment area. Light infestations would be indicated by sightings only at night when a light is turned on. For heavy infestations of Cockroaches, 12 - 24 bait points are recommended per 10 m² (100 sq. ft.) of treatment area. Heavy infestations are indicated when cockroaches are readily seen upon the opening of cabinets, pantries or closets. Each bait point should equal about 0.25 to 0.50 grams of product. A spot approximately 4 mm (1/8 inch) in diameter equals 0.25 grams of product, while a spot approximately 6 mm (1/4 inch) in diameter equals 0.50 grams of product.

Baits should be reapplied when all bait is consumed or is no longer palatable, approximately every three months. Applications should not be made more than once a month, unless bait in that area is consumed.

APPLICATION IN ALL EXCEPT FOOD/FEED HANDLING AREAS OF FOOD/FEED HANDLING ESTABLISHMENTS

For treatment in residential and the non-food/non-feed areas of institutional, warehousing and commercial establishments, place bait into cracks and crevices; holes; pipe chases; undersides of furniture; under drain plates; in or under trash containers; hidden surfaces around sinks and storage areas; behind baseboards; around doors and windows; inside, behind and under cabinets, drawers and shelving; under and behind appliances such as stoves and refrigerators; and in attics and crawl spaces. Also apply in points between different elements of construction, between equipment and floors, openings leading to voids and hollow spaces in walls, equipment legs and bases and crawl spaces where insects hide. During follow-up visits, inspect bait placements and re-apply when bait has been consumed. Care should be taken to avoid depositing the product onto exposed surfaces. If gel contacts an exposed surface, remove gel and wash exposed surface. This product may also be used in food/feed areas of food/feed handling establishments as outlined below.

Non-food/feed areas are areas such as garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage areas (after canning or bottling).

APPLICATION IN FOOD/FEED AREAS OF FOOD/FEED HANDLING ESTABLISHMENTS

Application within food/feed areas of food/feed handling establishments is limited to Crack and Crevice treatment only. Apply Prescription Treatment brand Avert AF Cockroach Gel Bait directly into cracks and crevices. Apply small amounts of gel (up to 0.50 grams) per point. Do not apply bait to areas where food utensils or food processing surfaces may become contaminated. Do not apply bait to areas that are routinely washed such as cracks and crevices in tops of tables, food/feed preparation and prepared food holding surfaces as bait may be removed by washing. Care should be taken to avoid depositing gel onto exposed surfaces. If gel contacts an exposed surface, remove gel and wash exposed surface.

Food/feed areas include areas for receiving, storage, packing (canning, bottling, wrapping, boxing), preparing edible waste storage, and enclosed processing systems (mills, dairies, edible oils, syrups). Serving areas are also considered food/feed areas when food is exposed and facility is in operation. Apply bait in small amounts between different elements of construction, between equipment and floors, openings leading into voids and hollow spaces in floors, walls, ceilings, equipment legs and bases, around plumbing pipes, doors and windows, behind and under refrigerators, cabinets, sinks, closets, stoves and other equipment where cockroaches hide. During follow-up visits, inspect bait placements and re-apply when bait has been consumed.

Application of this product in food/feed areas of food/feed handling establishments other than as a crack and crevice treatment is not permitted. Do not treat food preparation surfaces.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN. CAUTION: Wash hands and exposed skin before eating, drinking or smoking and after handling. Wash all contaminated clothing thoroughly before re-use. Contents under pressure. Do not puncture or incinerate container. Do not use or store near heat or open flame. Exposure to temperatures above 54° C may cause bursting. Do not treat directly into any electronic equipment such as radios, televisions, computers, etc. Do not treat where electrical short circuits might result. Do not use on or contaminate fruit, vegetables or other food or feed crops.

Do not apply to humans, animals, clothing or bedding. Do not apply to areas accessible to children. Do not contaminate feed or food products or food preparation surfaces, dishes, kitchen utensils and food containers. To prevent staining, exposed bait should be cleaned up immediately with a damp cloth or sponge.

FIRST AID

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 – 20 min. Remove contact lenses, if present, after the first 5 min, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 – 20 min. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Treat symptomatically

STORAGE: Store in a cool dry area away from heat or open flame and inaccessible to children.

DISPOSAL: Wastes resulting from the use of this product may be disposed of at an approved waste disposal facility. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill and for clean-up of spills.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on this label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.