

2023-2066  
2023-10-24  
Proposed Label on 2023-05-08

**(Container)**

# Durango™ Herbicide

<b>GROUP</b>	<b>9</b>	<b>HERBICIDE</b>
--------------	----------	------------------

**Water soluble herbicide for nonselective weed control in CROPLAND SYSTEMS AND IN NON-CROPLAND AREAS.**

**AGRICULTURAL and INDUSTRIAL READ THE LABEL AND BOOKLET BEFORE USING**

**KEEP OUT OF REACH OF CHILDREN**

**ACTIVE INGREDIENT:** Glyphosate (present as dimethylamine salt) 480 g/L  
solution

**REGISTRATION NO. 29774 PEST CONTROL PRODUCTS ACT**  
**CAUTION: EYE AND SKIN IRRITANT**

**POTENTIAL SKIN SENSITIZER**

**Net Contents:** 7.5 L, 10 L, 115 L, 450 L and Bulk

Albaugh, LLC  
1525 NE 36th Street  
Ankeny, IA 50021  
USA  
1-800-247-8013

2023-2066  
2023-10-24  
Proposed Label on 2023-05-08

## **PRECAUTIONS**

May irritate eyes and skin  
Avoid contact with eyes or with skin

### **KEEP OUT OF REACH OF CHILDREN**

Wear long sleeved shirt, long pants and chemical resistant gloves during mixing, loading, application, clean up and repair. In addition, wear goggles or a face shield during mixing and loading.

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind directions, temperature inversions, application equipment and sprayer settings.

## **PHYSICAL OR CHEMICAL HAZARDS**

Spray solutions of this product should be mixed, stored and applied only in stainless steel, aluminum, fibreglass, plastic and plastic-lined steel containers. **DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS.** This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

## **FIRST AID**

**Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.**

**If swallowed:** Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. **DO NOT** induce vomiting unless told to do so by a poison control centre or doctor. **DO NOT** give anything by mouth to an unconscious person.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

## **TOXICOLOGICAL INFORMATION**

No specific antidote. Employ supportive care. Treatment should be based on judgment of the physician in response to reactions of the patient.

## **AGRICULTURAL CHEMICAL**

DO NOT ship or store with food, feeds, drugs or clothing.

## **ENVIRONMENTAL PRECAUTIONS**

Avoid direct applications to any body of water. **DO NOT** contaminate water by disposal of waste or cleaning of equipment. **TOXIC** to aquatic organisms and non-target terrestrial plants. Observe buffer zones specified under **DIRECTIONS FOR USE**. To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

## **STORAGE**

Avoid contamination of seed, feed, and foodstuffs.  
Soak up small amounts of spill with absorbent clays.

2023-2066

2023-10-24

Proposed Label on 2023-05-08

**DISPOSAL**

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

2023-2066

2023-10-24

Proposed Label on 2023-05-08

**NOTICE TO USER:** This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

2023-2066  
2023-10-24  
Proposed Label on 2023-05-08

**(Booklet)**



**Dow AgroSciences**

# Durango™ Herbicide

<b>GROUP</b>	<b>9</b>	<b>HERBICIDE</b>
--------------	----------	------------------

**Water soluble herbicide for nonselective weed control in CROPLAND SYSTEMS AND IN NON-CROPLAND AREAS.**

**AGRICULTURAL and INDUSTRIAL READ THE LABEL AND BOOKLET BEFORE USING**

**KEEP OUT OF REACH OF CHILDREN**

**ACTIVE INGREDIENT:** Glyphosate (present as dimethylamine salt) 480 g/L  
solution

**REGISTRATION NO. 29774 PEST CONTROL PRODUCTS ACT CAUTION: EYE AND SKIN IRRITANT**

**POTENTIAL SKIN SENSITIZER**

**Net Contents:** 7.5 L, 10 L, 115 L, 450 L and Bulk

Albaugh, LLC  
1525 NE 36th Street  
Ankeny, IA 50021  
USA  
1-800-247-8013

2023-2066  
2023-10-24  
Proposed Label on 2023-05-08

**TABLE OF CONTENTS**

Page (s)

PRECAUTIONS

DIRECTIONS FOR USE

GENERAL INFORMATION

MIXING AND APPLICATION

    Precautions

    Mixing and Application Equipment Information

WEEDS CONTROLLED

    Annual Weeds

    Perennial Weeds

    Woody Brush and Trees

CROPLAND USES

    ANNUAL WEED CONTROL

        Annual Weed Control with Durango Herbicide

2023-2066

2023-10-24

Proposed Label on 2023-05-08

Annual Weed Control with Durango Herbicide Tank Mixtures for Summerfallow & Minimum Tillage Systems

Addition of Surfactant

Additional Important Information For Annual Weed Control

Weed Control in Glyphosate Tolerant Canola

Weed Control in Glyphosate Tolerant Corn

Durango Herbicide plus Lontrel™ 360

Weed Control in Glyphosate Tolerant Soybean

#### PERENNIAL WEED CONTROL

Perennial Weed Control With Durango Herbicide

Special Notes for Perennial Weed Control:

Quackgrass

Surfactant Information

Canada Thistle

Toadflax

Dandelion

Alfalfa Control with 2,4-D Tank-Mix

All Perennial Weeds

#### CROPLAND SITUATIONS

Prior to Planting - All Crops

Post Harvest Stubble Treatment

Spot treatment In-Crop

Grazing Restrictions

Summerfallow Treatment

Minimum and Zero Tillage Cropping Systems

Minimum and Zero Tillage Tank Mixtures

Durango Herbicide plus Pardner

Durango Herbicide plus Pursuit

Forage Legumes and Grasses

Pasture Renovation

Forage Seed Production

Preharvest Control of Quackgrass, Canada Thistle, Milkweed, Toadflax and Dandelion

Timing of Preharvest Applications

Restricted Use – Aerial Preharvest Application

Tree Plantings

Tree, Vine and Berry Crops

Selective Equipment

#### NON-CROPLAND USES

Weed Control In Non-Cropland Areas with Durango Herbicide

Application Information for Non-Cropland Uses

Ground Applications

Purple Loosestrife Control

Selective Equipment for Non-Cropland Uses

Turfgrass

Injection Applications

Cut Stump Application

Resistance Management Recommendations

#### PRECAUTIONS

May irritate eyes and skin

Avoid contact with eyes or with skin

**KEEP OUT OF REACH OF CHILDREN**

2023-2066

2023-10-24

Proposed Label on 2023-05-08

Wear long sleeved shirt, long pants and chemical resistant gloves during mixing, loading, application, clean up and repair. In addition, wear goggles or a face shield during mixing and loading.

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind directions, temperature inversions, application equipment and sprayer settings.

### **PHYSICAL OR CHEMICAL HAZARDS**

Spray solutions of this product should be mixed, stored and applied only in stainless steel, aluminum, fibreglass, plastic and plastic-lined steel containers. **DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS.** This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

### **FIRST AID**

**Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.**

**If swallowed:** Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. **DO NOT** induce vomiting unless told to do so by a poison control centre or doctor. **DO NOT** give anything by mouth to an unconscious person.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

### **TOXICOLOGICAL INFORMATION**

No specific antidote. Employ supportive care. Treatment should be based on judgment of the physician in response to reactions of the patient.

### **AGRICULTURAL CHEMICAL**

**DO NOT** ship or store with food, feeds, drugs or clothing.

### **ENVIRONMENTAL PRECAUTIONS**

Avoid direct applications to any body of water. **DO NOT** contaminate water by disposal of waste or cleaning of equipment. **TOXIC** to aquatic organisms and non-target terrestrial plants. Observe buffer zones specified under **DIRECTIONS FOR USE**. To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

### **STORAGE**

Avoid contamination of seed, feed, and foodstuffs.

Soak up small amounts of spill with absorbent clays.

### **DISPOSAL**

#### **Recyclable Containers:**

**DO NOT** reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:



2023-2066  
2023-10-24  
Proposed Label on 2023-05-08

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

#### **Returnable Containers:**

DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

#### **Refillable Containers**

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. DO NOT reuse this container for any other purpose.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

### **DIRECTIONS FOR USE GENERAL INFORMATION**

The restricted entry interval is 12 hours after application for all agricultural uses.

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests. DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Glyphosate is not to be applied using hand-wicking or hand-daubing methods.

DO NOT apply this product using aerial spray equipment except under conditions as specified within this label.

Field sprayer application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) coarse classification. Boom height must be 60 cm or less above the crop or ground.

Airblast or mist blower application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** direct spray above plants to be treated. **DO NOT** apply when wind speed is greater than 16 km/h at the application site as measured outside of the treatment area on the upwind side. For airblast applications, turn off outward pointing nozzles at row ends and outer rows.

Aerial application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply when wind speed is greater than 16 km/h at flying height at the site of application. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) coarse classification. To reduce drift caused by turbulent wingtip vortices, the nozzle distribution along the spray boom length **MUST NOT** exceed 65% of the wing- or rotorspan.

Durango Herbicide, a water soluble liquid, mixes readily with water for application as a foliage spray for the control or destruction of most herbaceous plants. It may be applied through most standard industrial

2023-2066

2023-10-24

Proposed Label on 2023-05-08

or field type sprayers after dilution and thorough mixing with water in accordance with the booklet instructions. This herbicide moves through the plant from the point of foliage contact to and into the root system. Visible effects on most annual weeds occur within 2 to 4 days but on most perennial weeds may not occur until 7 to 10 days. Extremely cool or cloudy weather at treatment time may slow down activity of this product and delay visual effects of control. Visible effects are a gradual wilting and yellowing of the plant which advances to complete browning of above ground growth and deterioration of underground plant parts.

Delay application until vegetation has emerged to the stages described for control of such vegetation under the annual and perennial weed control sections of this booklet to provide adequate leaf surface to receive the spray. Unemerged plants arising from underground rhizomes or root stocks of perennials will not be affected by the spray and will continue to grow. For this reason best control of most perennial weeds is obtained when treatment is made at late growth stages approaching maturity.

Always use the higher rate of this product per hectare within the recommended range when weed growth is heavy or dense, or weeds are growing in an undisturbed (non-cultivated) area.

DO NOT treat weeds under poor growing conditions such as drought stress, disease or insect damage, as reduced weed control may result. Reduced results may also occur when treating weeds heavily covered with dust.

This product does not provide residual weed control. For subsequent residual weed control follow a label approved herbicide program. Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used.

Heavy rainfall immediately after application may wash the chemical off the foliage and a repeat treatment may be required. DO NOT apply if rainfall is forecast for the time of application.

DO NOT mix with any surfactant, pesticide, herbicide oils or any other material other than water unless specified in this booklet. For best results, spray coverage should be uniform and complete. DO NOT spray weed foliage to the point of runoff.

#### **MIXING AND APPLICATION PRECAUTIONS**

**ATTENTION: AVOID CONTACT WITH FOLIAGE, GREEN STEMS, OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT.**

**APPLY THESE SPRAY SOLUTIONS IN PROPERLY MAINTAINED AND CALIBRATED EQUIPMENT CAPABLE OF DELIVERING DESIRED VOLUMES.**

**AVOID DRIFT - EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURING DESIRABLE PLANTS AND CROPS.** Even minute quantities of spray drift can cause severe damage or destruction to nearby crops, plants or other areas on which treatment is not intended, or may cause other unintended consequences.

**DO NOT USE IN GREENHOUSES. REDUCED RESULTS MAY OCCUR IF WATER CONTAINING SOIL IS USED, SUCH AS WATER FROM PONDS AND UNLINED DITCHES.**

Clean sprayer and parts immediately after using this product by thoroughly flushing with water. DO NOT contaminate water sources by disposal of wastes or cleaning of equipment.

**NOTE:** Use of this product in any manner not consistent with this booklet may result in injury to persons, animals or crops, or other unintended consequences. Keep container closed to prevent spills and contamination.

2023-2066

2023-10-24

Proposed Label on 2023-05-08

## **MIXING AND APPLICATION EQUIPMENT INFORMATION MIXING**

For ground or industrial type sprayers, fill the spray tank with one-half the required amount of water. Add the proper amount of herbicide (see "Weed Control" sections of this booklet) and mix well before adding the remaining portion of water. Placing the filling hose below the surface of the liquid solution will prevent excessive foaming. Removing hose from tank immediately will avoid back siphoning into water source. Use of mechanical agitators may cause excessive foaming. Bypass lines should terminate at the bottom of the tank.

For use in knapsack sprayers, it is suggested that the proper amount of this herbicide be mixed with water in a larger container. Fill sprayer with the mixed solution.

### **APPLICATION EQUIPMENT BOOM EQUIPMENT**

**For control of perennial weeds and woody brush and trees listed on this booklet using conventional boom equipment--** Apply this product in 50 to 300 L of clean water per hectare as a broadcast spray using no more than 275 kPa pressure. See "**Weed Control**" sections of this booklet for rates to control specific weeds.

**For control of annual weeds listed on this booklet using conventional boom equipment--**Apply this product in 50 to 100 L of clean water per hectare as a broadcast spray, except as otherwise stated on this label using no more than 275 kPa pressure. See "**Weed Control**" sections of this booklet for rates to control specific weeds.

### **HAND HELD AND HIGH VOLUME EQUIPMENT**

**(use coarse sprays only)**

**For control of weeds and woody brush and trees listed in the "Weed Controlled" section of this label using knapsack sprayers or high volume spraying equipment utilizing handguns or other suitable nozzle arrangements.** Unless otherwise specified, make a 0.75% solution of this product in water (0.75 litre of this product in 100 litres of water) and apply to foliage of vegetation to be controlled. For best results, use a 1.5% solution (1.5 litres of this product in 100 litres of water) on harder to control perennials such as field bindweed, hemp dogbane, milkweed and Canada thistle.

Applications should be made on a spray-to-wet basis. Spray coverage should be uniform and complete. DO NOT spray to point of runoff. Hand gun applications should be properly directed to avoid spraying desirable plants.

### **SELECTIVE EQUIPMENT**

Selective equipment such as **WIPER** and **ROLLER** applicators can be used for weed control in soy and dry beans, orchards, vineyards, cranberries, strawberries and non-crop areas. For information regarding use of this product with selective equipment, refer to "**Selective Equipment**" section of this label.

### **AERIAL EQUIPMENT (PREHARVEST APPLICATION ONLY)**

Aerial application can only be used for weed control in preharvest applications. Refer to "Restricted Use Aerial Preharvest Application for Prairie Provinces Only (Including Peace River Region of B.C.)" section for more information.

### **WEEDS CONTROLLED**

This product controls many annual and perennial grasses, broadleaf weeds, and woody brush and trees when applied as recommended and under conditions described. For information on how to control specific weeds including herbicide rate refer to the "**Annual Weed Control**" and "**Perennial Weed Control**" sections of this label. The following is a partial list of weeds controlled:

2023-2066  
2023-10-24  
Proposed Label on 2023-05-08

**ANNUAL WEEDS**

**Annual Grasses**

**Barnyard Grass**

Echinochloa crusgalli

**Blue Grass (annual)**

Poa annua

**Crab Grass (large)**

Digitaria sanguinalis

**Crab Grass (smooth)**

Digitaria ischaemum

**Downy Brome**

Bromus tectorum

**Fall Panicum**

*Panicum dichotomiflorum*

**Giant Foxtail**

Setaria faberii

**Green Foxtail**

Setaria viridis

**Persian Darnel**

Lolium

persicum

**Volunteer**

**Barley**

Hordeum spp.

**Volunteer Corn**

Zea Mays

**Volunteer Wheat**

Triticum spp.

**Wild Oats**

Avena fatua

**Wild Proso Millet**

*Panicum*

*miliaceum*

**Yellow Foxtail**

Setaria glauca

**Other**

**Dodder**

Cuscuta spp.

2023-2066  
2023-10-24  
Proposed Label on 2023-05-08

**Annual Broadleaf Weeds**

**Chickweed**

*Stellaria media*

**Cleavers**

*Galium aparine*

**Cocklebur**

*Xanthium strumarium* **Corn**

**Spurry**

*Spergula arvensis*

**Cowcockle**

*Saponaria vaccaria*

**Eastern Black Flowering**

**Nightshade**

*Solanum ptycanthum*

**Fleabane (Canada)**

*Erigeron canadensis*

**Flixweed**

*Descurania sophia* **Green**

**Smartweed**

*Polygonum scabrum*

**Hempnettle**

*Galeopsis tetrahit*

**Kochia**

*Kochia scoparia* **Lady's-**

**Thumb**

*Polygonum persicaria*

**Lamb's-Quarters (common)**

*Chenopodium album*

**Narrow-leaved Hawk's Beard**

*Crepis tectorum*

**Narrow-leaved Vetch**

*Vicia angustifolia*

**Night-flowering Catchfly**

*Silene noctiflora*

**Pennsylvania**

**Smartweed**

*Polygonum pensylvanicum*

**Prickly Lettuce**

*Lactuca scariola*

**Ragweed (common)**

*Ambrosia artemisiifolia*

**Redroot Pigweed**

*Amaranthus retroflexus*

**Round-Leaved**

**Mallow**

*Malva pusilla*

**Russian Thistle**

*Salsola pestifer*

**Shepherd's Purse**

*Capsella bursa-pastoris*

**Smooth Pigweed**

*Amaranthus hybridus*

**Sowthistle (annual)**

*Sonchus oleraceus*

**Stinkweed**

*Thlaspi arvense*

**Storksbill**

*Erodium cicutarium*

**Volunteer Canola**

*Brassica* spp

**Volunteer Flax**

*Linaria* spp

**Wild Buckwheat**

*Polygonum convolvulus*

**Wild Mustard**

*Sinapsis arvensis*

**Wild Tomato**

*Solanum triflorum*

**Velvetleaf**

*Abutilon theophrasti*

2023-2066  
2023-10-24  
Proposed Label on 2023-05-08

**PERENNIAL  
WEEDS**

**Perennial Grasses / Sedges**

**Blue Grass (Canada)**

*Poa compressa*

**Blue Grass (Kentucky)**

*Poa pratensis*

**Brome Grass (smooth)**

*Bromus inermis*

**Cattail (common)**

*Typha latifolia*

**Foxtail Barley**

*Hordeum*

*jubatum*

**Quackgrass**

*Agropyron repens*

**Yellow Nutsedge**

*Cyperus esculentus*

**Wire-stemmed**

**Muhly**

*Muhlenbergia*

*frondosa*

**Perennial Broadleaved Weeds**

**Alfalfa**

*Medicago* spp.

**Cottontop**

*Eriophorum chamissionis*

**Curled Dock**

*Rumex crispus*

**Dandelion**

*Taraxacum officinale*

**Field Bindweed**

*Convolvulus arvensis*

**Hemp Dogbane**

*Apocynum cannabinum*

**Hoary Cress**

*Cardaria draba*

**Knotweed (Japanese)**

*Polygonum cuspidatum*

**Milkweed**

**(common)**

*Asclepias*

*syriaca*

**Poison Ivy**

*Rhus radicans*

**Purple Loosestrife**

*Lythrum salicaria*

**Sow**

**Thistle**

**(perennial)**

*Sonchus arvensis*

**Thistle (Canada)**

*Cirsium arvense*

**Toad Flax**

*Linaria vulgaris*

**Wormwood**

**(Absinth)**

*Artemisia absinthium*

2023-2066  
2023-10-24  
Proposed Label on 2023-05-08

**Woody Brush And Trees**

**Alder**

*Alnus* spp.

**Birch**

*Betula* spp.

**Broadleaved meadowsweet**

*Spiraea latifolia*

**Canadian rhododendron**

*Rhododendron canadense*

**Cedar**

*Thuja* spp.

**Cherry**

*Prunus* spp.

**Douglas Fir**

*Pseudotsuga* spp.

**Hemlock**

*Tsuga* spp.

**Maple**

*Acer* spp.

**Mountain-fly honeysuckle**

*Lonicera villosa*

**Pine**

*Pinus* spp.

**Poplar**

*Populus* spp.

**Raspberry / Salmonberry**

*Rubus* spp.

**Sheep laurel**

*Kalmia angustifolia*

**Snowberry (Western)**

*Symphoricarpos occidentalis*

**Sweet fern**

*Comptonia peregrina*

**Willow**

*Salix* spp.

**Withrod**

*Viburnum cassinoides*

2023-2066  
 2023-10-24  
 Proposed Label on 2023-05-08

**CROPLAND USES**

**CROPLAND USES INCLUDE:**

In cropping systems before planting of all crops; in minimum tillage systems, post emergent in glyphosate tolerant soybean, canola and corn; preharvest applications in wheat, barley, oats, canola (rapeseed), flax (including low linolenic acid varieties), peas, lentils, dry beans, soybeans and forages; in pasture renovation; in forage, legume and grass establishments; in tree crops including apple, pear, cherry, plum, peach, apricot, filbert, hazelnut, walnut, chestnut; in grapes, cranberries, blueberries and strawberry; in asparagus, in sugar beets; in tree plantings; and grasses for seed production.

**ALWAYS READ PRECAUTIONARY STATEMENTS, GENERAL INFORMATION and MIXING and APPLICATION PRECAUTIONS PRIOR TO SPECIFIC APPLICATION INFORMATION IN ANY LABEL SECTION.**

**ANNUAL WEED CONTROL**

The following tables provide rates and specific application instructions for control of the annual weeds listed.

**ANNUAL WEED CONTROL WITH DURANGO HERBICIDE**

<b>RATE L/HA</b>	<b>GROWTH STAGE</b>	<b>WEEDS CONTROLLED</b>	<b>COMMENTS (Apply in 50 - 100 L/ha water)</b>
0.56	weeds up to 8 cm in height	wild oats, green foxtail, volunteer barley, volunteer wheat  volunteer canola (rapeseed), wild mustard, lady's-thumb, stinkweed	.for wild oats apply at 1 - 3 leaf stage  .add 350 mL of the surfactant Agral 90, or Ag Surf, or Companion□  .for heavy wild oat infestations use 0.75 L/ha rate.
0.75	weeds 8 cm to 15 cm in height	all annual grasses listed above  all annual broadleaved weeds listed above plus flixweed <sup>†</sup> and kochia <sup>†</sup>	.add 350 mL of surfactant registered for use as listed above.  <sup>†</sup> suppression only. Refer to higher rates of this table or tank-mix table for control options.



2023-2066  
 2023-10-24  
 Proposed Label on 2023-05-08

0.94 - 1.4	weeds up to 15 cm in height	<p>all annual grasses listed above plus downey brome, giant foxtail, and persian darnel</p> <p>all annual broadleaved weeds listed above plus lamb's quarters, redroot pigweed, hempnettle, flixweed, Russian thistle, volunteer flax, common ragweed<sup>†</sup>, Canada fleabane<sup>†</sup>, wild buckwheat<sup>††</sup>, narrowleaved hawk's beard<sup>†††</sup>,</p>	<p>.no surfactant required</p> <p><sup>†</sup> for tank-mix weed control options see annual weed control with tank mixture section</p> <p><sup>††</sup> DO NOT use these rates on plants greater than 8 cm in height</p> <p><sup>†††</sup> for 3 - 4 leaf stage use 1.4 L/ha rate</p> <p><sup>††††</sup> for weeds 8 cm to 15 cm in height use 1.4 L/ha rate</p>
1.69	weeds up to 15 cm in height	<p>all annual grasses listed above plus crab grass and annual blue grass.</p> <p>all annual broadleaved weeds listed above plus kochia, prickly lettuce, shepherd's purse, annual sow thistle, and narrowleaved vetch</p>	.for additional annual broadleaved weed control options, refer to tank-mix table
2.63	weeds over 15 cm in height	all annual grasses and broadleaved weeds listed above	.for additional annual broadleaved weed control options refer to tank-mix table

**NOTE:** For spot treatment, 0.56 – 2.63 L/ha is approximately equivalent to 6-26 mL/100 m<sup>2</sup>, respectively.

**ANNUAL WEED CONTROL WITH DURANGO HERBICIDE TANK MIXTURES FOR SUMMERFALLOW & MINIMUM TILLAGE SYSTEMS**

TANK MIXTURES	RATE L/ha	WEEDS CONTROLLED	COMMENTS (Apply in 50 - 100 L/ha water; add 350mL/ha of surfactant)
---------------	-----------	------------------	--

<p>Durango Herbicide</p> <p style="text-align: center;">+</p> <p>Banvel II</p>	<p>0.56 – 0.75</p> <p style="text-align: center;">+</p> <p>0.29</p>	<p>Volunteer cereal, wild oats, green foxtail,</p> <p>Volunteer canola (rapeseed), wild mustard, flixweed<sup>†</sup>, lamb's quarters, lady's thumb, stinkweed, kochia, Russian thistle, cow cockle, redroot pigweed<sup>††</sup>, wild buckwheat<sup>††</sup>.</p>	<p>This tank-mix is registered for <b>summerfallow use only</b>. Weeds should be less than 15 cm tall and actively growing for best results.</p> <p>Use higher rate if weeds are beyond 8 cm in height.</p> <p><sup>†</sup>Durango Herbicide applied at 0.75 L/ha rate only.</p> <p><sup>††</sup>Suppression only. See other tank mixtures for control options.</p>
<p>Durango Herbicide</p> <p style="text-align: center;">+</p> <p>Pardner</p>	<p>0.56 – 0.75</p> <p style="text-align: center;">+</p> <p>1.25</p>	<p>Volunteer cereals, green foxtail, volunteer canola (rapeseed), wild mustard, lady's thumb, stinkweed, wild buckwheat<sup>†</sup></p> <p>Redroot pigweed<sup>††</sup>, kochia<sup>††</sup>, wild oats<sup>††</sup></p>	<p>This tank-mix is registered <b>only for use in summerfallow, and prior to wheat, oats and barley in minimum tillage systems</b>. Weeds should be less than 15 cm tall and actively growing for best results.</p> <p>Use higher rate if weeds are beyond 8 cm in height.</p> <p><sup>†</sup>use Durango Herbicide at 0.75 L/ha rate only for wild buckwheat control.</p> <p><sup>††</sup>0.75 L rate, suppression only. See other tank mixtures for control options.</p>
<p>Durango Herbicide</p> <p style="text-align: center;">+</p> <p>2,4-D#</p>	<p>0.56 – 0.75</p> <p style="text-align: center;">+</p> <p>1.2</p>	<p>Volunteer cereals, wild oats<sup>†</sup> and green foxtail<sup>†</sup> volunteer canola (rapeseed), wild mustard, flixweed, redroot pigweed, lady's thumb, stinkweed, kochia.</p> <p>Lamb's quarters<sup>††</sup>, Russian thistle<sup>††</sup>.</p>	<p>This tank-mix is registered for <b>summerfallow use only</b>. Weeds should be less than 15 cm tall and actively growing for best results. Use higher rate if weeds are beyond 8 cm in height</p> <p><sup>†</sup>use Durango Herbicide at 0.75 L/ha rate only for wild oat and green foxtail control.</p> <p><sup>††</sup>suppression only. See other tank mixtures for control options.</p>

2023-2066  
 2023-10-24  
 Proposed Label on 2023-05-08

<p>Durango Herbicide + 2,4-D ##</p>	<p>0.94-1.4 + 0.6-0.9<sup>4</sup>  or  1.2-1.5<sup>5</sup></p>	<p>Volunteer cereals, wild oats, green foxtail, downy brome, giant foxtail, and Persian darnel.</p> <p>Volunteer canola, (rapeseed) (non-glyphosate tolerant), wild mustard, flixweed, redroot pigweed, lady's-thumb, stinkweed, kochia, lamb's-quarters, hempnettle, Russian</p>	<p>Weeds should be less than 15 cm tall and actively growing for best results.</p> <p>Use higher rate if weeds are beyond 8 cm in height.</p> <p>No surfactant required.</p> <p>† DO NOT use these rates on plants greater than 8 cm in height.</p> <p>†† For 3-4 leaf stage use 1.4 L/ha rate.</p>
---	--	---	---

		<p>thistle, volunteer flax, common ragweed<sup>†</sup>, Canada fleabane, wild buckwheat<sup>††</sup>, narrowleaved hawk's beard<sup>†††</sup></p> <p>Glyphosate tolerant volunteer canola (1-4 leaf stage)<sup>4</sup>, bluebur<sup>4</sup>, burdock<sup>4</sup>, cocklebur<sup>4</sup>, common plantain<sup>4</sup>, daisy fleabane<sup>4</sup>, false flax<sup>4</sup>, false ragweed<sup>4</sup>, goat's beard<sup>4</sup>, mustards<sup>4</sup> (except dog and tansy), prickly lettuce<sup>4</sup>, ragweeds<sup>4</sup>, Russian pigweed<sup>4</sup>, shepherd's purse<sup>4</sup>, stinging nettle<sup>4</sup>, sweet clover<sup>4</sup>, thymeleaved spurge<sup>4</sup>, wild radish<sup>4</sup>, wild sunflower<sup>4</sup></p> <p>Glyphosate tolerant volunteer canola (4-6 leaf stage)<sup>5</sup>, annual sow thistle<sup>5</sup>, common chickweed<sup>5</sup>, common purslane<sup>5</sup>, dog and tansy mustard<sup>5</sup>, oakleaved goosefoot<sup>5</sup>, groundsel<sup>5</sup>,</p>	<p>††† For weeds 8 cm to 15 cm in height use 1.4 L/ha rate.</p> <p><sup>4</sup> 2,4-D at 0.6 – 0.9 L/ha (280 – 420 g ai/ha).</p> <p><sup>5</sup> 2,4-D at 1.2 – 1.5 L/ha (560 – 700 g ai/ha).</p> <p>Use this tank mix prior to seeding or after seeding but before crop emergence in <b>wheat, winter wheat, barley and rye.</b></p>
--	--	--	---

		hairy galinsoga <sup>5</sup> , hawkweed <sup>5</sup> , heal-all <sup>5</sup> , knotweed <sup>5</sup> , peppergrass <sup>5</sup> , pineapple weed <sup>5</sup> , prostrate pigweed <sup>5</sup> , purslane <sup>5</sup> , sheep sorrel <sup>5</sup> , smartweed <sup>5</sup> , tumble pigweed <sup>5</sup> , velvetleaf <sup>5</sup> , volunteer canola <sup>5</sup>	
Durango Herbicide + MCPA### 500 g/L formulation, if another formulation is used, adjust rate accordingly	0.94-1.4 + 0.5 –0.7 <sup>1</sup> OR 0.5 –1.0 <sup>2</sup>	Volunteer cereals, wild oats, green foxtail, downy brome, giant foxtail, and Persian darnel.  Volunteer canola (rapeseed) (non- glyphosate tolerant), wild mustard, flixweed, redroot pigweed, lady’s thumb, stinkweed, kochia, lamb’s quarters, hempnettle, Russian thistle, volunteer flax, common ragweed <sup>†</sup> , Canada fleabane, wild buckwheat <sup>††</sup> ,	Weeds should be less than 15 cm tall and actively growing for best results.  Use higher rate if weeds are beyond 8 cm in height.  No surfactant required.  <sup>†</sup> DO NOT use these rates on plants greater than 8 cm in height.  <sup>††</sup> For 3-4 leaf stage use 1.4 L/ha rate.  <sup>†††</sup> For weeds 8 cm to 15 cm in height use 1.4 L/ha rate.

2023-2066

2023-10-24

Proposed Label on 2023-05-08

		narrowleaved hawk's beard <sup>†††</sup>  Volunteer glyphosate tolerant	<sup>1</sup> MCPA amine at 0.5 – 0.7 L/ha (250-350 g ai/ha) prior to peas.
--	--	---	--

		canola (1-4 leaf stage) <sup>1,2</sup> , bluebur <sup>3</sup> , burdock <sup>3</sup> (before 4 leaf stage), false flax <sup>3</sup> , flixweed <sup>3</sup> , lamb's quarters <sup>3</sup> , mustards <sup>3</sup> (except dog and tansy), prickly lettuce <sup>3</sup> , ragweeds <sup>3</sup> , redroot pigweed <sup>3</sup> , Russian pigweed <sup>3</sup> , shepherd's purse <sup>3</sup> , stinkweed (field pennycress) <sup>3</sup> , vetch <sup>3</sup> , wild radish <sup>3</sup> , wild sunflower <sup>3</sup>	<sup>2</sup> MCPA at 0.5 – 1.0 L/ha (250-500 g ai/ha) prior to wheat, barley, oats, corn (field and sweet)####, rye and flax.  <sup>3</sup> MCPA at 0.7 – 1.0 L/ha (350 – 500 g ai/ha) only.  Use this tank mix prior to seeding in <b>wheat, barley, rye, oats, corn (field and sweet)####, flax, and field peas###.</b>
--	--	---	---

2023-2066

2023-10-24

Proposed Label on 2023-05-08

<p>Durango Herbicide + Bucril M herbicides</p>	<p>0.94-1.4 + 0.5 –1.0<sup>1</sup></p>	<p>Volunteer cereals, wild oats, green foxtail, downy brome, giant foxtail, and Persian dandel.</p> <p>Volunteer canola (rapeseed) (non-glyphosate tolerant), wild mustard, flixweed, redroot pigweed, lady's thumb, stinkweed, kochia, lamb's quarters, hempnettle, Russian thistle, volunteer flax, common ragweed<sup>†</sup>, Canada fleabane, wild buckwheat<sup>††</sup>, narrowleaved hawk's beard<sup>†††</sup></p> <p>Volunteer glyphosate tolerant Canola (1-4 leaf stage)<sup>1,2</sup></p> <p>Seedlings up to the 4-leaf stage<sup>2</sup>: green smartweed, pale smartweed, lady's thumb, cow cockle, redroot pigweed, flixweed, bluebur, shepherd's purse, kochia<sup>3</sup>, Russian thistle<sup>3</sup>, scentless chamomile<sup>4</sup>, volunteer sunflower, night flowering catchfly, cocklebur, velvetleaf<sup>5</sup>, ball mustard, American nightshade</p> <p>Seedlings up to the 6-leaf stage<sup>2</sup>: wild tomato Seedlings up to the 8-leaf stage<sup>2</sup>: wild buckwheat, tartary buckwheat, common buckwheat, stinkweed, wild mustard, wormseed mustard, lamb's quarters, common ragweed, common groundsel</p>	<p>Weeds should be less than 15 cm tall and actively growing for best results.</p> <p>Use higher rate if weeds are beyond 8 cm in height.</p> <p>No surfactant required.</p> <p><sup>†</sup> DO NOT use these rates on plants greater than 8 cm in height.</p> <p><sup>††</sup> For 3-4 leaf stage use 1.4 L/ha rate.</p> <p><sup>†††</sup> For weeds 8 cm to 15 cm in height use 1.4 L/ha rate.</p> <p><sup>1</sup> Bucril M at 0.5 – 1.0 L/ha (280-560 g ai/ha) for all crops listed.</p> <p><sup>2</sup> Bucril M at 1.0 L/ha (560 g ai/ha only).</p> <p><sup>3</sup> Spray before plants are 5 cm high.</p> <p><sup>4</sup> Spring annuals only.</p> <p><sup>5</sup> Spray before plants are 8 cm high.</p> <p>Use this tank mix prior to seeding in <b>wheat, barley, rye, oats, corn, flax, canary seed and seedling grasses (including brome grass, crested wheatgrass, intermediate wheat grass, slender wheatgrass, tall wheatgrass, Russian wild rye, Timothy, orchard grass, creeping red fescue, meadow fescue, meadow foxtail, seedling tall fescue, seedling meadow brome grass, seedling streambank wheatgrass and reed canary grass.</b></p>
--	--	---	--

2023-2066

2023-10-24

Proposed Label on 2023-05-08

		Perennials (top growth) <sup>2</sup> :	
		Canada thistle, perennial sow thistle	

2023-2066  
 2023-10-24  
 Proposed Label on 2023-05-08

<p>Durango Herbicide + MCPA Amine (500 g/L formulation, if another formulation is used, adjust rate accordingly)</p>	<p>0.94-1.4 +          0.5 –0.7</p>	<p>Volunteer cereals, wild oats, green foxtail, downy brome, giant foxtail, and Persian darnel.</p> <p>Volunteer canola (rapeseed) (non-glyphosate tolerant), wild mustard, flixweed, redroot pigweed, lady’s thumb, stinkweed, kochia, lamb’s quarters, hempnettle, Russian thistle, volunteer flax, common ragweed<sup>†</sup>, Canada fleabane, wild buckwheat<sup>††</sup>, narrowleaved hawk’s beard<sup>†††</sup></p> <p>Volunteer glyphosate tolerant canola (1-4 leaf stage)<sup>3</sup>, bluebur<sup>4</sup>, burdock<sup>4</sup> (before 4 leaf stage), false flax<sup>4</sup>, flixweed<sup>4</sup>, lamb’s quarters<sup>4</sup>, mustards<sup>4</sup> (except dog and tansy), prickly lettuce<sup>4</sup>, ragweeds<sup>4</sup>, redroot pigweed<sup>4</sup>, Russian pigweed<sup>4</sup>, shepherd’s purse<sup>4</sup>, stinkweed<sup>4</sup> (field pennycress), vetch<sup>4</sup>, wild radish<sup>4</sup>, wild sunflower<sup>4</sup></p>	<p>Weeds should be less than 15 cm tall and actively growing for best results.</p> <p>Use higher rate if weeds are beyond 8 cm in height.</p> <p>No surfactant required.</p> <p><sup>†</sup>DO NOT use these rates on plants greater than 8 cm in height.</p> <p><sup>††</sup> For 3-4 leaf stage use 1.4 L/ha rate.</p> <p><sup>†††</sup> For weeds 8 cm to 15 cm in height use 1.4 L/ha rate.</p> <p><sup>3</sup> MCPA amine at 0.5 – 0.7 L/ha (250 – 350 g ai/ha) prior to lentils and chickpeas.</p> <p><sup>4</sup> MCPA amine at 0.7 L/ha (350 g ai/ha) only.</p> <ul style="list-style-type: none"> <li>• Use this tank mix prior to seeding in <b>lentil and chickpea</b>.</li> </ul>
--	---	---	---

For foxtail barley suppression, refer to “**Annual Weed Control**” table

#0.56 kg ai/ha of 2,4-D.

#, ## Adjust rates accordingly for other 2,4-D formulations. Use only low volatile ester or amine formulations of 2,4-D.

### Use only amine formulations of MCPA prior to seeding in corn and field peas.

**ADDITION OF SURFACTANT**

All Durango Herbicide tank mixtures for annual weed control require the addition of the surfactant Agral 90, or Ag Surf, or Companion. Surfactant should be added at a rate of 350 mL per hectare, in 50 – 100 L of clean water.

**Additional Important Information for Annual Weed Control**

Allow at least 1 day after treatment before tillage

Annual weeds generally will continue to germinate from seed throughout the growing season. Repeat treatments may be necessary to control later germinating weeds, in some situations.



2023-2066  
2023-10-24

Proposed Label on 2023-05-08

For additional information and precautions, refer to the "General Information" and "Mixing and Application" sections of this label.

**WEED CONTROL IN GLYPHOSATE TOLERANT CANOLA**

**WARNING: APPLY DURANGO HERBICIDE ON GLYPHOSATE TOLERANT CANOLA VARIETIES ONLY.**

**NOTE: ALWAYS USE PEDIGREED (I.E. CERTIFIED) GLYPHOSATE TOLERANT CANOLA SEED. CANOLA WHICH IS NOT DESIGNATED AS GLYPHOSATE TOLERANT WILL BE DAMAGED OR DESTROYED BY THIS TREATMENT.**

- For additional information and precautions refer to the "General Information" and "Mixing and Application" sections of the Durango Herbicide label.
- Apply Durango Herbicide in glyphosate tolerant canola only as directed in the following weed control table.
- Some short-term, visual yellowing may occur when Durango Herbicide is applied at the late application 4 to 6 leaf stage of the crop. This effect is temporary and will not influence crop growth, maturity or yield.

DO NOT APPLY BY AIR.

The following table describes the rate and specific application instructions for control of annual and perennial weeds in glyphosate tolerant canola varieties.

**WEED CONTROL IN GLYPHOSATE TOLERANT CANOLA**

<b>Rate (L/ha)</b>	<b>Growth Stage Of Crop</b>	<b>Weeds Controlled</b>	<b>Comments (Apply in 50 - 100 L/ha water)</b>
------------------------	-------------------------------------	-------------------------	--

<p>0.60 - 1.4</p>	<p>0 to 6 leaf</p>	<p>Annual Grasses wild oats, green foxtail, volunteer barley, volunteer wheat, barnyard grass</p> <p>Annual Broadleaves stinkweed, redroot pigweed, wild mustard, Russian thistle, lamb's-quarters, non-glyphosate tolerant volunteer canola (rapeseed), hempnettle, lady's thumb, kochia, chickweed, corn spurry, wild tomato, cleavers<sup>†</sup>, wild buckwheat<sup>†</sup>, shepherd's purse<sup>†</sup>, cow cockle<sup>†</sup>, night-flowering catchfly<sup>†</sup>, smartweed<sup>†</sup>, storksbill<sup>†</sup>, flixweed<sup>†</sup>, narrowleaved hawk's beard<sup>†</sup>, round-leaved mallow<sup>††</sup></p> <p>Perennials (suppression) <sup>††</sup> Canada thistle, Perennial sow thistle, Dandelion</p> <p>Perennials (season long control) Quackgrass<sup>††</sup>, foxtail barley <sup>†††</sup>Canada thistle<sup>††††</sup>, Perennial sow thistle<sup>††††</sup></p>	<ul style="list-style-type: none"> <li>• No additional surfactant is required</li> <li>• Repeat applications may be required if a second flush of weeds germinates prior to canopy closure.</li> </ul> <p>Ensure the crop has not advanced beyond the recommended growth stage.</p> <p><sup>†</sup> Use the 0.94 L/ha rate for control of these weeds at all crop growth stages. The lower rate can be used for control of shepherd's purse, cow cockle and night-flowering catchfly at the 1-3 leaf stage of the crop, or for control of smartweed at the 4-6 leaf stage.</p> <p><sup>††</sup> A single application at the 0.94 L/ha rate is required</p> <p><sup>†††</sup> Sequential applications at the 0.94 L/ha rate are required.</p> <p><sup>††††</sup> Sequential applications at the 0.94 L/ha rate are required or a single application of 1.4 L/ha.</p> <p>For sequential applications, ensure the crop has not advanced beyond the recommended growth stage.</p>
			<p>Maximum 1.88 L/ha is allowed for the postemergence use.</p>

**Durango Herbicide plus Lontrel™ 360 Herbicide Tank Mixture**

For hard-to-control weeds (see list below) in glyphosate tolerant canola apply a tank mixture of 0.28 L/ha of Lontrel 360 with 0.94 L/ha of Durango Herbicide in 100L of water per hectare. Apply when canola is in the 2 - 6 leaf stage. Refer to the Lontrel 360 and the Durango Herbicide labels for lists of other weeds controlled, timing of application, water volumes and use precautions. **Apply this tank-mixture in glyphosate tolerant canola only.**

2023-2066  
 2023-10-24  
 Proposed Label on 2023-05-08

**Weeds Controlled**

Canada thistle (season-long top growth) dandelions  
 <15cm diameter (season-long top growth)  
 dandelions >15cm diameter (suppression) perennial  
 sowthistle (season-long top growth) wild buckwheat

**WEED CONTROL IN GLYPHOSATE TOLERANT SOYBEAN**

**WARNING: APPLY DURANGO HERBICIDE ON GLYPHOSATE TOLERANT SOYBEAN VARIETIES ONLY.**

**NOTE: ALWAYS USE PEDIGREED (CERTIFIED) SOYBEAN SEED DESIGNATED AS GLYPHOSATE TOLERANT. SOYBEANS WHICH ARE NOT DESIGNATED AS GLYPHOSATE TOLERANT WILL BE DAMAGED OR DESTROYED BY THIS TREATMENT.**

DO NOT APPLY BY AIR.

**WEED CONTROL IN GLYPHOSATE TOLERANT SOYBEAN**

<b>Rate (L/ha)</b>	<b>Growth Stage of Crop</b>	<b>Weeds Controlled<sup>†</sup></b>	<b>Comments (use 100-200 L/ha water volumes)</b>
1.88	First trifoliolate leaf stage through to flowering.	velvetleaf, common ragweed, common lambsquarters, redroot pigweed, smooth pigweed, cocklebur, green smartweed, ladythumb, Pennsylvania smartweed, eastern black flowering nightshade, wild mustard, wild buckwheat, foxtail (green, yellow, giant), barnyard grass, crabgrass (smooth, large), quackgrass, milkweed <sup>†</sup> , yellow nutsedge <sup>†</sup> , fall panicum, wild proso millet	<ul style="list-style-type: none"> <li>• A second application may be used for late weed flushes emerging after the initial treatment</li> <li>† suppression only.</li> <li>• This second application must be made no later than the flowering stage of the soybean.</li> </ul>
1.88-3.75	First trifoliolate leaf stage through to flowering.	Perennial sow thistle, Canada thistle, wire-stemmed muhly	<ul style="list-style-type: none"> <li>• A single application at the higher rate or a second (sequential) application of 1.88 L/ha will improve control in heavy weed infestations.</li> <li>• If sequential applications of 1.88</li> </ul>

2023-2066

2023-10-24

Proposed Label on 2023-05-08

			<p>L/ha are used they should be at least 2 weeks apart for best results on perennial weeds.</p> <ul style="list-style-type: none"><li>• This second application must be made no later than the flowering stage of the soybean.</li><li>• Perennial sow thistle and Canada thistle should be from the rosette stage to 50 cm in height and actively growing.</li><li>• Wire-stemmed muhly should be 10-20 cm in height and actively growing.</li><li>• Plants not fully emerged at the time of application will escape the treatment.</li></ul>
3.75	First trifoliolate leaf stage through to flowering.	All weeds listed above, plus milkweed <sup>††</sup> , yellow nutsedge <sup>††</sup> , field bindweed <sup>††</sup>	<ul style="list-style-type: none"><li>• Only one application per season at 3.75 L/ha.</li></ul> <p><sup>††</sup> Will also be controlled by sequential applications of 1.88 L/ha. Applications should be at least 2 weeks apart for optimum control.</p> <ul style="list-style-type: none"><li>• This second application must be made no later than the flowering stage of the soybean.</li><li>• Milkweed should be 15-60 cm in height and actively growing; nutsedge should be 5-</li></ul>

2023-2066  
2023-10-24  
Proposed Label on 2023-05-08

			15 cm in height and actively growing.  Plants not fully emerged at the time of application will not be controlled
--	--	--	---

† Weeds will be more easily controlled and early crop competition avoided with applications made when the weeds are small. Control of weeds greater than 25 cm in height will be inconsistent, although some weeds may be controlled.

#### **Durango Herbicide plus Pursuit Herbicide Tank Mixture**

For added residual control of late germinating eastern black nightshade, common lamb's quarters, redroot pigweed, velvetleaf, fall panicum and wild proso millet, Pursuit herbicide may be tank mixed with Durango Herbicide at a rate of 1.88 litres per hectare. Use 0.16 to 0.21 litres per hectare of Pursuit Herbicide and apply up to and including the 3rd trifoliolate leaf stage of the glyphosate tolerant soybeans in 100-200 litres per hectare of clean water. The higher rate is recommended for heavier infestations. This tank mix is recommended primarily for soybean systems with row spacings of 50 centimetres (20 inches) or more where a single application timing is desired.

Mixing: Add and mix Pursuit Herbicide as per instructions on the Pursuit Herbicide label and then add Durango Herbicide as per instructions on this label.

A PHI of 100 days is required for the tank mix of Durango Herbicide and Pursuit herbicide on glyphosate tolerant soybeans.

Only one application per season of Durango Herbicide at 1.88 litres per hectare tank mixed with Pursuit herbicide at 0.16 to 0.21 litres per hectare is permitted.

Refer to the Pursuit herbicide label for further safety precautions and handling instructions.

#### **Durango Herbicide plus Assure II Tank Mixture**

For control of volunteer glyphosate tolerant corn, Assure II Herbicide may be tank mixed with Durango Herbicide. Use 1.88 – 3.75 litres per hectare Durango Herbicide and 0.38 litre per hectare of Assure II Herbicide.

2023-2066  
2023-10-24  
Proposed Label on 2023-05-08

Apply in 100 – 300 litres per hectare of clean water.

Mixing: Add and mix Assure II Herbicide as per instructions on the Assure II Herbicide label and then add Durango Herbicide as per instructions on this label.

This tank mix is to be applied when the crop is from the first trifoliolate leaf stage through flowering and when the volunteer glyphosate tolerant corn is at the 2-6 leaf stage.

A PHI (preharvest interval) of 80 days is required for the tank mix of Durango Herbicide and Assure II Herbicide on glyphosate tolerant soybeans.

Refer to the Assure II Herbicide label for further safety precautions and handling

**WEED CONTROL IN GLYPHOSATE TOLERANT CORN**

WARNING: APPLY DURANGO HERBICIDE ON GLYPHOSATE TOLERANT CORN VARIETIES ONLY.

NOTE: ALWAYS USE PEDIGREED (CERTIFIED) CORN SEED DESIGNATED AS GLYPHOSATE TOLERANT. CORN WHICH IS NOT DESIGNATED AS GLYPHOSATE TOLERANT WILL BE DAMAGED OR DESTROYED BY THIS TREATMENT.

DO NOT APPLY BY AIR.

<b>Rate (L/ha)</b>	<b>Growth Stage of Crop</b>	<b>Weeds Controlled†</b>	<b>Comments (use 100-200 L/ha water volumes)</b>
--------------------	-----------------------------	--------------------------	--

2023-2066  
2023-10-24

Proposed Label on 2023-05-08

1.88	Up to and including 8 leaf stage.	Velvetleaf, common ragweed, common lambsquarters, redroot pigweed, smooth pigweed, cocklebur, green smartweed, ladythumb, Pennsylvania smartweed, eastern black flowering nightshade, wild mustard, wild buckwheat, foxtail (green, yellow, giant), barnyard grass, crabgrass (smooth, large), quackgrass, fall panicum, wild proso millet	A second application may be used for late weed flushes emerging after the initial treatment.
1.88		Wild oats, volunteer barley, volunteer wheat, stinkweed, wild mustard, Russian thistle, nonglyphosate tolerant canola (rapeseed), hempnettle, kochia, chickweed, corn spurry, wild tomato, cleavers, shepherd's purse, cow cockle, nightflowering catchfly, stork's-bill, flixweed, narrow-leaved hawk's beard	This second application must be made no later than the 8 leaf stage of the corn.

2023-2066  
2023-10-24  
Proposed Label on 2023-05-08

		Common milkweed, yellow nutsedge, roundleaved mallow, field bindweed	<p>For control of common milkweed, yellow nutsedge, roundleaved mallow and field bindweed use two applications of 1.88 L/ha.</p> <p>This second application must be made no later than the 8 leaf stage of the corn.</p> <p>Milkweed should be 15-60 cm in height and actively growing.</p> <p>Yellow nutsedge should be 5-15 cm in height and actively growing.</p>
--	--	---	--

1.88		Perennial sow thistle, Canada thistle, wirestemmed muhly	<p>A second (sequential) application of 1.88 L/ha will improve control in heavy weed infestations.</p> <p>If sequential applications are used they should be at least 2 weeks apart for best results on perennial weeds.</p> <p>This second application must be made no later than the 8 leaf stage of the corn.</p> <p>Perennial sow thistle and Canada thistle should be from the rosette stage to 50 cm in height and actively growing.</p> <p>Wire-stemmed muhly should be 10-20 cm in height and actively growing.</p>
------	--	--	---



2023-2066  
 2023-10-24  
 Proposed Label on 2023-05-08

			Plants not fully emerged at the time of application will escape treatment.
1.88 Durango Herbicide + 0.75–1.0 kg ai/ha atrazine <sup>†</sup>	Up to and including 5th leaf stage.	Residual control of lamb'squarters, redroot pigweed, common ragweed	Tank mix should be used when only a single application timing is desired. Use higher rate of atrazine for heavier weed infestations.
1.88 Durango Herbicide + 2.5-3.7 Marksman Herbicide	Up to and including 5th leaf stage.	Residual control of lamb'squarters, redroot pigweed, common ragweed, velvetleaf	Tank mix should be used when only a single application timing is desired. Use higher rate Marksman Herbicide for heavier weed infestations.

<sup>†</sup> 0.75-1.0 kg ai atrazine/ha is equivalent to 1.56-2.08 L/ha of Clean Crop Atrazine 480, or Aatrex Liquid 480, or 0.83-1.11 kg/ha Aatrex Nine-O.

<sup>†</sup>Weeds will be more easily controlled and early crop competition avoided with applications made when the weeds are small. Control of weeds greater than 25 cm in height will be inconsistent, although some weeds may be controlled.

2023-2066  
 2023-10-24  
 Proposed Label on 2023-05-08  
**PERENNIAL WEED CONTROL**

**ALWAYS READ PRECAUTIONARY STATEMENTS, GENERAL INFORMATION and MIXING and APPLICATION SECTIONS PRIOR TO SPECIFIC APPLICATION INFORMATION IN ANY LABEL SECTION.**

When applied as recommended under the conditions described, this product will control the perennial weeds listed in the following table:

**PERENNIAL WEED CONTROL WITH DURANGO HERBICIDE**

WEED	APPLICATION			COMMENTS
	GROWTH STAGE	RATE (L/ha)	WATER VOLUME (L/ha)	
<b>Quackgrass</b> (control, light to moderate infestations)	3 to 4 green leaves or more	1.88	50 - 300	.Apply in clean water using flat fan nozzles. .Allow 3 or more days after treatment before tillage. .Refer to " <b>Quackgrass</b> " notes for more information. .For higher water volumes (ie. 150 - 300 L/ha) an approved surfactant must be added at 0.5 litres per 100 litres of clean water (0.5% v/v). Refer to list of surfactants. See also below.
<b>Quackgrass</b> (long term control, heavy infestations, high water volumes)	3 to 4 green leaves or more	1.88 – 5.25	50 - 300	.Allow 3 or more days after treatment before tillage. .Rates higher than 1.88 L/ha will provide more consistent, longer term control, especially with heavier infestations and/or higher water volumes (ie 150-300 L/ha) .Refer to " <b>Quackgrass</b> " notes for more information.
<b>Canada Thistle</b>	rosette stage (summer-fallow)	1.88	50 - 100	.Apply in clean water using flat fan nozzles. .Allow 10 or more days after treatment before tillage. .Refer to notes in "Canada Thistle" section for more information.
<b>Canada Thistle</b>	bud stage or beyond	3.56 – 5.25	100 - 300	.Allow 5 or more days after treatment before tillage.

2023-2066

2023-10-24

Proposed Label on 2023-05-08

<b>Field Bindweed</b>	full bloom or beyond	5.25 – 9	100 - 300	.Allow 7 or more days after treatment before tillage.
-----------------------	----------------------	----------	-----------	---

<b>Common Milkweed<sup>†</sup></b>	bud to full bloom (preharvest)	1.88	50 - 100	.See preharvest application section  .Allow 7 or more days after treatment before tillage.
	bud to full bloom	9	100 - 300	.Reduced control may occur after full bloom  .Milkweed may not all be in the correct stage, therefore, repeat treatments may be required.
<b>Toadflax</b>	Vegetative Stage (summerfallow)	1.88	50-100	.Apply in clean water using flat fan nozzles
	Bud to Full Bloom (preharvest)			.Allow 7 or more days after treatment before tillage in summerfallow  .For more information, see summerfallow control, or preharvest control
<b>Alfalfa</b>	Early bud to full bloom stage. Fall applications only	2.8 – 3.75	50 - 300	.Allow 5 or more days after treatment before tillage. Use the higher rates when alfalfa populations are high or when heavy grass infestations are also present.  .For spring applications and control in minimum tillage systems using a 2,4-D tank-mix

2023-2066  
 2023-10-24  
 Proposed Label on 2023-05-08

<b>Dandelion</b>	< 15 cm	1.88	50 - 100	.Allow 3 or more days after treatment before tillage for all rates.
	> 15 cm	2.78 – 3.75	50 - 300	.Use the higher rate when infestations are heavy.  . Refer to notes in Dandelion Section for more information.
	Rosette to full bloom (preharvest)	1.88	50 - 100	.Allow 7 or more days after treatment before tillage. For more information, see preharvest control section.

<b>Foxtail barley</b>	Seedling to heading	1.88 – 3.75	50 -100	-Allow a minimum of 1 day after treatment before tillage or seeding.  -Use higher rates for larger, more established plants, heavy infestations or if plants are stressed
<b>Other Perennials</b> (see perennial weeds listing)	early heading or early bud stage	5.25 – 9	100 - 300	.Allow 7 or more days after application before tillage.

†**NOTE:** For spot treatment, mix 90 mL of product in 5 L clean water per 100 m<sup>2</sup>. (1.88 – 9 L/ha is approximately equivalent to 19 – 90 L/100 m<sup>2</sup>, respectively).

#### **SPECIAL NOTES FOR PERENNIAL WEED CONTROL QUACKGRASS**

For **season-long control on fall tilled ground:** Apply 1.88L/ha of this product in spring prior to seeding. Apply in 50 to 100 L/ha of clean water as described in the preceding table. Delay application until the majority of quackgrass plants have 4-5 green leaves. This stage usually occurs 1 to 4 weeks later on fall tilled ground than on undisturbed ground. Reduced control may result on ground tilled deeper than 15 cm.

#### **NOTE:**

This treatment will provide season-long control of quackgrass on fall tilled ground. Reduced control will be experienced versus this product on non-fall tilled ground. Repeat treatments may be necessary.

**Applications on forages** should be followed by tillage 3 days or later and should be made when good growing conditions exist.

2023-2066  
2023-10-24  
Proposed Label on 2023-05-08

**If a frost has occurred**, wait several days to determine if the quackgrass has recovered. Quackgrass can be treated after a mild frost provided there are 3 to 4 green leaves actively growing at the time of application. DO NOT apply after the first damaging frost in the fall.

**Surfactant Information:**

The following is a list of approved surfactants for use with Durango Herbicide for control of quackgrass:  
Agral 90 Companion Ag Surf Frigate

**Always refer to surfactant label for specific instructions regarding use of that product.**

**CANADA THISTLE**

**Control of Canada thistle at the rosette stage:** To ensure the proper timing of application the following steps must be followed:

1. Conduct summerfallow tillage as usual and perform the last tillage operation between July 15th and August 1.
2. Allow the thistles to regrow for a minimum of 5 weeks until they are a minimum of 15cm in diameter and in the rosette stage of growth.

**NOTE: Canada thistle can be treated after a mild frost** provided the leaves are still green and actively growing at the time of application. DO NOT apply after the first damaging frost in the fall.

**Durango Herbicide plus Banvel II Tank Mixtures**

For control of Canada thistle (and perennial sow thistle) in summerfallow or in post-harvest stubble, apply 1.28 L/ha Durango Herbicide plus 1.25 L/ha Banvel II in 100 – 200 L/ha of clean water. In addition, add 350 mL/ha of a non-ionic surfactant registered for use with this product, such as Agral 90, Ag Surf, or Companion.

For best results in summerfallow, cultivate in the spring and apply when the majority of thistles are 15 cm to 25 cm tall and before the bud stage. Cultivate 3 weeks after application.

In post harvest stubble, apply this tank mixture to actively growing thistles at least 2 weeks prior to a killing frost.

**NOTE:**

Grow only cereals, canola (including rapeseed), soybeans, field corn, sweet corn, or white beans after application of this tank mixture.

If application is made after September 1<sup>st</sup>, or if soil moisture levels are extremely low after application, crop injury may occur in the spring following application.

**TOADFLAX**

**Control of Toadflax in a Summerfallow Vegetative Stage**

To ensure the proper timing of application, the following steps must be followed:

1. Conduct summerfallow tillage as usual and perform the last tillage operation between July 10-21.
2. Allow toadflax to regrow for a minimum of 4-5 weeks until they are minimum of 15 cm tall and at a lush green vegetative stage.

Note: Toadflax can be treated after a mild frost provided the leaves are still green and actively growing at the time of application. DO NOT apply after the first damaging frost.

2023-2066  
2023-10-24  
Proposed Label on 2023-05-08

## **DANDELION**

Applications should be made up to and including bloom for best results. Follow-up control measures should be used to manage new dandelions germinating from seed to maintain control throughout the season.

### **ALFALFA CONTROL WITH 2,4-D TANK-MIX:**

The addition of 2,4-D may improve alfalfa control in situations where control may be more difficult to obtain, such as in minimum tillage systems where populations are heavy, and with spring applications.

For fall control of established stands of alfalfa, apply 1.88 – 3.75 L/ha Durango Herbicide– and 1.2 – 2.4 L/ha of any 500 g/L 2,4-D amine or low volatile ester formulation in 100 – 200 L water/ha. (Adjust product rates accordingly for other 2,4-D formulations).

For spring applications, use only the low rate of 2,4-D (i.e. 1.2 L/ha) and 1.88 – 3.75 L/ha Durango Herbicide. Only cereal crops not underseeded to legumes may be planted following spring applications of this tank-mix, and a 14 day interval between application and planting is required.

Use the higher Durango Herbicide rates when perennial grasses are prevalent.

### **ALL PERENNIAL WEEDS**

**Weed Stages:** Weeds must be at the proper stage for effective control. Refer to "**Perennial Weed Control with Durango Herbicide** "

**Nozzle Type:** For best results with conventional boom equipment apply this product with 50 to 300 L/ha of clean water using flat fan nozzles and no more pressure than 275 kPa.

**Rhizome Dormancy:** Reduced control may result if rhizomes have become dormant. Dormancy may occur if soil fertility is low and/or the land has not been tilled for several years.

**Mowing Effects:** Mowing prior to application will reduce effectiveness unless weeds are allowed to regrow to the proper stage before application.

**Tillage Effects:** Fall or spring tillage prior to spring applications and tillage between harvesting and fall applications will reduce the effectiveness on perennial weeds. Follow-up tillage after application should be delayed 5-7 days for best results (see Weed Control Table for specific tillage interval for each weed).

**Rainfall Effects:** Heavy rainfall immediately after application may wash the chemical off the foliage and a repeat treatment may be required. DO NOT apply if rainfall is forecast for the time of application.

**Regrowth from Germinating Seeds:** This product only controls emerged plants. Repeat treatments or other weed control measures may be required to control weeds regenerating from seeds or other underground parts.

**Frost Effects:** Heavy frosts prior to application may reduce control. DO NOT apply after the first damaging frost in the fall.

### **CROPLAND SITUATIONS**

**ALWAYS READ PRECAUTIONARY STATEMENTS, GENERAL INFORMATION and MIXING and APPLICATION SECTIONS PRIOR TO SPECIFIC APPLICATION INFORMATION IN ANY LABEL SECTION.**

2023-2066

2023-10-24

Proposed Label on 2023-05-08

This product can be applied as a broadcast spray or spot treatment prior to planting all crops, post harvest to annual crops, preharvest in wheat, barley, oats, canola (rapeseed), flax (including low linolenic acid varieties), lentils, peas, soybeans, dry beans and forages, and in summerfallow. It may also be applied as a broadcast spray in glyphosate tolerant canola, soybean or corn (refer to sections on Weed Control in Glyphosate Tolerant Canola, Soybean or Corn). It can also be applied as a directed spray in orchards, vineyards, blueberries and strawberry, and using selective equipment in soy and dry beans, orchards, vineyards, cranberries and strawberry (refer to specific sections below for more information). **For specific instructions on weed control in the following cropping situations, always refer to the Annual and Perennial Weed Control sections for more information.**

### **Prior to Planting - All Crops**

This product may be applied prior to planting all crops for control of emerged weeds listed on this label. Ensure weeds are at the desired stage at the time of application. This product does not provide preemergent weed control and newly germinating weeds may be a problem in the crop. **APPLY BEFORE SEEDING OR TRANSPLANTING.**

### **Post Harvest Stubble Treatment**

This product may be applied in the fall as a postharvest stubble treatment for control of perennial weeds such as quackgrass and Canada thistle. Allow weeds to regrow to the desired stage (20-25 cm tall for quackgrass and Canada thistle) before application and ensure they have a high proportion of green coloration. Straw should be removed or evenly spread to allow for proper regrowth and spray coverage. Heavy frosts prior to application may decrease control.

### **Spot Treatment (In-Crop)**

This product can be applied as an in-crop spot treatment in barley, corn, oats, soybeans, wheat, strawberry, blueberry, forage grasses and legumes including seed production. Applications should be made using the same rates and at the same growth stages as listed in the weed control tables or use a 0.75% solution for annual weeds and quackgrass and a 1.5% solution for other perennial weeds (a 0.75% solution equals 0.75 litre Durango Herbicide in 100 litres of spray solution). The 0.75 or 1.5 per cent solutions should be applied to wet, but not run-off. Applications can be made using a boom sprayer, hose and handgun, or hand sprayer in accordance with instructions in the "Application Equipment" section.

### **Grazing Restrictions**

Applications can be made up to heading of small grains, initial pod set on soy and dry beans, silking of corn and emergence of seed heads. The crop in the treated area will be killed. Take care to avoid drift for the same reason. **DO NOT APPLY IF CROP GROWTH HAS ADVANCED BEYOND SEED SET.**  
ALLOW 3

**TO 5 DAYS FOR DURANGO HERBICIDE TO TRANSLOCATE INTO ALL PLANT PARTS BEFORE GRAZING OR HARVESTING TREATED AREAS IN FORAGES.**

### **Summerfallow Treatment**

This product, or labelled tank mixtures, may be applied in summerfallow to control weeds listed on this label. Ensure weeds are at the desired growth stage and actively growing at application for best results. Reduced control may result if weeds are drought stressed. Weeds will continue to germinate from seed throughout the growing season. Repeat treatments may be necessary to control later germinating weeds.

### **Minimum and Zero Tillage Cropping Systems (All Field Crops, including cereals, oilseeds, pulses, forages, corn and potatoes)**

This product may be applied prior to seeding or after seeding, but before crop emergence for control of emerged weeds in minimum and zero tillage cropping systems for all field crops. Applications made too far in advance of seeding may allow weeds to emerge between application and crop emergence, as this product does not provide residual weed control.

2023-2066

2023-10-24

Proposed Label on 2023-05-08

#### **Minimum and Zero Tillage Tank Mixtures**

**Durango Herbicide plus bromoxynil (Pardner)** can be applied prior to seeding or after seeding, but before crop emergence in **wheat, barley and oats**. Refer to “**Annual Weed Control with Durango Herbicide Tank Mixtures**” table for information.

**Durango Herbicide plus Pursuit** can be applied prior to, or after, seeding, but before crop emergence in soybeans. Durango Herbicide will control emerged weeds listed on this label when applied as directed (refer to Annual and Perennial Weed Control sections). Pursuit will control weeds germinating from seed. Add the recommended rates of both products in 100 L of water/ha, following the instructions on the Pursuit herbicide label.

**ALWAYS REFER TO THE PURSUIT LABEL FOR FURTHER INFORMATION ON WEEDS CONTROLLED, APPLICATION DIRECTIONS, AND USE PRECAUTIONS. ONLY SOYBEANS, FIELD CORN, SPRING BARLEY, SPRING WHEAT AND WINTER WHEAT MAY BE PLANTED THE SEASON FOLLOWING A PURSUIT APPLICATION. WINTER WHEAT MAY BE PLANTED THE SAME YEAR AS A PURSUIT APPLICATION TO SOYBEANS, BUT NOT EARLIER THAN 120 DAYS AFTER THE APPLICATION.**

**DO NOT APPLY AFTER CROP EMERGENCE.**

**Durango Herbicide plus MCPA** can be applied prior to seeding in wheat, barley, rye, oats, corn (field and sweet; MCPA amine only), flax and field peas (MCPA amine only). Refer to “**Annual Weed Control with Durango Herbicide Tank Mixtures**” table for information.

**Durango Herbicide plus Buctril M** can be applied prior to seeding in **wheat, rye, corn, barley, oats, flax, canary seed and seedling grasses (including brome grass, crested wheatgrass, intermediate wheat grass, slender wheatgrass, tall wheatgrass, Russian wild rye, Timothy, Orchard grass, creeping red fescue, meadow fescue, meadow foxtail, seedling tall fescue, seedling meadow bromegrass, seedling streambank wheatgrass and reed canary grass** Refer to “**Annual Weed Control with Durango Herbicide Tank Mixtures**” table for information.

**Durango Herbicide plus MCPA amine** can be applied prior to seeding in **lentil and chickpea** Refer to “**Annual Weed Control with Durango Herbicide Tank Mixtures**” table for information.

#### **Forage Legumes and Grasses**

This product may be applied for control of emerged weeds prior to emergence of forage legumes and grasses. If the forages are to be under-seeded with a cover crop, this product must be applied prior to planting the cover crop.

#### **Pasture Renovation**

Use this product to control or suppress existing vegetation for zero-tillage seeding of legumes into established sod for pasture renovation. Delay spraying until weed growth is at least 20 cm in height and a maximum number of seedlings or shoots have emerged. Application can be made immediately before, during or after seeding, but before crop emergence.

#### **Forage Seed Production**

For spot treatment control of perennial weed problems such as quackgrass and Canada thistle in seed fields, apply as directed to vegetation that is at least 20 to 25 cm in height but before emergence of seed head. The crop in the treated areas will be killed. Take care to avoid drift outside target area for the same reason.

**PRE-HARVEST CONTROL OF QUACKGRASS, CANADA THISTLE, MILKWEED, TOADFLAX and DANDELION; SEASON-LONG CONTROL OF PERENNIAL SOW THISTLE, AND HARVEST MANAGEMENT**



2023-2066

2023-10-24

Proposed Label on 2023-05-08

For control of quackgrass, Canada thistle, common milkweed, toadflax and dandelion; and season-long control of perennial sow thistle, Durango Herbicide can be applied prior to harvest of wheat, barley (including malting barley), oats, canola (rapeseed), flax (including low linolenic acid varieties), lentils, peas, dry beans, soybeans and forages. DO NOT apply to crops if grown for seed production.

This treatment may also provide harvest management benefits, by drying down crop and weed vegetative growth, for example, where late flushes of annual weeds, green vegetative crop growth, or late tillering may interfere with harvest operations. EXTREMELY COOL, WET AND/OR CLOUDY WEATHER CONDITIONS BETWEEN THE TIME OF APPLICATION AND THE ANTICIPATED HARVEST DATE MAY SLOW DOWN ACTIVITY OF THIS PRODUCT, THEREBY DELAYING CROP DRYDOWN AND HARVEST DATE.

Durango Herbicide should be applied pre-harvest at 1.88 L/ha in 50 to 100 L/ha of clean water, by ground application only. Apply only when the crop has 30% or less grain moisture content. This stage typically occurs 7 to 14 days before harvest. For forage crops, apply this product at 1.88 – 3.75 L/ha 3-7 days prior to the last cut before rotation or forage renovation. Consult the table "Guidelines for Timing of Preharvest Applications" for visual indicators of this stage in each crop. For the best weed control results quackgrass should be actively growing and have at least 4 to 5 green leaves. Canada thistle and perennial sow thistle should be actively growing and at or beyond the bud stage for best results. Common milkweed should be at the bud to bloom stage and actively growing for best results. Applications for weed control (not for harvest management) must be made at the correct stage of both weed and crop growth.

Apply only during the period 7-14 days (or 3-7 days for forage applications) before harvest to ensure best weed control and to maximize harvest management benefits. Earlier application may reduce crop yield and/or quality, and may lead to excess glyphosate residues in the crop.

DO NOT expose or contaminate any body of water or non-target vegetation by direct application, spray drift, or when cleaning and rinsing spray equipment.

DO NOT APPLY BY AIR.

#### GUIDELINES FOR TIMING OF PREHARVEST APPLICATIONS

CROP(S)	PERCENT GRAIN MOISTURE	VISUAL SYMPTOMS
WHEAT/BARLEY/ OATS	Less than 30	Hard dough stage; a thumbnail impression remains on seed.
CANOLA	Less than 30	Pods are green to yellow; most seeds are yellow to brown.
FLAX (INCLUDING LOW LINOLENIC ACID VARIETIES)	Less than 30	Majority (75%-80%) of bolls are brown.
PEAS	Less than 30	Majority (75%-80%) of pods are brown.
LENTILS	Less than 30	Lowermost pods (bottom 15%) are brown and seeds rattle.

2023-2066  
2023-10-24  
Proposed Label on 2023-05-08

DRY BEANS	Less than 30	Stems are green to brown in colour; pods are mature (yellow to brown in colour); 80%-90% leaf drop (original leaves).
SOYBEANS	Less than 30	Stems are green to brown in colour; pod tissue is dry and brown in appearance; 80%-90% leaf drop.
FORAGES	Not applicable	Normal stage for forage harvesting.

**RESTRICTED USE AERIAL PREHARVEST APPLICATION FOR PRAIRIE PROVINCES ONLY  
(Including PEACE RIVER REGION OF B.C.)**

**NOTICE TO USER:** This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way this is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

**NATURE OF RESTRICTION:** This product is to be used only in the manner authorized. For use only by aerial applicators and aerial application services approved by the provincial regulatory agency to apply this product with aerial application equipment. To qualify for consideration of provincial approval, the following requirements must be demonstrated to the provincial regulatory agency:

1. Aircraft used in the application of this product must have been configured and calibrated to acceptable standards at a recognized calibration (patternation) clinic within 20 months of the date of application. The spray system must not have been subjected to major changes (new nozzles, booms or configurations) since the calibration, and must meet critical drift management standards e.g.  
maximum boom width 65% of wing span; nozzle type, size and orientation to minimize drift and deliver droplet size VMD in the coarse (400-600 microns) or very coarse (600 - 1000 microns) range.

2023-2066

2023-10-24

Proposed Label on 2023-05-08

2. Aircraft used in the application of this product must carry a minimum of \$25,000 drift insurance in addition to any provincial requirements for general comprehensive insurance coverage.
3. Applicators using this product must have successfully completed an aerial application training course provided by Albaugh, LLC .
4. Aerial application services applying this product must employ on staff at least one pilot applicator with at least 250 hours of actual aerial application time and a minimum of 100 hours within the last 24 month period. All pilots who DO NOT meet the minimum experience standard must work under the direct daily supervision of a qualified pilot.

#### **DIRECTIONS FOR USE**

Durango Herbicide may be applied with aerial application equipment for control of quackgrass, Canada thistle, common milkweed, toadflax and dandelion, and season-long control of perennial sow thistle. Durango Herbicide can be applied prior to harvest of wheat, barley (including malting barley), oats, canola (rapeseed), flax (including low linolenic acid varieties), lentils, peas, dry beans, and soybeans. DO NOT apply to any crops if grown for seed production.

This treatment may also provide harvest management benefits, by drying down crop and weed vegetative growth, for example, where late flushes of annual weeds, green vegetative crop growth, or late tillering may interfere with harvest operations.

**EXTREMELY COOL, WET AND/OR CLOUDY WEATHER CONDITIONS BETWEEN THE TIME OF APPLICATION AND THE ANTICIPATED HARVEST DATE MAY SLOW DOWN ACTIVITY OF THIS PRODUCT, THEREBY DELAYING CROP DRYDOWN AND HARVEST DATE.**

Durango Herbicide should be applied at 1.88 L/ha in 20 - 50 L/ha of clean water with aerial application equipment. Apply only when the crop has 30% or less grain moisture content. This stage typically occurs 7 to 14 days before harvest. Consult the table "Guidelines for Timing of Preharvest Applications" for visual indicators of this stage in each crop. For the best weed control results quackgrass should be actively growing and have at least 4 to 5 green leaves. Canada thistle and perennial sow thistle should be actively growing and at or beyond the bud stage for best results. Common milkweed should be at the bud to bloom stage and actively growing for best results. Applications for weed control (not for harvest management) must be made at the correct stage of both weed and crop growth.

Apply only during the period 7 - 14 days before harvest to ensure best weed control and to maximize harvest management benefits. Earlier application may reduce crop yield and/or quality, and may lead to excess glyphosate residues in the crop.

#### **USE PRECAUTIONS**

**AVOID DRIFT ON TO IMPORTANT WILDLIFE HABITATS. EXTREME CARE MUST BE TAKEN WHEN APPLYING THIS PRODUCT TO PREVENT INJURING DESIRABLE PLANTS AND CROPS.**

Apply only in wind conditions in compliance with local and/or provincial regulations. DO NOT apply when other climatic conditions, including lesser wind velocities, will allow significant drift to occur.

Coarse sprays are less likely to drift, therefore DO NOT use nozzles or nozzle configurations which disperse spray as fine spray droplets. DO NOT angle nozzles forward into the airstream

2023-2066

2023-10-24

Proposed Label on 2023-05-08

and DO NOT increase spray volume by increasing nozzle pressure. See # 1 of the Nature of Restrictions for additional details.

DO NOT overspray or allow drift on to bodies of water, wetlands<sup>†</sup> and/or wetland vegetation (e.g., sloughs, swamps, bogs, marshes, potholes), shelterbelts, woodlots and other cover on the edge of fields.

DO NOT apply directly to roadside ditches, or apply under conditions that would favour drift into roadside

2023-2066  
2023-10-24  
Proposed Label on 2023-05-08  
ditches.

†A wetland is any land where the water table stands at or above the land surface for at least part of the year, and contains vegetation associated with wetlands such as bulrushes, sedges, cattails, etc.

Ensure uniform application - To avoid streaked, uneven or overlapped application, use appropriate marking devices, or equivalent electronic positioning systems (GPS).

Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of this product accumulated during spraying or from spills. **PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. LANDING GEAR ARE MOST SUSCEPTIBLE.** The maintenance of an organic coating (paint) which meets aerospace specification MIL-C 38412 may prevent corrosion.

### **Tree Plantings Shelterbelts and Nursery Stock (Woody Ornamentals)**

This product may be used to control listed annual or perennial weeds prior to planting, or as a post directed spray in established nurseries or shelterbelts of the following species:

#### **Deciduous**

Ash - Fraxinus spp.  
Caragana - Caragan spp.  
  
Cherry - Prunus spp.  
Elm - Ulmus spp.  
Lilac - Syringa spp.  
Maple - Acer spp.  
Mountain Ash - Sorbus spp.  
Poplar - Populus spp.  
Russian Olive - Elaeagnus spp.  
Willow - Salix spp.

#### **Coniferous**

Fir – Abies spp.  
Juniper - Juniperus spp.  
Pine - Pinus spp.  
Spruce - Picea spp.  
Yew - Taxus spp.

**NOTE:** This product is not recommended for use as an over-the-top broadcast spray in forest tree nurseries or in Christmas tree plantations. Application in such sites should be limited to directed sprays. DO NOT treat Christmas tree plantations in the year of anticipated harvest.

### **TREE, VINE and BERRY CROPS**

This product is recommended for annual and perennial weed control in established vineyards or orchards, in blueberry, cranberry and strawberry, or for site preparation prior to transplanting tree and vine crops. Applications may be made with boom equipment, shielded sprayers, hand-held and high volume orchards guns, or with wiper applicator equipment (orchards, vineyards, cranberry and strawberry only). See the "Mixing and Application Equipment Information" section of this label and the following table for specific information on the use of equipment.

Repeat treatments may be necessary to control weeds originating from underground parts of untreated weeds or from seeds. This product does not provide residual or pre-emergent weed control. For subsequent weed control, follow a program using residual herbicides or use repeated applications of this product. DO NOT apply more than 26 litres of this product per hectare per year.

**EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF HERBICIDE SOLUTION, SPRAY,**

2023-2066  
2023-10-24

Proposed Label on 2023-05-08

**DRIFT, OR MIST WITH FOLIAGE OR GREEN BARK OF TRUNK, BRANCHES, SUCKERS, FRUIT, CANES OF BLUEBERRY BUSHES, OR OTHER PARTS OF TREES OR VINES. CONTACT OF THIS PRODUCT WITH OTHER THAN MATURED BROWN BARK CAN RESULT IN SERIOUS CROP DAMAGE.**

Reduced control may result when applications are made to annual or perennial weeds that have been mowed, grazed or cut and have not been allowed to regrow to the recommended stage for treatment.

**WEED CONTROL IN TREE, VINE and BERRY CROPS**

Crop	Rate (L/ha)	PreHarvest Interval (days)	Max. Appl. per Yr.	Weeds Controlled	Comments  (Refer to annual weed control and perennial weed control sections for specific rates for weed control)
Apples Apricot Cherry (Sweet/sour) Peaches Pears Plums	1.69 - 9	30	3	Annual and perennial weeds	
Apples Grapes	<b>Tank Mix</b>  1.69 - 9  +  Simazine  2.0 - 4.5 kg ai/ha		1	Annual and perennial weeds	Will provide seasonlong pre-emergent control  DO NOT apply to coarse, sandy or gravelly soil  Use according to the more restrictive label direction for each product in the mix  DO NOT apply to orchards or vineyards that have been established less than 1 or 3 years, respectively  Simazine rate is equivalent to 2.25 – 5.0 kg/ha Princep Nine-T, or 4.0-9.0 kg/ha Simadex

2023-2066

2023-10-24

Proposed Label on 2023-05-08

Grapes	1.69 – 9	14	3	Annual and perennial weeds	- Remove all sucker growth from the spray zone before spraying, except for the Concord variety of grape - Suckering should be conducted within 2 weeks prior to application DO NOT apply to vines which have been established less than 3 years
--------	----------	----	---	----------------------------	--

Highbush (cultivated) blueberry	2.1 – 4.2	30	1	Quack-grass	- Use as a directed spray, with no more than 275 kPa pressure
Lowbush blueberry	0.75 – 1.5% solution (spot application)	Apply in non-bearing year only	1	Woody brush	- Apply as a directed spray in mid-summer of the vegetative (nonbearing) year - See spot treatment section for instructions
Filberts Hazelnut (established plantations)	1.69 – 2.63	14	-	Annual weeds	- Use as a directed spray, with no more than 275 kPa pressure
Walnut Chestnut Japanese heartnut	1.69 – 9	-	2	Annual and perennial weeds	- Apply late spring and fall, post-harvest but prior to a killing frost - Apply in 200-300 L water as a directed spray, using no more than 275 kPa pressure - Apply alternatively as a 1.5% wiper solution (see Wiper Applications section)

2023-2066

2023-10-24

Proposed Label on 2023-05-08

Cranberry	15% Solution (0.75 L Durango Herbicide + 4L water)	30	1	Annual and perennial weeds	- Apply using wick or wiper applicators
Strawberry	0.75 – 1.5% solution (spot application) 25% solution (wiper application)	30	1	Emerged perennial weeds	- Apply when weeds are at a susceptible growth stage (see perennial weed control section) - See spot treatment section for instructions - See wiper application section for instructions
Sugar beets	0.75 – 1.5% solution (spot application)	Treated crop MUST NOT be harvested	1	Dodder species	- Apply when dodder is vigorously growing but before flowering. - See spot treatment section for instructions.
Asparagus	0.94-1.88	7	1	Fall seeded rye grass	Apply in spring before emergence of crop shoots.



**NOTE TO USER: READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR THE INDICATED SPECIAL USE APPLICATIONS: (NORTH AMERICAN GINSENG).**

The DIRECTIONS FOR USE for this product for the use(s) described on this label were developed by persons other than Albaugh, LLC. and accepted for registration by Health Canada under the User Requested Minor Use Label Expansion program.

Albaugh, LLC. itself makes no representation or warranty with respect to performance (efficacy) or crop tolerance (phytotoxicity) claims for this product when used on the crop(s) listed on this label.

Accordingly, the Buyer and User assume all risks related to performance and crop tolerance arising, and agree to hold Albaugh, LLC. harmless from any claims based on efficacy and/or phytotoxicity in connection with the use(s) described on this label.

**DIRECTIONS FOR USE**

**ALWAYS REFER TO THE PRODUCT LABEL FOR FURTHER INFORMATION ON WEEDS CONTROLLED, APPLICATION DIRECTIONS, AND USE PRECAUTIONS.**

**NORTH AMERICAN GINSENG**

**New Gardens (British Columbia only):** Apply this product in the fall after seeding but before freeze-up in new gardens only to control volunteer cereals. Apply when weeds are at the growth stages listed on the product label. Use a single application of 1.88 litres per hectare in 50 to 100 litres water per hectare. **DO NOT USE A FALL APPLICATION IN ESTABLISHED/EXISTING GARDENS.**

**Existing/Established Gardens:** Apply this product in the spring before the crop has emerged from the soil. Apply when weeds are at the growth stages described in the product label. A maximum of two 1.88 litres per hectare applications in 50 to 100 litres water per hectare may be made in a season. **DO NOT USE A FALL APPLICATION IN ESTABLISHED/EXISTING GARDENS.**

**SELECTIVE EQUIPMENT WIPER APPLICATORS**

This product may be applied with a wiper applicator, after dilution and thorough mixing with water, to listed weeds in soy and dry beans, grapes, orchards, cranberries and strawberry. Applications must be made before initial pod set in soy and dry beans. (It may also be used in any industrial, tree planting and non-crop site specified on this label.)

A wiper applicator applies the herbicide solution onto weeds by rubbing the weed with an absorbent material containing the herbicide solution. Wiper applicators include either roller or wick devices which physically wipe appropriate concentrations or amounts of this product directly onto the weed. Equipment must be designed, maintained and operated to prevent the herbicide solution from contacting desirable vegetation. Performance may be improved by reducing speed in areas of heavy weed infestations to insure adequate wiper saturation. Best results may be obtained if 2 applications are made in opposite directions.

2023-2066

2023-10-24

Proposed Label on 2023-05-08

**AVOID CONTACT WITH DESIRABLE VEGETATION.** Contact of the herbicide solution with desirable vegetation may result in damage or destruction. Applicators used above desired vegetation should be adjusted so that wiper contact point is at least 5 cm above the desirable vegetation. Droplets or foam of the herbicide solution settling on desirable vegetation may result in discoloration, stunting or destruction. Applications should be made when the weeds are a minimum of 15 cm above the desirable vegetation. Best results may be obtained when more of the weed is exposed to the herbicide solution. Weeds not contacted by the herbicide solution will not be affected. This may occur in dense clumps, severe infestations, or when the height of the weeds varies so that not all weeds are contacted. In these instances, repeat treatments may be necessary. See the Weed Control tables in this label for recommended stage of growth for specific weeds.

#### **NOTES**

- Maintain equipment in good operating condition. Avoid leakage or dripping onto desirable vegetation.
- Adjust height of applicator to insure proper contact with weeds.
- Keep wiping surfaces clean.
- Maintain recommended roller RPM on roller applicators while in use.
- Keep wiper material at proper degree of saturation with herbicide solution.
- DO NOT use wiper equipment when weeds are wet.
- DO NOT operate equipment at ground speeds below 4 and greater than 10 km/h. Weed control may be affected by speed of application equipment. As weed density increases, reduce equipment ground speed to insure good coverage of weeds.
- Be aware that on sloping ground the herbicide solution may migrate, causing dripping on the lower end and drying on the upper end of the wiper applicator.
- Variation in equipment design may affect weed control. With wiper applicators, the wiping material and its orientation must allow delivery of sufficient quantities of the recommended herbicide solution directly to the weed.
- Care must be taken with all types of wipers to insure that the absorbent material does not become over-saturated, causing the herbicide to drip onto desirable vegetation.
- With all equipment, drain and clean wiper parts immediately after using this product, by thoroughly flushing with water.

**For Roller Applicators**--Mix 0.38 to 0.75 L of this product in 10 L water to prepare a 3.8 to 7.5% solution. Roller speed should be maintained at 50 to 150 rpm.

**For Wick or other Wiper Applicators**--Mix 1 litre of this product in 3 litres of water to prepare a 25% solution.

#### **NON-CROPLAND USES**

##### **NON-CROPLAND USES INCLUDE:**

Industrial; recreational, rights-of-way, and public areas; turf grass renovation.

2023-2066

2023-10-24

Proposed Label on 2023-05-08

**ALWAYS READ PRECAUTIONARY STATEMENTS, GENERAL INFORMATION and MIXING and APPLICATION SECTIONS PRIOR TO SPECIFIC APPLICATION INFORMATION IN ANY LABEL SECTION.**

This product can be used to control annual and perennial weeds and woody brush and trees listed on this label in non-crop areas such as railroad, pipeline, highway, power and telephone rights-of-way; petroleum tank farms and pumping installations; roadsides; storage areas; lumberyards; fence rows; industrial plant sites; parking areas; school yards, parks, golf courses, other public areas; airports and similar industrial or non-crop areas.

**NOTE:** For all industrial, rights-of-way, recreational and public areas, repeat treatments may be necessary to control regeneration or new growth.

When applied as recommended under the conditions described, this product will control weeds in noncropland areas as listed in the following table.

**WEED CONTROL IN NON-CROPLAND AREAS WITH DURANGO HERBICIDE**

WEEDS	GROUND APPLICATION			COMMENTS
	BOOM APPLICATION		Hand Held High Volume Application % Solution	
	Rate † L / ha	Water Vol. † L /ha		
Annual grasses <u>and</u> <u>broadleaves</u>	1.69 – 2.63	50 - 100	1	.Actively growing weeds
<u>Perennial Weeds</u>				
Quackgrass	1.88	50 - 300	1	.Actively growing weeds .Add 0.5% v/v of a recommended surfactant when using water volumes greater than 150 L.
	3.56 –5.25	50 - 300	2	Higher rate for long term control and for heavy infestations
Canada Thistle (Bud Stage)	3.56 - 5.25	100 - 300	2	.See section on purple loosestrife control for application instructions  .Summer through fall is optimum
Purple loosestrife	4.5	300-600	0.75 – 1.5 (or 25% for wiper application)	
Other Perennials	5.25 – 9	100 –300	2	

2023-2066  
 2023-10-24  
 Proposed Label on 2023-05-08

<u>Brush and Trees</u>  Birch, Cherry, Poplar, Western Snowberry, Willow  Maple, Raspberry/ Salmonberry, Alder	2.25 – 4.5      4.5	100 – 300      100 – 300	1 - 2      2	.Summer through early fall  .Late Summer through fall  .Fall is optimum
<u>Turf Renovation</u>  Annual and Perennial Weeds	1.88 – 9.0	100 – 300	1 - 2	.Use higher end of the rate range for perennials
<u>Roadside Vegetation</u> (1-2 m wide along shoulders)  Annual Weeds (refer to Tank-Mix sections on product labels for specific weeds controlled)	1) 0.56- 0.75 + 1.25-2.5 L DyCleer 480  or  2) 0.56- 0.75 + 0.30 L DyCleer 480 + 1.2 L 2,4-D Amine 500	25-150	-	-Refer to annual weed control table in this label for appropriate product rate for specific weeds  -For 2,4-D amine formulations with a different guarantee, adjust the rate accordingly  -No application to standing water

2023-2066

2023-10-24

Proposed Label on 2023-05-08

<u>Residual Control</u> Annual and Perennial Weeds (the simazine component of this tank mixture will provide season long control of most germinating broadleaf weeds and grasses. It may also provide post-emergent activity on certain annual weeds)	1.88 – 9 + a) 2.5-5.6 kg Simazine 80W or + b) 4.0-9.0 L Simadex Flowable	200-400	-	-DO NOT apply to coarse, sandy or gravelly soil. One application per year.  -Use according to the most restrictive label directions for each product in the mixture.  -For other simazine formulations registered for industrial/non-cropland areas, use equivalent rates; i.e. 2.0-4.5 kg simazine/ha
--	---	---------	---	--

† For more information on rates, water volumes and application, refer to the "Annual and Perennial Weed Control" sections of this booklet.

## APPLICATION INFORMATION FOR NON-CROPLAND USES

### Foliar Applications

Spray coverage should be uniform and complete. DO NOT spray to the point of runoff. DO NOT allow spray drift to contact desirable vegetation as severe injury or destruction may occur. For woody brush and trees, early season applications may take 30-45 days for symptoms to develop on target species. Late season application may be made to species that have some autumn colours provided no major leaf drop has occurred. Control will be observed the following spring.

**EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF SPRAY WITH FOLIAGE OF DESIRABLE TURFGRASSES, TREES, SHRUBS, OR OTHER DESIRABLE VEGETATION SINCE SEVERE DAMAGE OR DESTRUCTION MAY RESULT.**

This product does not provide residual weed control. For subsequent weed control, follow a label approved herbicide program. Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used.

### GROUND APPLICATIONS: For all non-cropland uses

For woody brush and trees, apply 2.25 to 4.5 L of this product per hectare. Use ground boom or boomless, or mist blower equipment, or apply as a 0.75 to 1.5% solution using hand-held high volume equipment. Apply as directed in the recommended volume of clean water to foliage of actively growing vegetation. Use the 4.5 L/ha rate for Maple, Alder and Willow† species, as well as for hard to control perennial weed species. († Suppression only)

Spray coverage should be uniform and complete. DO NOT spray to the point of runoff. DO NOT allow spray drift to contact desirable vegetation as severe injury or destruction may occur. If weeds have been mowed or tilled, DO NOT treat until regrowth has reached the recommended stages.

DO NOT APPLY BY AIR.

### PURPLE LOOSESTRIFE CONTROL

- DO NOT TREAT PLANTS OVER OPEN WATER. Durango Herbicide is not registered for direct application to bodies of water.

2023-2066

2023-10-24

Proposed Label on 2023-05-08

- Treat when plants are actively growing at or beyond the bloom stage. If using hand-held equipment, spray-to-wet.
- For wiper applications, see specific section on this label.
- Where feasible, remove flower heads before treatment to ensure prevention of seed set.
- For large (>1.6 ha) monocultures of loosestrife, work from the periphery inward in successive years to allow competing vegetation to invade the treated area.
- A long-term control strategy should include measures to control both established plants and seedlings. Sprayed areas should be monitored to determine the appropriate follow-up management. Early detection and treatment of second and third generation seedlings is important to prevent reinfestation of purple loosestrife. Desirable native plant communities will then have a chance to become re-established.

### **Selective Application for All Non-Cropland Uses**

Selective equipment such as **WIPER** and **ROLLER** applicators can be used to control emerged weeds in non-crop areas and tree plantings. See "Selective Equipment" for more information.

### **Turfgrass**

When applied as directed, under conditions described, this product controls most existing vegetation. Apply this product at rates specified in the "Weed Control in Non-Cropland Areas" section

**DO NOT DISTURB SOIL OR UNDERGROUND PLANT PARTS BEFORE TREATMENT.**

Where existing vegetation is growing in a field or unmowed situation, apply this product to actively growing weeds at the stages of growth given in the "Weeds Controlled" section of this booklet. Where existing vegetation is growing under mowed turfgrass management, apply this product after omitting at least one regular mowing to allow sufficient growth for good interception of the spray and proper translocation into underground plant parts. Tillage or renovation techniques such as vertical mowing, coring or slicing should be delayed for 7 days after application to allow proper translocation into underground plant parts.

For maximum control of existing vegetation, delay establishment to determine if regrowth from escaped underground plant parts occurs. When repeat treatments are necessary, sufficient regrowth must be attained prior to application. Desirable turfgrasses may be established following the above procedures.

### **Injection Applications** - for all non-cropland uses

Woody vegetation may be controlled by injection application of this product. Apply using suitable equipment, which must penetrate into living tissue, at a rate of at least 0.38 mL (either undiluted or 1:1 with water) per 5 cm tree diameter at breast height (DBH). The cuts should be spaced evenly around the tree and below all major branches. Application may be made at any time of year, except when cold temperatures prevent adequate penetration of injection equipment, or in the spring during periods of heavy sap flow. Control of tree species with tree diameters greater than 20 cm may not be acceptable at this rate.

Total control may not be evident for 1-2 years following treatment.

#### **ALDER**

Alnus spp.

#### **BIRCH**

Betula spp.

#### **CEDAR**

Thuja spp.

#### **CHERRY**

Prunus spp.

#### **DOUGLAS FIR**

#### **HEMLOCK**

Tsuga spp.

#### **MAPLE†**

Acer spp.

#### **PINE**

Pinus spp.

#### **POPLAR**

Populus

spp.

2023-2066  
 2023-10-24  
 Proposed Label on 2023-05-08  
 Pseudotsuga spp.

**WILLOW** A list of species controlled includes:  
 Salix spp.

†This treatment may only provide suppression of Big-Leaf Maple. Late fall applications will provide optimum suppression of Big- Leaf Maple

**Cut Stump Application**

Woody vegetation may be controlled by the application of this product to freshly cut stumps to prevent regrowth. Because the treatment uses a concentrated solution, application must be made using low-pressure equipment, e.g. squirt bottle or similar device. This product must be applied immediately to the surface of the freshly cut stump i.e. within 5 minutes for optimum control at the prescribed rates. Only the cambial tissues of the cut surface should be treated. Apply the herbicide solution at a rate equivalent to at least 0.38 mL product for every 5 cm DBH.

DO NOT cover the remaining area nor any exposed roots, as this product does not penetrate bark well. This treatment may be used at any time of year, except during periods of heavy sap flow or when low temperatures prevent solution application due to freezing. A water soluble colourant may be added to the solution as a means of indicating which surfaces have been treated. Total control may not be evident until 1-2 years after treatment.

See the “Injection Applications” section of this label for the list of species controlled.

**Buffer Zones**

Use of the following spray methods or equipment **DO NOT** require a buffer zone: hand-held or backpack sprayer and spot treatment, inter-row hooded sprayer, low-clearance hooded or shielded sprayers that ensure spray drift does not come in contact with orchard crop fruit or foliage, soil drench and soil incorporation.

For application to rights-of-way and for forestry uses, buffer zones for protection of sensitive terrestrial habitats are not required; however, the best available application strategies which minimize off-site drift, including meteorological conditions (for example, wind direction, low wind speed) and spray equipment (for example, coarse droplet sizes, minimizing height above canopy), should be used. Applicators must, however, observe the specified buffer zones for protection of sensitive aquatic habitats.

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands) and sensitive aquatic habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs, wetlands and estuarine/marine water bodies).

Method of Application	Crop	Maximum number of applications	Buffer Zones (metres) required for protection of:	
			Aquatic Habitat	Terrestrial Habitat
<b>Agricultural and non-cropland systems</b>				

Agricultural crop system and ground boom application method	Rye, cranberry, pasture, summer fallow, all other crops for pre-seeding treatments only, filberts or hazelnut pre-seeding only, ginseng new garden, asparagus		1	1	1
	Ginseng - existing established garden, Canola – Roundup Ready hybrid for seed production		2	1	1
	Filberts or hazelnut		4	1	1
	Corn (glyphosate non-tolerant varieties including grain, silage and ornamental types), sugar beet (glyphosate non-tolerant varieties), strawberry, blueberry highbush and lowbush, walnut, chestnut, Japanese heartnut, turfgrass (prior to establishment or renovation)		2	1	2
	Wheat, barley, oats, soybean (glyphosate non-tolerant varieties), corn-sweet (glyphosate tolerant varieties), canola (glyphosate non-tolerant varieties), peas, dry beans, flax (including low linoleic acid varieties), lentils, chickpea, , corn (glyphosate tolerant varieties), forage grasses and legume including seed production		3	1	2
	Canola (glyphosate tolerant varieties), soybean (glyphosate tolerant varieties)		4	1	2
	Apple, apricot, cherry (sweet/sour), peaches, pears, plums, grapes		3	1	3
Agricultural crop system and airblast application method (including mist blower)	Pasture		1	20	30
	Turfgrass (Prior to establishment or renovation)		2	25	35
Non-cropland system and ground boom application method	Non-crop land and industrial uses: Industrial and rights of way areas, Recreational and public areas		3	1	3*
Non-cropland system and airblast application method (including mist blower)	Non-crop land and industrial uses: Industrial and rights of way areas, Recreational and public areas		3	1	30*
Agricultural crop system and aerial application method	Rye, corn (glyphosate nontolerant varieties), corn-sweet (glyphosate tolerant varieties), chickpea, all other crops for pre-seeding treatments only	Fixed and rotary wing	1	15	20



	Canola (glyphosate tolerant varieties)	Fixed and rotary wing	3	20	40
	Wheat, barley, oats, soybean (glyphosate non-tolerant varieties), canola (glyphosate non-tolerant varieties), peas, dry beans, flax (including low linoleic acid varieties), lentils	Fixed wing	2	20	35
		Rotary wing	2	20	30
	Forage grasses and legume including seed production	Fixed and rotary wing	1	20	40
	Soybean (glyphosate tolerant varieties)	Fixed wing	3	20	45
		Rotary wing	3	20	40
	Summer fallow	Fixed wing	1	20	45
		Rotary wing	1	20	40
	Corn (glyphosate tolerant varieties)	Fixed wing	2	20	50
		Rotary wing	2	20	45
	Pasture	Fixed wing	1	30	70
		Rotary wing	1	30	55
Non-cropland system and aerial application method	Non-crop land and industrial uses: rights-of way areas only	Fixed wing	3	100	NR
		Rotary wing	3	60	NR

\* Buffer zones for the protection of terrestrial habitats are not required for forestry uses or for use on rights-of-way including railroad ballast, rail and hydro rights-of-way, utility easements and roads.

NR = Not Required

For tank mixes, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Buffer Zone Calculator on the Pest Management Regulatory Agency web site.

#### **RESISTANCE MANAGEMENT RECOMMENDATIONS**

For resistance management, Durango Herbicide is a Group 9 herbicide. Any weed population may contain or develop plants naturally resistant to Durango Herbicide and other Group 9 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

2017-3496

2018-03-16

Page 58

- Where possible, rotate the use of Durango Herbicide or other Group 9 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(S) as effectively as the more resistance-prone partner.
- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor treated weed populations after herbicide application for signs of resistance development. (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.

For further information or to report suspected resistance, contact Albaugh, LLC

- . at 1-800-247-8013.

**NOTICE TO USER:** This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

All products listed are registered trademarks of their respective companies.