

CONTAINER LABEL FOR BOTTLE

Group	2	Herbicide
-------	----------	-----------

EVEREST® 70 WDG Herbicide
70% WATER DISPERSIBLE GRANULAR HERBICIDE

AGRICULTURAL

FOR PREPLANT, PREEMERGENCE AND POST-EMERGENCE APPLICATION ON SPRING WHEAT (EXCLUDING DURUM WHEAT), AND FOR POST-EMERGENCE APPLICATION ON SPRING WHEAT (INCLUDING DURUM WHEAT) FOR CONTROL OF WILD OAT, GREEN FOXTAIL, VOLUNTEER TAME OAT AND CERTAIN BROADLEAF WEEDS

FOR SALE AND USE IN MANITOBA, SASKATCHEWAN, ALBERTA AND PEACE RIVER REGION OF BRITISH COLUMBIA ONLY

ACTIVE INGREDIENT:

Flucarbazone..... 66%
(present as flucarbazone-sodium)

REGISTRATION NUMBER 26447

PEST CONTROL PRODUCTS ACT

READ THE LABEL AND BOOKLET BEFORE USING

Warning, contains the allergen milk

KEEP OUT OF REACH OF CHILDREN

STORE IN A COOL, DRY PLACE

NET CONTENTS: 300 GRAMS to 600 kg

For product information call: 1-800-438-6071

UPL AgroSolutions Canada Inc.
630 Freedom Business Center, Suite 402
King of Prussia, PA 19406

2021-3971
2021-12-02

FOR CHEMICAL EMERGENCY: spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on this label. It is an offence under the *PEST CONTROL PRODUCTS ACT* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

PROTECTIVE CLOTHING

When handling (mixing and loading) EVEREST 70 WDG Herbicide, wear long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes.

Read and follow handling precautions and protective clothing recommendations on broadleaf herbicide and surfactant tank-mix partner labels. Follow manufacturers instructions for cleaning/maintaining PPE (Personal Protective Equipment). If no such instructions for washables, use detergent and hot water. Keep PPE apart from other laundry and wash separately.

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove personal protective equipment immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Note: If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at: www.croplife.ca/.

FIRST AID

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If Inhaled: Move person to fresh air. If person is not breathing, call 911 or an

2021-3971

2021-12-02

ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

No specific antidote is available. Treat the patient symptomatically.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, or feed. Store in original container and out of reach of children, preferably in a locked storage area. EVEREST 70 WDG Herbicide is not affected by freezing.

SPILL CLEAN-UP: Follow safety precautions as directed for handling the product. Sweep spilled product with a broom and shovel into tightly closable container. If on a floor or hard surface, wash the surface or floor with detergent and water, then rinse. If on soil, collect surface soil contaminated with the product. If the product is clean, it may be used; otherwise, follow the DISPOSAL instructions.

DISPOSAL/DECONTAMINATION:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

<p>FOR CHEMICAL EMERGENCY: spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.</p>
--

BOOKLET LABEL FOR BOTTLE

Group	2	Herbicide
-------	----------	-----------

EVEREST® 70 WDG Herbicide
70% WATER DISPERSIBLE GRANULAR HERBICIDE

AGRICULTURAL

FOR PREPLANT, PREEMERGENCE AND POSTEMERGENCE APPLICATION ON SPRING WHEAT (EXCLUDING DURUM WHEAT), AND FOR POSTEMERGENCE APPLICATION ON SPRING WHEAT (INCLUDING DURUM WHEAT) FOR CONTROL OF WILD OAT, GREEN FOXTAIL, VOLUNTEER TAME OAT AND CERTAIN BROADLEAF WEEDS

FOR SALE AND USE IN MANITOBA, SASKATCHEWAN, ALBERTA AND PEACE RIVER REGION OF BRITISH COLUMBIA ONLY

ACTIVE INGREDIENT:

Flucarbazone..... 66%
(present as flucarbazone-sodium)

REGISTRATION NUMBER 26447

PEST CONTROL PRODUCTS ACT

READ THE LABEL AND BOOKLET BEFORE USING

Warning, contains the allergen milk.

KEEP OUT OF REACH OF CHILDREN

STORE IN A COOL, DRY PLACE

NET CONTENTS: 300 GRAMS to 600 Kg

For product information call: 1-800-438-6071

UPL AgroSolutions Canada Inc.
630 Freedom Business Center, Suite 402
King of Prussia, PA 19406

FOR CHEMICAL EMERGENCY: spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *PEST CONTROL PRODUCTS ACT* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

PRODUCT INFORMATION

EVEREST 70 WDG Herbicide is a selective herbicide for use at preplant, preemergence and postemergence timings in spring wheat (hard red spring, Canada Prairie spring, soft white spring, and extra strong or utility) and at postemergence timings only in durum wheat.

EVEREST 70 WDG Herbicide applied post emergence must be tank mixed with a surfactant listed on this label. For broader spectrum activity, EVEREST 70 WDG Herbicide may be tank mixed with a broadleaf herbicide listed on this label. See TANK MIXES section for recommended products.

EVEREST 70 WDG Herbicide is absorbed by foliage and roots of wild oat, green foxtail and certain broadleaf weeds. These weeds cease growth soon after application, removing the competitive effects of susceptible weeds. However complete weed control may not be seen for one to two weeks.

EVEREST 70 WDG Herbicide will control wild oat biotypes which have developed resistance to ACCase (Group 1) and triallate (Group 8) herbicides. It will also control green foxtail biotypes resistant to ACCase (Group 1) and dinitroaniline (Group 3) herbicides.

One 567 gram jug contains enough EVEREST 70 WDG Herbicide to treat 13-26 hectares (33-66 acres) depending on the target weeds to be controlled. For weeds controlled and staging of application for each rate refer to the recommended application tables.

567 g Jug Treats the Following:		
Use Rate	Acres	Hectares
21.5 g/ha	66	26
28.5 g/ha	50	20
35.7 g/ha	40	16
43.0 g/ha	33	13

DIRECTIONS FOR USE

Read the entire DIRECTIONS FOR USE before using EVEREST 70 WDG Herbicide.

- **Ground Application only. DO NOT APPLY BY AIR.**
- Do not mix, load or clean spray equipment within 10 metres of well-heads or aquatic systems, including marshes, ponds, ditches, streams, lakes, etc.
- Buffer Zones: Overspray or drift to sensitive habitats should be avoided. A buffer zone of 20 metres is required between the downwind edge of the boom and the closest edge of sensitive terrestrial habitats including forested areas, shelter belts, woodlots, hedgerows, and shrublands. A buffer zone of 35 metres is required between the downwind edge of the boom and the closest edge of sensitive aquatic habitats including sloughs, coulees, ponds, prairie potholes, lakes, rivers, streams, wetlands, and wildlife habitat at the edge of these bodies of water. Do not contaminate these habitats when cleaning and rinsing spray equipment or containers.
- Consult the label of the tank mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture.
- Do not apply during periods of dead calm, when winds are gusty or when wind speed is greater than 15 km/hr at 2 metres high above ground at the site of application.
- **Do not apply postemergence treatments if it is raining or if rainfall is expected within one (1) hour after application.**
- Do not graze treated fields or use green crop for feed. Wheat grain or straw from harvested treated fields may be fed to livestock.
- Do not allow this chemical to drift on to other crops, especially canola, tame oats, or other non target crops.
- Observe minimum interval to harvest of 80 days after treatment.
- Do not treat wheat underseeded to legumes.
- The maximum rate of glyphosate applied when tank mixed with Everest 70 WDG Herbicide for pre-plant, pre-emergence is 450 g ae/ha.

TIMING OF APPLICATION

EVEREST 70 WDG Herbicide can be applied up to 1 week prior to planting, or immediately after planting prior to crop emergence in spring wheat (excluding durum wheat), and postemergence in all types of spring wheat (including durum wheat).

Do not apply EVEREST 70 WDG Herbicide before crop emergence is complete in durum wheat.

Research has demonstrated that optimum wheat yield is obtained by early removal of target weeds prior to tillering of wheat. In postemergence use, apply EVEREST 70 WDG Herbicide to wheat that has a minimum of one leaf to a maximum of four leaves on the main stem, plus two tillers. Application beyond this stage may result in yield losses due to weed competition.

PREPLANT, POSTPLANT PREEMERGENCE AND PREPLANT OR POSTPLANT PREEMERGENCE FOLLOWED BY POSTEMERGENCE USE DIRECTIONS FOR SPRING WHEAT (EXCLUDING DURUM WHEAT)

Use of EVEREST 70 WDG Herbicide as a preplant or postplant preemergence application only or in combination with an additional postemergence application of EVEREST 70 WDG allows for early weed removal and reduced weed competition with wheat.

Do not apply more than 43 g/ha of EVEREST 70 WDG Herbicide (30 g ai/ha) per growing season.

Application Timing	Application Rate (g product/ha)	Weeds Controlled or Suppressed
Preplant or postplant preemergence ¹	21.5	Control: volunteer canola excluding imidazolinone-tolerant (Clearfield) canola varieties Suppression: wild oat
	28.5	Weeds controlled or suppressed with 21.5 g/ha plus Suppression: wild mustard, green foxtail
Preplant or postplant preemergence followed by Postemergence ^{2,3} Everest 70 WDG Herbicide	14.5 + 28.5	Control: green foxtail, wild oat, volunteer canola excluding imidazolinone-tolerant (Clearfield) canola varieties
	21.5 + 21.5	
	28.5 + 14.5	

¹Crop injury may be observed following preplant or postplant preemergence applications, but yield should be unaffected.

² For postemergence applications, use a registered non-ionic surfactant such as Agral 90, Ag-Surf, etc. at a rate of 0.25% v/v (0.25L per 100L total spray solution) according to use directions under POSTEMERGENCE USE DIRECTIONS on the Everest 70 WDG Herbicide label.

³ For tank mix directions and control of other weeds using postemergence applications, refer to the POSTEMERGENCE USE DIRECTIONS on the Everest 70 WDG Herbicide label.

GLYPHOSATE TANK MIXTURE

Everest 70 WDG Herbicide may be tank mixed with glyphosate present as the isopropylamine or potassium salt when applied preplant or postplant preemergence. Refer to the appropriate product label for use directions.

POSTEMERGENCE USE DIRECTIONS IN SPRING AND DURUM WHEAT

RECOMMENDED APPLICATIONS

CROP APPLICATION TIMING		
CROP	GROWTH STAGE	REMARKS
Spring Wheat Durum Wheat	1 leaf to 6 total leaves	1 leaf to 4 leaves on main stem, plus 2 tillers

GRASS WEED CONTROL IN SPRING AND DURUM WHEAT			
CROP	RATE	WEEDS	REMARKS
Spring Wheat Including: hard red spring, Canada Prairie spring, soft white spring and extra strong (utility) wheat Durum Wheat	21.5 g/ha	Green Foxtail (Wild millet)	Green foxtail should be no larger than 4-leaf, 2 tillers.
	28.5 g/ha	Weeds controlled at 21.5 g/ha plus Wild Oat Volunteer Tame Oat	Wild oat should be no larger than 4-leaf, 2 tillers. Use 28.5 g/ha when wild oat infestations are below 100 plants/m ² . Volunteer tame oat should be no larger than 4-leaf, 2 tillers.
	35.7 g/ha	Weeds controlled at 28.5 g/ha plus Wild Oat	Wild oat should be no larger than 4-leaf, 2 tillers. Use 35.7 g/ha when wild oat infestations are greater than 100 plants/m ² and plants are actively growing under ideal environmental conditions.
	43 g/ha	Weeds controlled at 35.7 g/ha plus Wild Oat	Wild oat should be no larger than 4-leaf, 2 tillers. Use 43 g/ha when wild oat infestations are greater than 100 plants/m ² and when plants are not actively growing due to poor environmental conditions or when tank mixing with a product that contains dicamba. [Refer to the TANK MIXES section listed on this label.]

BROADLEAF WEED CONTROL IN SPRING AND DURUM WHEAT		
WEED	RATE	REMARKS
Redroot Pigweed (<i>Amaranthus retroflexus</i>)	28.5 g/ha	For control of redroot pigweed, apply EVEREST 70 WDG Herbicide to weeds at the 2 to 6 leaf stage.
Wild Mustard (<i>Brassica kaber</i>)		For control of wild mustard apply EVEREST 70 WDG Herbicide to weeds at the 2 to 6 leaf stage

2021-3971
2021-12-02

Stinkweed (<i>Thlaspi arvense</i>)		For control of stinkweed apply EVEREST 70 WDG Herbicide to weeds at the 2 to 9 leaf stage
Volunteer Canola (<i>Brassica napus</i>)		For control of volunteer canola apply EVEREST 70 WDG Herbicide to weeds at the 2 to 6 leaf stage Will not control imidazolinone-tolerant (CLEARFIELD®) canola varieties
Green Smartweed (<i>Polygonum scabrum</i> Moench.)		For control of green smartweed apply EVEREST 70 WDG Herbicide to weeds at the 2 to 6 leaf stage
Shepherd's Purse (<i>Capsella bursa-pastoris</i>)		For control of shepherd's purse, apply EVEREST 70 WDG Herbicide to weeds at the 2 to 6 leaf stage.

Wheat exposed to water-logged or saturated soils, or temperature extremes such as heat or freezing weather, or drought, low fertility or plant disease at application time could show unacceptable injury symptoms. Weed control also may be reduced by these same conditions.

TANK MIXES

Use 28.5 g/ha when wild oat populations are below 100 plants/ m² and when plants are actively growing. For broader spectrum control of both annual grasses and broadleaf weeds, EVEREST 70 WDG Herbicide may be mixed with one of the broadleaf herbicides plus a surfactant listed in the following table. With all tank mix partners, read and follow the use directions, rates, precautions, timing, recropping restrictions, grazing interval restrictions and recommendations on the broadleaf herbicide and surfactant labels.

APPROVED TANK MIX PARTNERS IN SPRING WHEAT						
BROADLEAF HERBICIDE TANK MIX PARTNER	BROADLEAF HERBICIDE PRODUCT RATES	EVEREST 70 WDG HERBICIDE RATE			SURFACTANTS (0.25% v/v or 0.25 L/100L)	
		WILD OAT AND GREEN FOXTAIL	WILD OAT ONLY	GREEN FOXTAIL ONLY***		
1	2,4-D Amine	up to 560 g ae/ha	43 g/ha	28.5 g/ha	21.5 g/ha	Surf 92, Super Spreader, LI 700® Penetrating Surfactant, Agral® 90, Ag-Surf
2	2,4-D Ester	up to 560 g ae/ha	43 g/ha	28.5 g/ha	21.5 g/ha	Surf 92, Super Spreader, LI 700® Penetrating Surfactant, Agral® 90, Ag-Surf
3	**Ally® + 2,4-D Amine	5 or 7.5 g/ha Ally + up to 560 g ae/ha 2,4-D Amine	***43 g/ha	***28.5 g/ha	21.5 g/ha	Agral® 90, Ag-Surf
4	**Ally® + 2,4-D Ester	5 or 7.5 g/ha Ally + up to 560 g ae/ha 2,4-D Ester	***43 g/ha	***28.5 g/ha	21.5 g/ha	Agral® 90, Ag-Surf
5	Attain® Herbicide Tank Mix	0.6 L/ha Attain A + 1 L/ha Attain B	***43 g/ha	***28.5 g/ha	21.5 g/ha	Agral® 90, Ag-Surf
6	Buctril® M	1 L/ha	43 g/ha	28.5 g/ha	21.5 g/ha	Surf 92, Super Spreader, LI 700® Penetrating Surfactant, Agral® 90, Ag-Surf
7	Curtail® M	1.5 – 2.0 L/ha	***43 g/ha	***28.5 g/ha	21.5 g/ha	Agral® 90, Ag-Surf
8	DyVel®	1.25 L/ha	43 g/ha	Do Not Use	21.5 g/ha	Agral® 90, Ag-Surf

2021-3971

2021-12-02

9	*DyVel® DS	850 mL/ha	43 g/ha	Do Not	21.5	Agral® 90,
---	------------	-----------	---------	--------	------	------------

APPROVED TANK MIX PARTNERS IN SPRING WHEAT

	BROADLEAF HERBICIDE TANK MIX PARTNER	BROADLEAF HERBICIDE PRODUCT RATES	EVEREST 70 WDG HERBICIDE RATE			SURFACTANTS (0.25% v/v or 0.25 L/100L)
			WILD OAT AND GREEN FOXTAIL	WILD OAT ONLY	GREEN FOXTAIL ONLY***	
				Use	g/ha	Ag-Surf
10	Estaprop® Dichlorprop-D	1.75 L/ha	43 g/ha	28.5 g/ha	21.5 g/ha	Surf 92, Super Spreader, LI 700® Penetrating Surfactant, Agral® 90, Ag-Surf
11	Express Pack®	10 g/ha Express + 0.625 L/ha 2,4-D	43 g/ha	28.5 g/ha	21.5 g/ha	Agral® 90, Ag-Surf
12	Frontline™ 2,4-D Herbicide Tank-Mix	100 ml/ha Frontline 2,4-D A + 1.0 L/ha Frontline 2,4-D B	43 g/ha	28.5 g/ha	21.5 g/ha	Agral® 90, Ag-Surf
13	Frontline™ Herbicide Tank-Mix	100 ml/ha Frontline A + 0.84 L/ha Frontline B	43 g/ha	28.5 g/ha	21.5 g/ha	Agral® 90, Ag-Surf
14	MCPA Amine	up to 560 g ae/ha	43 g/ha	28.5 g/ha	21.5 g/ha	Agral® 90, Ag-Surf
15	MCPA Ester	up to 560 g ae/ha	43 g/ha	28.5 g/ha	21.5 g/ha	Agral® 90, Ag-Surf
16	Pardner®	1 L/ha	43 g/ha	Do Not Use	21.5 g/ha	Agral® 90, Ag-Surf
17	Prestige® Herbicide Tank-Mix	0.8 L/ha Prestige A + 2.0 L/ha Prestige B	***43 g/ha	***28.5 g/ha	21.5 g/ha	Agral® 90, Ag-Surf
18	** Refine Extra® + 2,4-D Amine	20 g/ha Refine Extra + up to 560 g ae/ha 2,4-D Amine	43 g/ha	28.5 g/ha	21.5 g/ha	Surf 92, Super Spreader, LI 700® Penetrating Surfactant, Agral® 90,Ag-Surf
19	** Refine Extra® + 2,4-D Ester	20 g/ha Refine Extra + up to 560 g ae/ha 2,4-D Ester	43 g/ha	28.5 g/ha	21.5 g/ha	Agral® 90, Ag-Surf
20	* Target®	1 to 1.5 L/ha	***43 g/ha	Do Not Use	21.5 g/ha	Agral® 90, Ag-Surf
21	Spectrum™ Herbicide Tank-Mix	100 mL/ha Spectrum A + 1.5 L/ha Spectrum B	43 g/ha	28.5 g/ha	21.5 g/ha	Agral® 90, Ag-Surf

22	Thumper®	1 L/ha	43 g/ha	28.5 g/ha	21.5 g/ha	Surf 92, Super Spreader, LI 700® Penetrating Surfactant, Agral® 90, Ag-Surf
23	** Unity® Herbicide	0.5 L/ha Unity EC + 10.6	***43	Do Not	21.5	Agral® 90,

APPROVED TANK MIX PARTNERS IN SPRING WHEAT

BROADLEAF HERBICIDE TANK MIX PARTNER	BROADLEAF HERBICIDE PRODUCT RATES	EVEREST 70 WDG HERBICIDE RATE			SURFACTANTS (0.25% v/v or 0.25 L/100L)
		WILD OAT AND GREEN FOXTAIL	WILD OAT ONLY	GREEN FOXTAIL ONLY***	
Tank Mix	g/ha Unity 75 WG	g/ha	Use	g/ha	Ag-Surf

APPROVED TANK MIX PARTNERS IN DURUM WHEAT

BROADLEAF HERBICIDE TANK MIX PARTNER	BROADLEAF HERBICIDE PRODUCT RATES	EVEREST 70 WDG HERBICIDE RATE***			SURFACTANTS (0.25% v/v or 0.25 L/100L)	
		WILD OAT AND GREEN FOXTAIL	WILD OAT ONLY	GREEN FOXTAIL ONLY		
1	2,4-D Amine	420 g ae/ha	43 g/ha	28.5 g/ha	21.5 g/ha	Agral® 90, Ag-Surf
2	2,4-D Ester	420 g ae/ha	43 g/ha	28.5 g/ha	21.5 g/ha	Agral® 90, Ag-Surf
3	Frontline™ Herbicide Tank-Mix	100 mL/ha Frontline A + 0.84 L/ha Frontline B	43 g/ha	28.5 g/ha	21.5 g/ha	Agral® 90, Ag-Surf
4	Spectrum™ Herbicide Tank-Mix	100 mL/ha Spectrum A + 1.5 L/ha Spectrum B	43 g/ha	28.5 g/ha	21.5 g/ha	Agral® 90, Ag-Surf

NOTE:

- * Reduction in wild oat control may be observed with this tank mix partner.
- ** Addition of a second surfactant is not required for EVEREST 70 WDG Herbicide tank mixes with the herbicides Ally, Refine Extra and Unity Herbicide Tank Mix. Use only one of the recommended surfactants from the EV EREST 70 WDG Herbicide label at an application rate of 0.25% v/v (0.25 L per 100 L total spray solution).
- *** Apply specified rate in at least 100 L of water per hectare (45 L/A or 10 gallons/A).

MIXING INSTRUCTIONS

Ensure the spray tank is clean. In-line strainers and nozzle screens should be clean and 50-mesh or coarser.

1. Fill the spray tank 1/3 to 1/2 full with clean water and begin agitation or bypass.
2. Add the appropriate EVEREST 70 WDG Herbicide rate directly to the spray tank. Vigorous agitation is required to dissolve the EVEREST 70 WDG Herbicide. Maintain sufficient agitation during both mixing and application.
3. Add the broadleaf weed herbicide.
4. Add the surfactant, then complete filling tank with balance of water needed.

2021-3971
2021-12-02

SPRAYING INSTRUCTIONS

Unless otherwise specified in the tank mix partner section of this label, apply in a spray volume of 50 – 100 L/ ha (22.5 – 45 L/A or 5 – 10 gallons/A) at 207-345 kPa (30-50 PSI) pressure to ensure proper weed coverage.

Flat fan nozzles of 80 ° or 110 ° are recommended for optimum coverage. Do not use floodjet nozzle or control droplet application equipment. Nozzles may be oriented 45 ° forward to enhance crop penetration and to give better weed coverage.

SPRAYER CLEAN-UP

1. Drain the tank and thoroughly rinse spray tank, boom and hoses with clean water. Pay particular attention to flushing out any visible deposits.
2. Fill the tank with clean water and 1% v/v (1 L/100L) household ammonia. Flush the hoses, boom and nozzles with the cleaning solution. Circulate for at least 15 minutes. Flush hoses, boom and nozzles once more, then drain the tank;
3. Clean nozzles and screens in a separate container using the ammonia and water;
4. Repeat #2;
5. Rinse tank, boom and hoses with clean water.
 - Do not clean sprayer near desirable vegetation, wells, or other water sources;
 - Dispose of all rinsings in accordance with provincial regulations;
 - Check tank mix partner label for any additional clean-up procedures.

ROTATIONAL CROPS

The following crops may be planted 11 months after an application of EVEREST 70WDG Herbicide.

Soil Zones and Rotational Crops			
Gray-Wooded	Black	Dark Brown	Brown
Spring wheat Barley Canola Field peas*	Spring wheat Barley Canola Field peas* Field bean Flax Durum wheat	Spring wheat Barley Canola Field peas* Flax Durum wheat	Spring wheat

* Field peas can be successfully grown the year following an Everest application providing the following three criteria are all met.

1. Soil pH must be below 7.5
2. Organic Matter content must be above 4%.
3. Precipitation must be equal to or above the 10 year average (minimum 1 00mm within 60 days of application in year of application).

Or, fields can be summer fallowed.

As Everest 70 W DG Herbicide is degraded by soil microbes, environmental conditions that decrease microbial activity must be considered when making rotational cropping decisions. These environmental conditions include prolonged drought and/or cold temperatures within the following cropping season, as well as soils with both low O M (less than 2%) and high pH (greater than 7.5). If these conditions exist, a soil bioassay may be necessary to ensure rotational crop safety.

Do not plant crops other than those listed above in the year following application of

EVEREST 70 WDG Herbicide on wheat.

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, EVEREST 70 WDG Herbicide is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to EVEREST 70 WDG Herbicide and other Group 2 wild oat herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of EVEREST 70 WDG Herbicide or other Group 2 wild oat herbicides with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted.
- Herbicide use should be based on an IPM program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical), cultural, biological and other chemical control practices.
- Monitor treated weed populations for resistance development.
- Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment and planting clean seed.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact UPL AgroSolutions Canada Inc at 1-800-438-6071.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

PROTECTIVE CLOTHING

When handling (mixing and loading) EVEREST 70 WDG Herbicide, wear long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes.

Read and follow handling precautions and protective clothing recommendations on broadleaf herbicide and surfactant tank-mix partner labels. Follow manufacturer's instructions for cleaning/maintaining PPE (Personal Protective Equipment). If no such instructions for washables, use detergent and hot water. Keep PPE apart from other laundry and wash separately.

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove personal protective equipment immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Note: If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at: www.croplife.ca/.

FIRST AID

If swallowed: Call a poison control centre immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

No specific antidote is available. Treat the patient symptomatically.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, or feed. Store in original container and out of reach of children, preferably in a locked storage area. EVEREST 70 WDG Herbicide is not affected by freezing.

SPILL CLEAN-UP: Follow safety precautions as directed for handling the product. Sweep spilled product with a broom and shovel into tightly closable container. If on a floor or hard surface, wash the surface or floor with detergent and water, then rinse. If on soil, collect surface soil contaminated with the product. If the product is clean, it may be used; otherwise, follow the DISPOSAL instructions.

DISPOSAL/DECONTAMINATION:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.

2021-3971
2021-12-02

3. Make the empty container unsuitable for further use.

2021-3971
2021-12-02

4. Dispose of the container in accordance with provincial requirements.
5. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

**FOR CHEMICAL EMERGENCY: spill, leak, fire, exposure, or
accident call CHEMTREC 1-800-424-9300.**

UPL, the UPL logo and EVEREST are registered trademarks of a UPL Corporation Limited Group Company.

©2021 UPL Corporation Limited Group Company.

® All other products are registered trademarks of their respective companies.

™ All other products are trademarks of their respective companies.

2021-3971
2021-12-02

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.