

2022-0231

2022-01-31

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41

Advantage™ II Medium Dog
Insert (English)

<Elanco Logo>

Advantage™ II medium dog

Flea Adulticide/Larvicide/Ovicide and Lice Adulticide

For Dogs 7 weeks and older weighing between 4.6 and 11 kg
Once-A-Month Topical Flea and Lice Treatment

Liquid

READ THE LABEL AND INSERT BEFORE USING

FOR USE ONLY ON DOGS
Do not use on other animals

KEEP OUT OF REACH OF CHILDREN

WARNING



POISON

EYE IRRITANT

DOMESTIC

ACTIVE INGREDIENT: Imidacloprid 9.1% and Pyriproxyfen 0.46%

REG. NO. 27583 P.C.P. ACT

CONTENTS: One to Six 1.0 mL Tubes

Elanco Canada Limited
1919 Minnesota Court, Suite 401
Mississauga, Ontario L5N 0C9
1-800-265-5475

2022-0231

2022-01-31

42

43 Advantage II medium dog provides dual protection against fleas and lice. Advantage II
44 medium dog kills 98-100 % of the existing fleas on dogs within 1 day after treatment.
45 Protection against further flea infestation can last up to 4 weeks. Retreatment at
46 monthly intervals is required if reinfestation is continuous or if flea control is desired.
47 Fleas, eggs, and larvae are killed following contact with skin debris from an Advantage
48 II medium dog treated pet. Advantage II medium dog provides multi-stage flea control
49 effectively breaking all flea lifecycle stages for quick and lasting control of flea
50 populations. Advantage II medium dog treats lice infestations on dogs.

51

52 Do not shampoo or immerse the animal in water immediately after treatment. The
53 animal can be shampooed, immersed in water or exposed to rain 3 days after
54 treatment. Advantage II medium dog retains its ovicidal and larvicidal efficacy up to 4
55 weeks following a shampoo treatment, swimming or exposure to rain. The adulticidal
56 effect may be shortened by repeated shampooing, water immersion or exposure to rain.
57 Reapplication may be required after such activities.

58

59 The successive feeding activity of fleas on pets may elicit a hypersensitivity skin
60 disorder known as flea allergy dermatitis (FAD). Treatment of pets with Advantage II
61 medium dog rapidly kills fleas and may reduce the incidence of this condition. For
62 effective flea and lice control, treatment of the pet should be combined with sanitation of
63 any area used by the pet. Vacuum floors, carpets and furniture (discard vacuum bag
64 after use) and wash the pet's bedding, living quarters and surrounding areas. If pest
65 problems persist, an insecticidal premise treatment may be required. A veterinarian
66 should be consulted when pest infestation continues to be a problem on the pet.

67

68 PRECAUTIONS:

69

70 HUMANS: KEEP OUT OF THE REACH OF CHILDREN. Harmful if swallowed, or
71 absorbed through the skin. Causes eye irritation. Do not get in eyes. Avoid contact with
72 skin or clothing. Wear rubber gloves during application of this product. After handling or
73 applying, wash hands (or any other skin that came in contact with the product), with
74 soap and water. Avoid contact with pets until dry.

75

76 PETS: Do not use on animals under 7 weeks of age. Care should be taken to prevent
77 exposure of other pets who may be in close physical contact with the treated animal.
78 Individual sensitivities, while rare, may occur after using ANY pesticide product for pets.
79 Smaller animals are more likely to experience an adverse reaction. Observe carefully. If
80 signs persist, or become more severe, consult a veterinarian immediately.

81

Side Effects: Monitor your dog after application. Side effects may include: skin irritation such as redness or scratching; changes in behaviour such as agitation or lethargy; or gastrointestinal effects such as vomiting or loss of appetite. If these or other side effects occur, consult your veterinarian or Elanco at 1-800-265-5475.

82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124

FIRST AID:

HUMANS: IF IN EYES, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

Call a poison control centre or doctor for treatment advice.

IF SWALLOWED, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person. **IF ON SKIN OR CLOTHING,** take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

PETS: If signs of sensitivity occur, bathe the pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical or veterinary attention.

TOXICOLOGICAL INFORMATION: Treat the patient symptomatically.

DISPOSAL: Do not reuse empty container. Wrap empty tube and cap in newspaper and dispose in household garbage.

DIRECTIONS FOR USE:

This product contains imidacloprid and pyriproxyfen. Do not apply another pest control product such as a shampoo, collar, or powder that contains imidacloprid or pyriproxyfen to the treated animal after applying Advantage II Medium Dog.

Do NOT administer by mouth. For dogs 7 weeks and older weighing between 4.6 and 11 kg. This product is packaged in single dosage tubes. This package contains one to six monthly 1.0 mL dosages. Treat your dog monthly in the manner described below. Do not apply a dosage higher than one 1.0 mL tube at one time. Consult a veterinarian before using on sick, aged, pregnant or nursing animals or animals receiving drug or other pesticide treatment. A veterinarian should be consulted when pest infestation continues to be a problem on the pet. Retreatment at monthly intervals is required if reinfestation is continuous or if flea control is desired.

Use only on dogs. Do not use on other animals.

Do not use on puppies under 7 weeks of age.

Remove one applicator from the package.

Hold applicator tube in upright position. Pull cap off tube.

Turn cap around and place other end of cap back on tube.

Twist cap to break seal, then remove cap from tube.

2022-0231

2022-01-31

125 The dog should be standing for easy application. The entire contents of the Advantage
126 II medium dog tube should be applied evenly to two or three spots on the top of the
127 back from the shoulder to the base of the tail. At each spot part the hair until the skin is
128 visible. Place the tip of the tube on the skin and gently squeeze to dispense a portion of
129 the solution on the skin. Do not apply an excessive amount of solution at one spot that
130 could cause some of the solution to run off the side of the dog. Do not get this product in
131 your pet's eyes or administer by mouth. Until Advantage II medium dog dries, keep pet
132 off furniture and away from hardwood floors.

133

134 Discard the empty tube as described in "DISPOSAL".

135

136 RETAIN PACKAGING INFORMATION FOR FUTURE REFERENCE

137

138 Advantage, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.

139 ©2022 Elanco or its affiliates.

140

2022-0231
2022-01-31

141
142
143 <Elanco Logo>

Advantage™ II Medium Dog
Carton (English)

Advantage™ II medium dog

Flea Adulticide/Larvicide/Ovicide and Lice Adulticide

For Dogs 7 weeks and older weighing between 4.6 and 11 kg
Once-A-Month Topical Flea and Lice Treatment

Liquid

READ THE LABEL AND INSERT BEFORE USING

FOR USE ONLY ON DOGS
Do not use on other animals

KEEP OUT OF REACH OF CHILDREN

WARNING



POISON

EYE IRRITANT

DOMESTIC

ACTIVE INGREDIENT: Imidacloprid 9.1% and Pyriproxyfen 0.46%

REG. NO. 27583 P.C.P. ACT

CONTENTS: One to Six 1.0 mL Tubes

Elanco Canada Limited
1919 Minnesota Court, Suite 401
Mississauga, Ontario L5N 0C9
1-800-265-5475

2022-0231

2022-01-31

182 Advantage II medium dog provides dual protection against fleas and lice. Advantage II
183 medium dog kills 98-100 % of the existing fleas on dogs within 1 day after treatment.
184 Protection against further flea infestation can last up to 4 weeks. Retreatment at
185 monthly intervals is required if reinfestation is continuous or if flea control is desired.
186 Fleas, eggs, and larvae are killed following contact with skin debris from an Advantage
187 II medium dog treated pet. Advantage II medium dog provides multi-stage flea control
188 effectively breaking all flea lifecycle stages for quick and lasting control of flea
189 populations. Advantage II medium dog treats lice infestations on dogs.

190
191 Do not shampoo or immerse the animal in water immediately after treatment. The
192 animal can be shampooed, immersed in water or exposed to rain 3 days after
193 treatment. Advantage II medium dog retains its ovicidal and larvicidal efficacy up to 4
194 weeks following a shampoo treatment, swimming or exposure to rain. The adulticidal
195 effect may be shortened by repeated shampooing, water immersion or exposure to rain.
196 Reapplication may be required after such activities.

197
198 **PRECAUTIONS:**

199
200 **HUMANS: KEEP OUT OF THE REACH OF CHILDREN.** Harmful if swallowed, or
201 absorbed through the skin. Causes eye irritation. **DO NOT** get in eyes. Avoid contact
202 with skin or clothing. Wear rubber gloves during application of this product. After
203 handling or applying, wash hands (or any other skin that came in contact with the
204 product), with soap and water. Avoid contact with pets until dry.

205
206 **PETS:** Do not use on animals under 7 weeks of age. Care should be taken to prevent
207 exposure of other pets who may be in close physical contact with the treated animal.
208 Individual sensitivities, while rare, may occur after using ANY pesticide product for pets.
209 Smaller animals are more likely to experience an adverse reaction. Observe carefully. If
210 signs persist, or become more severe, consult a veterinarian immediately.

211

Side Effects: Monitor your dog after application. Side effects may include: skin irritation such as redness or scratching; changes in behaviour such as agitation or lethargy; or gastrointestinal effects such as vomiting or loss of appetite. If these or other side effects occur, consult your veterinarian or Elanco at 1-800-265-5475.

212
213 **FIRST AID:**

214
215 **HUMANS: IF IN EYES,** hold eye open and rinse slowly and gently with water for 15-20
216 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue
217 rinsing eye.

218 Call a poison control centre or doctor for treatment advice.

219 **IF SWALLOWED,** call a poison control centre or doctor immediately for treatment
220 advice. Have person sip a glass of water if able to swallow. Do not induce vomiting
221 unless told to do so by a poison control centre or doctor. Do not give anything by mouth

2022-0231

2022-01-31

222 to an unconscious person. IF ON SKIN OR CLOTHING, take off contaminated clothing.
223 Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control
224 centre or doctor for treatment advice.

225

226 PETS: If signs of sensitivity occur, bathe the pet with mild soap and rinse with large
227 amounts of water. If signs continue, consult a veterinarian immediately.

228 Take container, label or product name and Pest Control Product Registration Number
229 with you when seeking medical or veterinary attention.

230

231 TOXICOLOGICAL INFORMATION: Treat the patient symptomatically.

232

233 DISPOSAL: Wrap empty tube and cap in newspaper and dispose in household
234 garbage.

235

2022-0231
2022-01-31

236
237
238
239
240
241
242
243

244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262

<Elanco Logo>

Advantage™ II Medium Dog
Tube (English)

Advantage™ II medium dog

READ THE LABEL AND INSERT BEFORE USING

WARNING



POISON

EYE IRRITANT

DOMESTIC

EXTERNAL USE ONLY

ACTIVE INGREDIENT: Imidacloprid 9.1% and Pyriproxyfen 0.46%

REG. NO. 27583 P.C.P. ACT

Elanco Canada Limited
Mississauga, Ontario L5N 0C9

1.0 mL