

2020-11-17  
Sub.No. 2020-3957

IPG EXTRA  
STABILIZED CHLORINATING TABLETS  
CONTROLS BACTERIA AND ALGAE in Spa Waters  
DOMESTIC

REG. NO. 27565 P.C.P. ACT

ACTIVE INGREDIENT:  
AVAILABLE CHLORINE (present as Trichloro-s-triazinetrione).....90%

CAUTION

POISON



CORROSIVE



xx g per tablet

Net Contents  
{Format}

READ THE LABEL BEFORE USING  
KEEP OUT OF REACH OF CHILDREN

International Pool Group Inc  
7 Villandry Street Nepean  
(Ontario) K2G 6X7  
1-866-470-7264

## DIRECTIONS FOR USE

Weight per tablet: xx g

### WARNING:

Direct tablet contact may cause bleaching of spa surfaces. Automatic feeding devices employing these tablets must be totally free from traces of other spa sanitizers to avoid the probability of an explosion.

1. Maintain recommended daily levels as determined by testing kit.

Free available chlorine: 3.0 – 5.0 ppm

pH: 7.2 – 7.8

Total alkalinity: 125 – 150 ppm

Calcium hardness: 150 – 200 ppm

2. Superchlorinate a freshly filled spa by adding 20 g of IPG EXTRA per 1,000 L of spa water. Maintain water circulation for at least 1 hour before entering the spa (always measure your free available chlorine level before using the spa to make sure it falls within the recommended levels). Repeat superchlorination procedure after each use period.

3. If using an automatic chlorinator, fill the chamber with IPG EXTRA tablets and adjust flow rate, according to manufacturer's recommendations to maintain proper chlorine residual. Adding tablets to the spa skimmer basket is an alternative chlorine feed method.

4. Maintain proper chlorine residual by adding 10 g of IPG EXTRA per 1,000 L of spa water, daily or as needed.

5. Persistent foam formation, turbid, or malodorous water are indicators of inadequate sanitizer levels. To correct such conditions, superchlorinate spa water by adding 30 g of IPG EXTRA per 1,000 L of spa water. Maintain circulation at least 1 hour and wait four hours before using (always measure your free available chlorine level before using the spa to make sure it falls within the recommended levels).

6. Note on Superchlorination: IPG EXTRA contains stabilizer (cyanuric acid) which will accumulate in spa water.

Use of IPG EXTRA for superchlorination will raise stabilizer level at a faster rate. Keep stabilizer level within 30-100 ppm range by dilution with fresh water. Superchlorination with IPG EXTRA will also lower total alkalinity. Make sure that the total alkalinity stays within recommended levels.

7. NOTE: Maximum spa water usage temperature is 40°C. Bathers should not stay more than 15 minutes in spa water at 40°C. For proper sanitation, spa must be completely drained periodically. The number of days between COMPLETE SPA DRAINAGE is equal to the volume of spa water in litres, divided by 10 times the maximum number of daily spa users. Refill spa with water and repeat DIRECTIONS FOR USE.

## PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN. Strong oxidizing agent. May cause fire or explosion if contaminated. Dangerous gas formed when mixed with acid. Do not mix with any other chemical. It is recommended to wear rubber gloves when handling this product. Avoid contact with skin and eyes. Harmful if swallowed. Keep container closed and store in a cool, dry location.

## DISPOSAL

Do not reuse empty container. Wrap and dispose of empty container with household garbage. For information on disposal of unused, unwanted product, contact the manufacturer or the Provincial Regulatory Agency. Contact the manufacturer and the Provincial Regulatory Agency in case of a spill and for clean-up of spills.

## FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice. If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

## TOXICOLOGICAL INFORMATION

Treat symptomatically.