

2010-2186
08-FEB-2011

Canada Placepack text: paste products:

PCP No. 29504 - DOMESTIC

Live Print Area

W=1.75 x H=1.375

Front:

Difethialone Paste Place Packs Rodenticide
Diféthialone paquets de pâte raticide à disposer
REG. NO./NO. D'ENR 29504 PCP ACT/LOI PA
DIFETHIALONE 0.0025%
KEEP OUT OF REACH OF CHILDREN
(*INSERT DIAMOND Skull & Crossbones*)
WARNING, POISON; contains the allergens
wheat and soy.
GARDER HORS DE LA PORTÉE DES
ENFANTS
AVERTISSEMENT, POISON; contient les
allergènes blé et soja.

Back:

DOMESTIC DOMESTIQUE
Liphatech Inc., 3600 W. Elm Street,
Milwaukee, WI 53209 (414) 351-1476
MADE IN FRANCE
CONTENU NET CONTENTS: 10 g
READ THE ENTIRE LABEL ON THE
OUTER CARTON BEFORE USE
LIRE L'ÉTIQUETTE AU COMPLET SUR
L'EXTÉRIEUR DU CARTON AVANT
L'UTILISATION

"Main Carton Label"

Difethialone Paste Place Packs Rodenticide

Kills Norway Rats, Roof Rats, House Mice and Warfarin-Resistant House Mice
Rats and Mice may consume a lethal dose in a single feeding

DOMESTIC

GUARANTEE: Difethialone..... 0.0025% (when packed)

REG. NO. 29504 P.C.P. ACT

KEEP OUT OF REACH OF CHILDREN
READ THE LABEL BEFORE USING
WARNING, CONTAINS THE ALLERGENS WHEAT AND SOY

WARNING
[Diamond Skull & Crossbones Symbol]
POISON

NET CONTENTS: 1000 g / 1 Kg (100 x 10 g)

Liphatech Inc.
3600 W. Elm Street
Milwaukee, WI
53209 U.S.A.
(414) 351-1476

DIRECTIONS FOR USE:

For the control of Norway rats, Roof rats and House mice inside and in areas adjacent to homes.

Use only in sites listed on the product label.

CARCASS DISPOSAL: To protect pets and non-target wildlife, carefully dispose of animals killed by this bait. Wrap poisoned carcasses securely with paper. Dispose of them by: placing in a closed container in the garbage OR burying deep enough to prevent scavengers from reaching them.

When this bait is used in an area accessible to birds, it must be placed in a bait station that prevents birds from eating the bait.

Selection of Treatment Areas: Treated baits should be placed in tamper-resistant bait stations, or in locations not accessible to children, pets, wildlife or domestic animals. Tamper-resistant bait stations must have the following characteristics: be constructed of high-strength material and resistant to destruction by children and animals; have an entrance which prevents children and non-target animals from reaching bait; have an internal structure which prevents bait from being shaken loose; have an access panel which fastens securely and locks; capable of being securely fastened to a surface; be clearly labelled: **WARNING POISON**.

Place bait in locations where rats and mice travel and will most likely find and consume the bait. Placement along walls, in corners, by gnawed openings, in and around burrows, and in concealed locations where rats and mice feed is desirable. Maintain a steady supply of fresh bait. Remove any spoiled or contaminated bait promptly. Do not place bait in areas where there is a possibility of contaminating food or surfaces that may come in contact with food.

APPLICATION DIRECTIONS:

For Norway Rats (*Rattus norvegicus*) and Roof Rats (*Rattus rattus*): Norway rats usually inhabit the ground floor of structures and burrow in the ground. They will range approximately 15 to 100 m from their nesting area seeking food and water. Roof rats, however, are excellent climbers and usually infest the upper stories of structures. Hence, special attention must be paid to the ability of this species to climb by placing baits in the roof rats activity area. For both species, eliminate as much of the naturally occurring food as possible. Place 10 to 20 packs of bait per placement at intervals of 5 to 10 meters. Larger amounts (up to 50 packs) may be necessary at points of very high rat activity. Maintain an uninterrupted supply of fresh bait for 10 days, or until signs of rat activity cease.

For House Mice (*Mus musculus*): House mice are excellent climbers and will infest all floors in a structure. Mice will only range approximately 10 m from their nesting area seeking food. Eliminate as much of the naturally occurring food as possible. Place 1 pack of bait per placement at intervals of 2 to 4 meters. If a bait placement is not disturbed after several days, move the placement approximately 1 meter. Two to five packs of bait may be necessary at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for 15 days, or until signs of mouse activity cease.

Note: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. To prevent reinfestation, eliminate rodent food, water and harbourage as much as possible. If reinfestation occurs, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN, PETS, DOMESTIC ANIMALS AND WILDLIFE. Keep away from feed or food. May irritate eyes. May be harmful or fatal if swallowed or absorbed through the skin because this material may reduce the clotting ability of blood and cause bleeding. Do not open pre-measured place packs. Chemical-resistant gloves must be worn when disposing of dead rodents, unconsumed bait and empty containers. Do not get in eyes, on skin, or on clothing. Wash arms, hands, and face with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds.

FIRST AID:

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person. If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. If on skin or clothing: Take off contaminated clothing. Rinse skin with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

Vitamin K1 in the form of intramuscular or subcutaneous injections, or by oral ingestion are suggested remedial treatments for anticoagulant poisoning. The severity of the case measured by establishing prolonged prothrombin times (P.T.) will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments. Contains wheat and soy - may cause allergic reaction in sensitive persons.

STORAGE AND DISPOSAL:

Store bait in original, tightly sealed container. Store in a cool, dry place. Do not contaminate water, food or feed by storage or disposal. Do not reuse empty container. Wrap and dispose of empty container with household garbage. Unused or partially used products should be disposed at provincially or municipally designated hazardous waste disposal sites. For information on disposal of unused or unwanted product, contact the manufacturer or the provincial regulatory agency.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

OPTIONAL LABEL STATEMENTS

Bitrex™ is a trademark of Macfarlan Smith Ltd., Edinburgh, Scotland

CONTAINS BITTERING AGENT

RATS AND MICE CAN CONSUME A LETHAL DOSE IN ONE NIGHT'S FEEDING WITH FIRST DEAD RODENTS APPEARING FOUR OR FIVE DAYS AFTER FEEDING BEGINS

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.