[Front Panel English Copy]

Hagen Flea Collar for Puppies and Small Dogs with Integral Buckle

* Kills fleas up to 4 months and American Dog and Brown Dog ticks up to 2 months

DOMESTIC

READ THE LABEL AND ENCLOSED LEAFLET BEFORE USING. FOR USE ONLY ON DOGS.
KEEP OUT OF REACH OF CHILDREN.

GUARANTEE: Propoxur 9.48%

REG. NO 27609 P.C.P. ACT

Warning, contains the allergen Soy.

Distributed by: Rolf C. Hagen Inc.

3225 Sartelon Montreal, QC H4R 1E8

NET CONTENTS 12.5g (38.5cm)

[Side Panel English Copy]

Hagen Flea Collar For Puppies and Small Dogs with Integral Buckle

[Insert Leaflet Copy]

Hagen Flea Collar For Puppies and Small Dogs with Integral Buckle

DOMESTIC

READ THE LABEL AND LEAFLET BEFORE USING. FOR USE ONLY ON DOGS. KEEP OUT OF REACH OF CHILDREN.

DIRECTIONS FOR USE:

One collar/animal.

For best results, before placing this collar on, bathe and shampoo pet with a pyrethrin based product such as Hagen Flea & Tick Shampoo for Dogs. This will produce a "quick kill" effect.

Do not open protective pouch until ready to use. Remove the collar from the package and place around dog's neck. Thread flat end of the collar through buckle hole and tighten as necessary, so that at least two ridges are covered by the buckle, leaving ample room for expansion as the dog grows. Cut off any excess length and discard in garbage. Collar must be worn loosely, but securely enough to prevent easy removal. After collar placement, adult ticks will be killed in a few days and will fall off or may be easily removed. American Dog and Brown Dog ticks appear on dogs in three stages. In the first two stages, ticks are small and are difficult to see. The collar will also control these immature stages. Collar will kill ticks encountered in first 2 months of wear. Kills dog fleas for up to 4 months. For control of ticks, replace this collar no more than once every 2 months. For the control of fleas, replace this collar no more than once every 4 months. Replace every 2 months for continual pest control in situations where there is an ongoing risk of tick infestation. Replace every 4 months for continual pest control in situations where there is an ongoing risk of flea infestation. To release the collar, squeeze between thumb and finger and pull sharply. Do not use on nursing dogs, or dogs under 12 weeks of age. When collar is first worn, observe neck area every few days for irritation or adverse reaction. Consult a veterinarian before using on sick, aged, pregnant or nursing animals or animals receiving drug or other pesticide treatment. RETAIN PACKAGING INFORMATION FOR FUTURE REFERENCE.

Temporary wetting will not reduce effectiveness. If your dog goes swimming or is out in the rain, it is not necessary to remove the collar. The continuous action collar will rapidly replace any flea and tick killing material removed. For effective flea and tick control, treatment of the pet should be combined with the sanitation of any area used by the pet. Vacuum floors, carpets, and furniture (discard vacuum bag after use) and wash pet's bedding, living quarters, and surrounding areas. If pest problems persist, an insecticidal premise treatment may be required. For best results, use in combination with a complete flea eradication program (consult your local pet retailer for more information). A veterinarian should be consulted when pest infestation continues to be a problem on the pets.

PRECAUTIONS: HUMANS: KEEP OUT OF REACH OF CHILDREN. Do not allow children to handle this collar. Dust will form on the collar during storage. Do not get dust or collar in the mouth. Harmful if swallowed. Do not get dust in eyes. Wash hands (or any other skin that came into contact with the product) thoroughly with soap and water after handling. Collar is intended for use only as an insecticide generator and is not to be taken internally by humans or animals. PETS: Do NOT use this product on dogs at

the same time or within 30 days before or after treatment with, or exposure to, other cholinesterase-inhibiting pesticides, drugs or chemicals.

FIRST AID: HUMANS: If dust gets in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. If dust gets on skin or clothing, take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice. If ingested, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention. PETS: Sensitivities may occur after using ANY pesticide for pets. If signs of sensitivity occur, remove collar and bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately. If the collar is swallowed or chewed, consult a veterinarian.

TOXICOLOGICAL INFORMATION: This product contains a pesticide that is a cholinesterase inhibitor. Atropine is antidotal.

DISPOSAL: Wrap and dispose of any excess collar, packaging and expired collars with garbage to keep out of reach of children.

GUARANTEE: Propoxur...9.48% REG. NO. 27609 P.C.P. ACT Warning, contains the allergen Soy.

NET CONTENTS: 12.5 g (38.5 cm)

Distributed by: Rolf C. Hagen Inc. 3225 Sartelon Montreal, QC H4R 1E8

Made in U.S.A.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.