

2021-3866  
2024-08-29

<b>GROUP</b>	<b>29</b>	<b>INSECTICIDE</b>
--------------	-----------	--------------------

**Label**

## **Beleaf® 50SG Insecticide**

**COMMERCIAL**

**SOLUBLE GRANULES**

**READ THE LABEL AND THE PAMPHLET BEFORE USING**

**ACTIVE INGREDIENT: Flonicamid..... 50.0%**

**Warning, contains the allergens milk and sulfites**

**Contains 500 grams of Flonicamid per kilogram of formulated product**

**CAUTION – EYE IRRITANT**

**REGISTRATION NO: 29796    PEST CONTROL PRODUCTS ACT**

**Net Contents: 0.5 – 100 kg**

**ISK Biosciences Corporation  
7470 Auburn Road  
Suite A  
Concord, Ohio 44077 U.S.A.  
1-877-706-4640**

## **PRECAUTIONS**

### **KEEP OUT OF REACH OF CHILDREN**

**DO NOT** take internally. Harmful if swallowed.

May irritate eyes.

Avoid contact with eyes.

Avoid prolonged contact with skin. Wash exposed areas of skin thoroughly with soap and warm water after handling or using. Remove contaminated clothing and wash before re-use.

Avoid breathing spray mist.

Wear coveralls over a long-sleeved shirt and long pants, chemical-resistant gloves, socks and shoes, and protective eyewear (goggles or face shield) during mixing, loading, application, clean-up and repair. In addition, wear chemical-resistant headgear during open cab airblast/air assisted application. Chemical-resistant headgear includes Sou'Wester hat, chemical-resistant rain hat or large brimmed waterproof hat and hood with sufficient neck protection. Gloves and protective eyewear are not required during application within a closed cab.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. For pome fruit, stone fruit, hops, outdoor ornamentals (excluding conifers), and sweet corn please refer to the individual use descriptions for activity-specific restricted entry intervals.

Apply only to agricultural crops when the potential for drift to areas of human habitation and human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

**DO NOT** apply using handheld mistblower/airblast or handheld fogging equipment in greenhouses.

**DO NOT** apply using handheld mist blower/airblast or handheld fogging equipment.

**DO NOT** apply by air.

### **FIRST AID**

**IF SWALLOWED:** Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

In Case of Emergency: Call 1-888-484-7546.

## **TOXICOLOGICAL INFORMATION**

Treat symptomatically.

## **ENVIRONMENTAL PRECAUTIONS**

Toxic to certain beneficial arthropods (which may include predatory and parasitic insects, spiders and mites). Minimize spray drift to reduce harmful effects on beneficial arthropods in habitats next to the application site such as hedgerows and woodland.

TOXIC to non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

Greenhouse use: Toxic to certain beneficial arthropods (which may include predatory and parasitic insects, spiders, and mites). May harm certain beneficial arthropods, including those used in greenhouse production. Avoid application when beneficial arthropods are in the treatment area.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

## **STORAGE**

Store this product away from food or feed. Store product in original container, in a secured, dry place separate from other pesticides and fertilizer.

## **DISPOSAL**

Recyclable containers:

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

Recyclable containers:

DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

Refillable containers:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

Disposal of unused, unwanted product:

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

**NOTICE TO USER**

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

Beleaf® is a registered trademark of Ishihara Sangyo Kaisha, Ltd.

**Pamphlet**

## **Beleaf® 50SG Insecticide**

**COMMERCIAL**

SOLUBLE GRANULES

READ THE LABEL AND THE PAMPHLET BEFORE USING

ACTIVE INGREDIENT:..... Flonicamid 50.0%

Warning, contains the allergens milk and sulfites

Contains 500 grams of Flonicamid per kilogram of formulated product

CAUTION – EYE IRRITANT

REGISTRATION NO: 29796    PEST CONTROL PRODUCTS ACT

ISK Biosciences Corporation  
7470 Auburn Road  
Suite A  
Concord, Ohio 44077 U.S.A.  
1-877-706-4640

## **PRECAUTIONS**

### **KEEP OUT OF REACH OF CHILDREN**

**DO NOT** take internally. Harmful if swallowed.

May irritate eyes.

Avoid contact with eyes.

Avoid prolonged contact with skin. Wash exposed areas of skin thoroughly with soap and warm water after handling or using. Remove contaminated clothing and wash before re-use.

Avoid breathing spray mist.

Wear coveralls over a long-sleeved shirt and long pants, chemical-resistant gloves, socks and shoes, and protective eyewear (goggles or face shield) during mixing, loading, application, clean-up and repair. In addition, wear chemical-resistant headgear during open cab airblast/air assisted application. Chemical-resistant headgear includes Sou'Wester hat, chemical-resistant rain hat or large brimmed waterproof hat and hood with sufficient neck protection. Gloves and protective eyewear are not required during application within a closed cab.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. For pome fruit, stone fruit, hops, outdoor ornamentals (excluding conifers), and sweet corn please refer to the individual use descriptions for activity-specific restricted entry intervals.

Apply only to agricultural crops when the potential for drift to areas of human habitation and human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

**DO NOT** apply using handheld mistblower/airblast or handheld fogging equipment in greenhouses.

**DO NOT** apply using handheld mist blower/airblast or handheld fogging equipment.

**DO NOT** apply by air.

### **FIRST AID**

**IF SWALLOWED:** Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

In Case of Emergency: Call 1-888-484-7546.

## **TOXICOLOGICAL INFORMATION**

Treat symptomatically.

## **ENVIRONMENTAL PRECAUTIONS**

Toxic to certain beneficial arthropods (which may include predatory and parasitic insects, spiders, and mites). Minimize spray drift to reduce harmful effects on beneficial arthropods in habitats next to the application site such as hedgerows and woodland.

**TOXIC** to non-target terrestrial plants. Observe spray buffer zones specified under **DIRECTIONS FOR USE**.

**Greenhouse use:** Toxic to certain beneficial arthropods (which may include predatory and parasitic insects, spiders, and mites). May harm certain beneficial arthropods, including those used in greenhouse production. Avoid application when beneficial arthropods are in the treatment area.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.  
Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

## **STORAGE**

Store this product away from food or feed. Store product in original container, in a secured, dry place separate from other pesticides and fertilizer.

## **DISPOSAL**

### Recyclable containers:

**DO NOT** reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

Recyclable containers:

DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

Refillable containers:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

Disposal of unused, unwanted product:

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills



## **DIRECTIONS FOR USE**

Beleaf<sup>®</sup> 50SG Insecticide can be used against the listed pests on flax, brassica (cole) leafy vegetables, cucurbit vegetables, fruiting vegetables, leafy vegetables (excluding Brassica), hop, tuberous and corm vegetables, root vegetables (excluding sugar beets), legume vegetables (succulent or dried, excluding soybeans, Crop Group 6), pome fruit, stone fruit, alfalfa grown for seed, herbs (Crop Group 25), greenhouse cucumbers, greenhouse strawberry, greenhouse tomatoes, greenhouse peppers, greenhouse lettuce, greenhouse Chinese cabbage, bok choy, Bushberries (Crop Subgroup 13-07B), low growing berries (Crop Subgroup 13-07G), non-grass Animal Feeds (Crop Group 18), greenhouse ornamentals (including cut flowers), outdoor ornamentals (excluding conifers) and sweet corn.

The rate of application is dependent upon the insect species present, the level of insect pressure, and the amount of foliage present. Begin applications before populations begin to build or at economic thresholds according to local economic guidelines. Refer to local Cooperative Extension Guideline and/or time applications for scouting results. Thorough plant coverage is essential for good performance.

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

DO NOT allow releases, effluent or runoff from greenhouses containing this product to enter lakes, streams, pond or other waters.

### **Mode of Action**

Beleaf<sup>®</sup> 50SG Insecticide is a member of the pyridinecarboxamide class of chemistry. Beleaf<sup>®</sup>50SG Insecticide controls target pests by contact and ingestion provoking rapid and irreversible feeding cessation. Aphids could remain on the plant until they desiccate.

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

DO NOT allow effluent or runoff from greenhouses containing this product to enter lakes, streams, pond or other waters.

**Field sprayer application:** DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Boom height must be 60 cm or less above the crop or ground.

**Airblast application:** DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT direct spray above plants to be treated. Turn off outward pointing nozzles at row ends and outer rows. DO NOT apply when wind speed is greater than 16 km/h at the application site as measured outside of the treatment area on the upwind side.

**Chemigation:** DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Applications MUST be conducted WITHOUT the use of end guns.

DO NOT apply by air.

**Spray Buffer zones:**

A spray buffer zone is NOT required for:

- uses with hand-held application equipment permitted on this label,
- soil drench.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands).

Method of application	Crop		Spray Buffer Zones (metres) Required for the Protection of:
			Terrestrial habitat
Field sprayer	Flax, Cucurbit vegetables, Fruiting vegetables, Hop, Tuberos and corn vegetables, Root vegetables, Alfalfa grown for seed, Herbs (Crop Group 25), Outdoor ornamentals, Bushberries (Crop Subgroup 13-07B), Low growing berry subgroup 13-07G, Legume vegetables – Succulent or Dried (except soybean) (Crop Group 6), Non-grass animal feeds Crop Group 18, Crop subgroup 4- 13A: Leafy vegetables – leafy greens, Crop subgroup 4- 13B: Leafy vegetables – Brassica leafy greens, Crop group 5-13: Brassica head and stem vegetables, Celtuce, Florence fennel, Kohlrabi, Crop subgroup 22B: Stalk, stem and leaf petioles – leaf petiole vegetables, sweet corn		1
Airblast	Pome fruit, stone fruit, outdoor ornamentals, and Balsam Fir and Fraser Fir, including Christmas trees	Early growth stage	1
		Late growth stage	1

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pesticides portion of the Canada.ca website.

### **Resistance Management**

For resistance management, please note that Beleaf® 50SG Insecticide contains a Group 29 insecticide. Any insect population may contain individuals naturally resistant to Beleaf® 50SG Insecticide and other Group 29 insecticides. The resistant individuals may dominate the insect population if this group of insecticides is used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but are specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay insecticide resistance:

- Where possible, rotate the use of Beleaf® 50SG Insecticide or other Group 29 insecticides with different groups that control the same pests. Avoid application of more than 3 consecutive sprays of Beleaf® 50SG Insecticide or other insecticides in the same group in a year.
- Use tank mixtures with insecticides from a different group when such use is permitted.
- Insecticide use should be based on an IPM program that includes scouting and record keeping, and considers cultural, biological and other chemical control practices.
- Monitor treated pest populations for resistance development.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact ISK Biosciences Corporation at 1-440-357-4640 (Long distance charges apply).

### **Use Restrictions**

DO NOT use in residential areas. Residential areas are defined as any use site where bystanders, including children, could be exposed during or after application. This includes areas in and around homes, schools, public buildings or any other areas where they could be exposed.

#### **Crop Rotation Restrictions**

Following application of Beleaf® 50SG Insecticide, any crop listed on this label may be planted at any time.

All other crops may be planted 30 days after the last application of Beleaf® 50SG Insecticide.

### **Pre-Harvest Intervals**

<b>Crop</b>	<b>PHI Days</b>
Brassica Head and Stem Vegetables	0
Cucurbit Vegetables	0
Flax	7
Fruiting Vegetables (except cucurbits)	0
Hop	31
Leafy Greens	0
<i>Brassica</i> Leafy Greens	0

Leaf Petioles Vegetables	0
Tuberous and Corm Vegetables	7
Root Vegetables (except sugar beets)	3
Pome Fruit	21
Stone Fruit	14
Cucumber, greenhouse	0
Tomato, greenhouse	0
Pepper, greenhouse	0
Lettuce, greenhouse	0
Herbs (Crop Group 25)	7
Low growing berry subgroup 13-07G	0
Bushberries (Crop Subgroup 13-07B)	0
Kohlrabi	0
Celtuce	0
Florence fennel	0
Greenhouse Strawberry	0
Crop Group 18 Non-grass Animal Feeds	7
Legume Vegetables -- Succulent or Dried (except soybean) (Crop Group 6)	7
Sweet corn	0
Greenhouse Chinese cabbage, bok choy	0

### **Mixing and Loading Instructions**

Ensure the spray system is clean and free of residues from previous applications. Fill the spray tank 1/2 full with clean water. Ensure the agitation system is operating and sufficient to provide uniform spray mixing during application and until the spray tank has been emptied. Complete filling the spray tank to the desired level.

Avoid the overnight storage of Beleaf<sup>®</sup> 50SG Insecticide spray mixtures.

Premixing Beleaf<sup>®</sup> 50SG Insecticide solutions in nurse tanks is not recommended. Prepare no more spray mixture than is needed for the immediate application.

Do not use liquid fertilizer as a carrier for Beleaf<sup>®</sup> 50SG Insecticide.

### **Adjuvant Use Requirements and Precautions**

Beleaf<sup>®</sup> 50SG Insecticide should not be used with spray adjuvants.

### **Tank Mixing and Compatibility**

This product may be tank mixed with a supplement or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

Conduct a jar test with intended tank-mix product(s) prior to preparation of large volumes to ensure compatibility. Use the following procedure: 1) Pour the recommended proportions of the

products into a suitable container of water, 2) Mix thoroughly and 3) Allow to stand 5 minutes. If the combination remains mixed or can be re-mixed readily, it is considered physically compatible. Any physical incompatibility in the jar test indicates that Beleaf<sup>®</sup>50SG Insecticide must not be used in the tank-mix. Since not all crop varieties have been tested with all possible tank mix combinations, the combination should also be tested for crop safety on a small portion of the crop to ensure that phytotoxicity will not occur.

### **Spray Equipment Clean Out**

After spraying Beleaf<sup>®</sup>50SG Insecticide and before using sprayer equipment for any other applications, the sprayer must be thoroughly cleaned.

### **Application Information**

All ground application equipment must be properly maintained and calibrated using appropriate carriers. Use the largest droplet size consistent with good pest control.

Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage. Finished spray volumes should be increased under extreme pest populations or dense plant foliage.

### **Ground Application**

Utilize a boom and nozzle sprayer equipped with the appropriate nozzles, spray tips and screens and adjusted to provide optimum spray distribution and coverage at the appropriate operating pressures. Utilize nozzles that produce minimal amounts of fine spray droplets. Do not exceed 205 kPa (30 psi) spray pressure unless otherwise required by the manufacturer of drift reducing nozzles. Sprayers should be adjusted to position spray tips a minimum of 46 cm (18 inches) above the crop. Be aware that overlaps and slower ground speeds while starting, stopping or turning while spraying may result in higher application rates.

### **Chemigation Application**

Apply this product only through drip (trickle) (greenhouse cucumbers, greenhouse peppers, and greenhouse sweet potato slips only) irrigation systems. Do not apply this product through any other type of irrigation system. Do not connect any irrigation system, including greenhouse systems, used for pesticide application to a public water system. Crop injury, lack of effectiveness or illegal residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact Provincial Extension specialists, equipment manufacturers, or other experts. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must also contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must

include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed causes non-uniform distribution and/or favors drift beyond the area intended for treatment. Do not apply by chemigation if the area to be treated is within 100 metres of a residential area or park.

Beleaf<sup>®</sup> 50SG should be applied continuously for the duration of the water application. Beleaf<sup>®</sup> 50SG Insecticide should be diluted in sufficient volume to insure accurate application over the area to be treated. Use the appropriate amount of water to carry the product to the target pest. Agitation generally is not required when suitable diluents are used. A diluents test should be conducted to ensure that phase separation would not occur during dilution and application. Failure to achieve a uniform dilution throughout the time of application may result in undesirable residues or less than desirable control.

Using Water from Public Water Systems: **DO NOT APPLY Beleaf<sup>®</sup> 50SG INSECTICIDE THROUGH ANY IRRIGATION SYSTEM PHYSICALLY CONNECTED TO A PUBLIC WATER SYSTEM.** Public water system means a system for the provision to the public of piped water for human consumption, if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year. Beleaf<sup>®</sup> 50SG Insecticide may be applied through irrigation systems, which may be **supplied** by a public water system **only if** water from the water system is discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. Before beginning chemigation, always make sure that the air gap exists and that there is no blockage of the overflow of the reservoir tank.

**Brassica Head and Stem Vegetables (0 day PHI)**

**Crop Group 5-13**

**Broccoli; Brussels sprouts; Cabbage; Cabbage, Chinese (napa); Cauliflower, as well as cultivars, varieties, and hybrids of these commodities.**

PESTS	RATE OF APPLICATION		COMMENTS
	Kilograms Beleaf <sup>®</sup> 50SG Insecticide per Hectare	Kilograms Active Ingredient per Hectare	
Aphids	0.12 to 0.16	0.06 to 0.08	Use higher rates for greater pest populations and/or dense foliage.
<p>Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use a minimum of 94 litres per hectare when applied by ground. Finished spray volumes should be increased when plant foliage is dense.</p> <p>Apply before populations reach economic thresholds or as populations begin to increase but before damaging populations become established. Scout fields and reapply if necessary.</p> <p>Allow a minimum of 7 days between applications.</p> <p>Do not apply more than 3 applications per year.</p> <p>Do not apply more than 0.16 kilograms per hectare Beleaf<sup>®</sup> 50SG Insecticide (0.08 kilograms a.i. per hectare) per application. Do not apply more than 0.48 kilograms per hectare Beleaf<sup>®</sup> 50SG Insecticide (0.24 kilograms a.i. per hectare) per year.</p>			

## Cucurbit Vegetables (0 day PHI)

### Crop Group 9

**Chayote (fruit); Chinese waxgourd (Chinese preserving melon); Citron melon; Cucumber; Gherkin; Gourd, edible (includes hyotan, cucuzza, hechima, Chinese okra); Momordica spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber); Muskmelon (hybrids and/or cultivars of Cucumis melo) (includes true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon); Pumpkin; Squash, summer (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini); Squash, winter (includes butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash); Watermelon (includes hybrids and/or varieties of Citrullus spp.).**

PESTS	RATE OF APPLICATION		COMMENTS
	Kilograms Beleaf <sup>®</sup> 50SG Insecticide per Hectare	Kilograms Active Ingredient per Hectare	
Aphids	0.12 to 0.16	0.06 to 0.08	Use higher rates for greater pest populations and/or dense foliage.
Suppression of Tarnished Plant Bug	0.20 (3 applications) or 0.30 (2 applications)	0.10 (3 applications) or 0.15 (2 applications)	Apply when tarnished plant bugs first appear in the field and before populations reach high levels. Beleaf will stop tarnished plant bug feeding rapidly but it may take several days to see a reduction in tarnished plant bug numbers. Reapply when new insects are detected.
<p>Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use a minimum of 94 litres per hectare when applied by ground. Finished spray volumes should be increased when plant foliage is dense.</p> <p>Apply before populations reach economic thresholds or as populations begin to increase but before damaging populations become established. Scout fields and reapply if necessary.</p> <p>Allow a minimum of 7 days between applications.</p> <p>Do not apply more than 3 applications per year.</p> <p>Do not apply more than 0.30 kilograms per hectare Beleaf<sup>™</sup> 50SG Insecticide (0.15 kilograms a.i. per hectare) per application. Do not apply more than 0.60 kilograms per hectare Beleaf<sup>™</sup> 50SG Insecticide (0.30 kilograms a.i. per hectare) per year.</p>			



**Fruiting Vegetables (except cucurbits) (0 day PHI)**

**Crop Group 8-09**

**African eggplant; bell pepper; currant tomato; eggplant; garden huckleberry; goji berry; groundcherry; martynia; okra; pea eggplant; pepino; nonbell pepper; scarlet eggplant; sunberry; tomatillo; tomato; cultivars, varieties, and/or hybrids of these.**

PESTS	RATE OF APPLICATION		COMMENTS
	Kilograms Beleaf <sup>®</sup> 50SG Insecticide per Hectare	Kilograms Active Ingredient per Hectare	
Aphids	0.12 to 0.16	0.06 to 0.08	Use higher rates for greater pest populations and/or dense foliage.
Suppression of Tarnished Plant Bug	0.20 (3 applications) or 0.30 (2 applications)	0.10 (3 applications) or 0.15 (2 applications)	Apply when tarnished plant bugs first appear in the field and before populations reach high levels. Beleaf will stop tarnished plant bug feeding rapidly but it may take several days to see a reduction in tarnished plant bug numbers. Reapply when new insects are detected.
<p>Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use a minimum of 94 litres per hectare when applied by ground. Finished spray volumes should be increased when plant foliage is dense.</p> <p>Apply before populations reach economic thresholds or as populations begin to increase but before damaging populations become established. Scout fields and reapply if necessary.</p> <p>Allow a minimum of 7 days between applications.</p> <p>Do not apply more than 3 applications per year.</p> <p>Do not apply more than 0.30 kilograms per hectare Beleaf<sup>™</sup> 50SG Insecticide (0.15 kilograms a.i. per hectare) per application. Do not apply more than 0.60 kilograms per hectare Beleaf<sup>™</sup> 50SG Insecticide (0.30 kilograms a.i. per hectare) per year.</p>			

**Hop** (31 day PHI):

PESTS	RATE OF APPLICATION		COMMENTS
	Kilograms Beleaf <sup>®</sup> 50SG Insecticide per Hectare	Kilograms Active Ingredient per Hectare	
Aphids	0.12 to 0.16	0.06 to 0.08	Use higher rates for greater pest populations and/or dense foliage.

Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use a minimum of 94 litres per hectare when applied by ground. Finished spray volumes should be increased when plant foliage is dense.

Apply before populations reach economic thresholds or as populations begin to increase but before damaging populations become established. Scout fields and reapply if necessary.

Allow a minimum of 7 days between applications.

Do not apply more than 3 applications per year.

Do not apply more than 0.16 kilograms per hectare Beleaf<sup>®</sup> 50SG Insecticide (0.08 kilograms a.i. per hectare) per application. Do not apply more than 0.48 kilograms per hectare Beleaf<sup>®</sup> 50SG Insecticide (0.24 kilograms a.i. per hectare) per year.

For hops, do not enter or allow worker entry for the purpose of performing stripping and training activities in treated areas during the restricted entry interval (REI) of 31 days.

## Leafy Greens (0 day PHI)

### Crop Subgroup 4-13A

**Amaranth, Chinese; Amaranth, leafy; Aster, Indian; Blackjack; Cat's Whiskers; Cham-chwi; Cham-namul; Chervil, fresh leaves; Chipilin; Chrysanthemum, garland; Cilantro, fresh leaves; Corn salad; Cosmos; Dandelion; Dang-gwi; Dillweed, fresh leaves; Dock; Dol-nam-mul; Ebolo; Endive; Escarole; Famefower; Feather cockscomb; Good King Henry; Huazontle; Jute leaves; Lettuce, bitter; Lettuce (head and leaf); Orach; Parsley, fresh leaves; Plantain buckhorn; Primrose, English; Purslane, garden; Purslane, winter; Radicchio (Red Chicory); Spinach; Spinach, Malabar; Spinach, New Zealand; Spinach, tree; Swiss Chard; Tanier Spinach; Violet, Chinese**

PESTS	RATE OF APPLICATION		COMMENTS
	Kilograms Beleaf <sup>®</sup> 50SG Insecticide per Hectare	Kilograms Active Ingredient per Hectare	
Aphids	0.12 to 0.16	0.06 to 0.08	Use higher rates for greater pest populations and/or dense foliage.
Suppression of Tarnished Plant Bug	0.20 (3 applications) or 0.30 (2 applications)	0.10 (3 applications) or 0.15 (2 applications)	Apply when tarnished plant bugs first appear in the field and before populations reach high levels. Beleaf will stop tarnished plant bugs feeding rapidly but it may take several days to see a reduction in tarnished plant bug numbers. Reapply when new insects are detected.
<p>Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use a minimum of 94 litres per hectare when applied by ground. Finished spray volumes should be increased when plant foliage is dense.</p> <p>Apply before populations reach economic thresholds or as populations begin to increase but before damaging populations become established. Scout fields and reapply if necessary.</p> <p>Allow a minimum of 7 days between applications.</p> <p>Do not apply more than 3 applications per year.</p> <p>Do not apply more than 0.30 kilograms per hectare Beleaf<sup>®</sup> 50SG Insecticide (0.15 kilograms a.i. per hectare) per application. Do not apply more than 0.60 kilograms per hectare Beleaf<sup>®</sup> 50SG Insecticide (0.30 kilograms a.i. per hectare) per year.</p>			

## Brassica Leafy Greens (0 day PHI)

### Crop Group 4-13B

**Arugula; Broccoli raab; Broccoli, Chinese; Cabbage, Abyssinian; Cabbage, seakale; Chinese cabbage, bok choy; Collards; Cress, garden; Cress, upland; Hanover salad; Kale; Maca; Mizuna; Mustard greens; Radish, leaves; Rape greens; Rocket, wild; Shepherd's purse; Turnip greens; Watercress as well as cultivars, varieties and hybrids of these commodities.**

PESTS	RATE OF APPLICATION		COMMENTS
	Kilograms Beleaf <sup>®</sup> 50SG Insecticide per Hectare	Kilograms Active Ingredient per Hectare	
Aphids	0.12 to 0.16	0.06 to 0.08	Use higher rates for greater pest populations and/or dense foliage.
<p>Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use a minimum of 94 litres per hectare when applied by ground. Finished spray volumes should be increased when plant foliage is dense.</p> <p>Apply before populations reach economic thresholds or as populations begin to increase but before damaging populations become established. Scout fields and reapply if necessary.</p> <p>Allow a minimum of 7 days between applications.</p> <p>Do not apply more than 3 applications per year.</p> <p>Do not apply more than 0.16 kilograms per hectare Beleaf<sup>®</sup> 50SG Insecticide (0.08 kilograms a.i. per hectare) per application. Do not apply more than 0.48 kilograms per hectare Beleaf<sup>®</sup> 50SG Insecticide (0.24 kilograms a.i. per hectare) per year.</p>			

**Stalk and Stem Vegetables (0 day PHI)**

**Celtuce, Florence Fennel and Kohlrabi only**

PESTS	RATE OF APPLICATION		COMMENTS
	Kilograms Beleaf <sup>®</sup> 50SG Insecticide per Hectare	Kilograms Active Ingredient per Hectare	
Aphids	0.12 to 0.16	0.06 to 0.08	Use higher rates for greater pest populations and/or dense foliage.
<p>Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use a minimum of 94 litres per hectare when applied by ground. Finished spray volumes should be increased when plant foliage is dense.</p> <p>Apply before populations reach economic thresholds or as populations begin to increase but before damaging populations become established. Scout fields and reapply if necessary.</p> <p>Allow a minimum of 7 days between applications.</p> <p>Do not apply more than 3 applications per year.</p> <p>Do not apply more than 0.16 kilograms per hectare Beleaf<sup>®</sup> 50SG Insecticide (0.08 kilograms a.i. per hectare) per application. Do not apply more than 0.48 kilograms per hectare Beleaf<sup>®</sup> 50SG Insecticide (0.24 kilograms a.i. per hectare) per year.</p>			

## Leaf Petioles Vegetables (0 day PHI)

### Crop Subgroup 22B

**Cardoon; Celery; Celery, Chinese; Fuki; Rhubarb; Udo; Zuiki as well as Cultivars, Varieties and hybrids of these commodities.**

PESTS	RATE OF APPLICATION		COMMENTS
	Kilograms Beleaf® 50SG Insecticide per Hectare	Kilograms Active Ingredient per Hectare	
Aphids	0.12 to 0.16	0.06 to 0.08	Use higher rates for greater pest populations and/or dense foliage.
Suppression of Tarnished Plant Bug	0.20 (3 applications) or 0.30 (2 applications)	0.10 (3 applications) or 0.15 (2 applications)	Apply when tarnished plant bugs first appear in the field and before populations reach high levels. Beleaf will stop tarnished plant bugs feeding rapidly but it may take several days to see a reduction in tarnished plant bug numbers. Reapply when new insects are detected.
<p>Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use a minimum of 94 litres per hectare when applied by ground. Finished spray volumes should be increased when plant foliage is dense.</p> <p>Apply before populations reach economic thresholds or as populations begin to increase but before damaging populations become established. Scout fields and reapply if necessary.</p> <p>Allow a minimum of 7 days between applications.</p> <p>Do not apply more than 3 applications per year.</p> <p>Do not apply more than 0.30 kilograms per hectare Beleaf® 50SG Insecticide (0.15 kilograms a.i. per hectare) per application. Do not apply more than 0.60 kilograms per hectare Beleaf® 50SG Insecticide (0.30 kilograms a.i. per hectare) per year.</p>			

## Tuberous and Corm Vegetables (7 day PHI)

### Crop Subgroup 1C

**Artichoke, Chinese; Artichoke, Jerusalem; Casava, bitter and sweet; Chayote (root); Chufa; Dasheen (taro); Ginger; Potato; Sweet potato; Yam (true).**

PESTS	RATE OF APPLICATION		COMMENTS
	Kilograms Beleaf <sup>®</sup> 50SG Insecticide per Hectare	Kilograms Active Ingredient per Hectare	
Aphids	0.12 to 0.16	0.06 to 0.08	Use higher rates for greater pest populations and/or dense foliage.
Psyllid (suppression)	0.20	0.10	

Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use a minimum of 94 litres per hectare when applied by ground. Finished spray volumes should be increased when plant foliage is dense.

Apply before populations reach economic thresholds or as populations begin to increase but before damaging populations become established. Scout fields and reapply if necessary.

Allow a minimum of 7 days between applications.

Do not apply more than 3 applications per year.

**For applications to control aphids:** Do not apply more than 0.16 kilograms per hectare Beleaf<sup>®</sup> 50SG Insecticide (0.08 kilograms a.i. per hectare) per application.

**For applications for suppression of psyllids:** Do not apply more than 0.20 kilograms per hectare Beleaf<sup>®</sup> 50SG Insecticide (0.10 kilograms a.i. per hectare) per application.

**Root Vegetables (except sugar beets) (3 day PHI)**

**Crop Subgroup 1B**

**Beet, garden; Burdock, edible; Carrot; Celeriac; Chervil, turnip-rooted; Chicory; Ginseng; Horseradish; Parsley, turnip-rooted; Parsnip (*Pastinaca sativa*); Radish; Radish, oriental; Rutabaga; Salsify; Salsify, black; Salsify, Spanish; Skirret; Turnip.**

PESTS	RATE OF APPLICATION		COMMENTS
	Kilograms Beleaf <sup>®</sup> 50SG Insecticide per Hectare	Kilograms Active Ingredient per Hectare	
Aphids	0.12 to 0.16	0.06 to 0.08	Use higher rates for greater pest populations and/or dense foliage.
Suppression of Tarnished Plant Bug	0.20 (3 applications) or 0.30 (2 applications)	0.10 (3 applications) or 0.15 ( 2 applications)	Apply when Tarnished Plant Bugs first appear in the field and before populations reach high levels. Beleaf will stop Tarnished Plant Bug feeding rapidly but it may take several days to see a reduction in Tarnished Plant Bug numbers. Reapply when new insects are detected.
<p>Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use a minimum of 94 litres per hectare when applied by ground. Finished spray volumes should be increased when plant foliage is dense.</p> <p>Apply before populations reach economic thresholds or as populations begin to increase but before damaging populations become established. Scout fields and reapply if necessary.</p> <p>Allow a minimum of 7 days between applications.</p> <p>Do not apply more than 3 applications per year.</p> <p>Do not apply more than 0.30 kilograms per hectare Beleaf<sup>®</sup> 50SG Insecticide (0.15 kilograms a.i. per hectare) per application. Do not apply more than 0.60 kilograms per hectare Beleaf<sup>®</sup> 50SG Insecticide (0.30 kilograms a.i. per hectare) per year.</p>			



**Pome Fruit (21 day PHI)**

**Crop Group 11-09**

**Apple; azarole; crabapple; mayhaw; medlar; pear; pear, Asian; quince; quince, Chinese; quince, Japanese; tejocote; cultivars, varieties, and/or hybrids of these.**

PESTS	RATE OF APPLICATION		COMMENTS
	Kilograms Beleaf® 50SG Insecticide per Hectare	Kilograms Active Ingredient per Hectare	
Aphids	0.12 to 0.16	0.06 to 0.08	Use higher rates for greater pest populations and/or dense foliage.
Suppression of Tarnished Plant Bug	0.20	0.10	

Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use a minimum of 94 litres per hectare when applied by ground. Finished spray volumes should be increased when plant foliage is dense.

Apply before populations reach economic thresholds or as populations begin to increase but before damaging populations become established. Scout fields and reapply if necessary.

Allow a minimum of 7 days between applications.

Do not apply more than 3 applications per year.

Do not apply more than 0.20 kilograms per hectare Beleaf® 50SG Insecticide (0.10 kilograms a.i. per hectare) per application. Do not apply more than 0.60 kilograms per hectare Beleaf® 50SG Insecticide (0.30 kilograms a.i. per hectare) per year.

**DO NOT** enter or allow worker entry for the purpose of performing thinning by hand in treated areas during the restricted entry interval (REI) of 3 days.

## Stone Fruit (14 day PHI)

### Crop Group 12-09

**Apricot; apricot, Japanese; cherry, black; cherry, Nanking; cherry, sweet; cherry, tart; nectarine; peach; plum; plum, American; plum, beach; plum, Canada; plum, cherry; plum, Chickasaw; plum, Damson; plum, Japanese; plum, Klamath; plum, prune; plumcot; sloe; cultivars, varieties, and/or hybrids of these.**

PESTS	RATE OF APPLICATION		COMMENTS
	Kilograms Beleaf <sup>®</sup> 50SG Insecticide per Hectare	Kilograms Active Ingredient per Hectare	
Aphids	0.12 to 0.20	0.06 to 0.10	Use higher rates for greater pest populations and/or dense foliage.
Suppression of Tarnished Plant Bug	0.20	0.10	

Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use a minimum of 94 litres per hectare when applied by ground. Finished spray volumes should be increased when plant foliage is dense.

Apply before populations reach economic thresholds or as populations begin to increase but before damaging populations become established. Scout fields and reapply if necessary.

Allow a minimum of 7 days between applications.

Do not apply more than 3 applications per year.

Do not apply more than 0.20 kilograms per hectare Beleaf<sup>®</sup> 50SG Insecticide (0.10 kilograms a.i. per hectare) per application. Do not apply more than 0.60 kilograms per hectare Beleaf<sup>®</sup> 50SG Insecticide (0.30 kilograms a.i. per hectare) per year.

**DO NOT** enter or allow worker entry for the purpose of performing thinning by hand in treated areas during the restricted entry interval (REI) of 3 days.

**Flax (7 day PHI)**

PESTS	RATE OF APPLICATION		COMMENTS
	Kilograms Beleaf <sup>®</sup> 50SG Insecticide per Hectare	Kilograms Active Ingredient per Hectare	
Aphids	0.12 to 0.16	0.06 to 0.08	Use higher rates for greater pest populations and/or dense foliage.
<p>Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use a minimum of 94 litres per hectare when applied by ground. Finished spray volumes should be increased when plant foliage is dense.</p> <p>Apply before populations reach economic thresholds or as populations begin to increase but before damaging populations become established. Scout fields and reapply if necessary.</p> <p>Allow a minimum of 7 days between applications.</p> <p>Do not apply more than 3 applications per year.</p> <p>Do not apply more than 0.16 kilograms per hectare Beleaf<sup>®</sup> 50SG Insecticide (0.08 kilograms a.i. per hectare) per application. Do not apply more than 0.48 kilograms per hectare Beleaf<sup>®</sup> 50SG Insecticide (0.24 kilograms a.i. per hectare) per year.</p>			

## Non-grass Animal Feeds (Forage, Fodder, Straw and Hay) (7 day PHI)

### Crop Group 18

**Alfalfa; clover (*Trifolium* spp., *Melilotus* spp.); lespedeza; lupin; sainfoin; trefoil; vetch; vetch, crown; vetch, milk.**

PESTS	RATE OF APPLICATION		COMMENTS
	Kilograms Beleaf <sup>®</sup> 50SG Insecticide per Hectare	Kilograms Active Ingredient per Hectare	
Aphids	0.12 to 0.16	0.06 to 0.08	Use higher rates for greater pest populations and/or dense foliage.
Tarnished Plant Bug	0.20 (3 applications) or 0.30 (2 applications)	0.10 (3 applications) or 0.15 (2 applications)	

Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use a minimum of 94 litres per hectare when applied by ground. Finished spray volumes should be increased when plant foliage is dense.

Apply before populations reach economic thresholds or as populations begin to increase but before damaging populations become established. Scout fields and reapply if necessary.

Allow a minimum of 7 days between applications.

Do not apply more than 3 applications per year.

Do not apply more than 0.30 kilograms per hectare Beleaf<sup>®</sup> 50SG Insecticide (0.15 kilograms a.i. per hectare) per application. Do not apply more than 0.60 kilograms per hectare Beleaf<sup>®</sup> 50SG Insecticide (0.30 kilograms a.i. per hectare) per year.

**NOTE TO USER: READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR THE INDICATED SPECIAL USE APPLICATIONS:**

The DIRECTIONS FOR USE for the uses described in this section of the label were developed by persons other than ISK Biosciences Corporation under the User Requested Minor Use Label Expansion program. For these uses, ISK Biosciences Corporation has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.

**Alfalfa grown for seed (7 day PHI)**

PESTS	RATE OF APPLICATION		COMMENTS
	Kilograms Beleaf® 50SG Insecticide per Hectare	Kilograms Active Ingredient per Hectare	
Aphids	0.12 to 0.16 Maximum	0.06 to 0.08 Maximum	Use higher rate for greater pest populations and/or dense foliage.
Tarnished Plant Bug	0.20 (3 applications) or Maximum 0.30 (2 applications)	0.10 (3 applications) or Maximum 0.15 (2 applications)	

Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use a minimum of 100 litres per hectare when applied by ground. Finished spray volumes should be increased when plant foliage is dense.

Apply before populations reach economic thresholds or as populations begin to increase but before damaging populations become established. Scout fields and reapply if necessary.

Allow a minimum of 7 days between applications.

Do not apply more than 3 applications per year.

Do not apply more than 0.30 kilograms per hectare Beleaf® 50SG Insecticide (0.15 kilograms a.i. per hectare) per application. Do not apply more than 0.60 kilograms per hectare Beleaf® 50SG Insecticide (0.30 kilograms a.i. per hectare) per year.

Do not use seed or treated forage for human or animal consumption.

**Cucumber, Greenhouse (0 day PHI)**

PESTS	RATE OF APPLICATION	COMMENTS
Thrips Aphids Lygus bugs Whiteflies <sup>2</sup>	0.3 g product/L (30 g product/100 L)	Foliar application. Apply sufficient volume to ensure good coverage, up to 1000 L/ha. The maximum volume should be used when plant foliage is dense. Do not apply more than 1 foliar application per crop cycle.
	12.5 mg product/plant	Apply using drip irrigation <sup>1</sup> or drench by hand using sufficient water volume to ensure delivery of the product to the roots.

<sup>1</sup>See Chemigation Section of this label for additional information.

<sup>2</sup> Suppression only for foliar application.

Apply before populations reach economic thresholds or as populations begin to increase but before damaging populations become established.

Allow a minimum of 7 days between applications.

Do not apply more than 2 applications per crop cycle (no more than 1 application per crop cycle may be foliar).

**Peppers, Greenhouse (0 day PHI)**

PESTS	RATE OF APPLICATION	COMMENTS
Thrips Aphids Lygus bugs	0.3 g product/L (30 g product/100 L)	Foliar application. Apply sufficient volume to ensure good coverage, up to 1000 L/ha. The maximum volume should be used when plant foliage is dense. Do not apply more than 1 foliar application per drop cycle.
	30 mg product/m <sup>2</sup>	Apply using drip irrigation <sup>1</sup> or drench by hand using sufficient water volume to ensure delivery of the product to the roots.

<sup>1</sup>See Chemigation Section of this label for additional information.

Apply before populations reach economic thresholds or as populations begin to increase but before damaging populations become established.

Allow a minimum of 7 days between applications.

Do not apply more than 2 applications per crop cycle (no more than 1 application per crop cycle may be foliar).

**Chinese cabbage, bok choy; and cultivars, varieties, and/or hybrids of these,  
Greenhouse (0 day PHI)**

PESTS	RATE OF APPLICATION	COMMENTS
Whiteflies	0.2 g product/L (20 g product/100 L)	Foliar application. Apply sufficient volume to ensure good coverage. The spray volume can vary from 500 to 1000 L/ha, depending on the size of the plants. The maximum volume should be used when plant foliage is dense. Use of this product is restricted to plant growth stages for which thorough coverage can be achieved by spray volumes of 1000 L/ha or less.
<p>Apply before populations reach economic thresholds or as populations begin to increase but before damaging populations become established.            Allow a minimum of 7 days between applications.            Do not apply more than 2 applications per crop cycle.</p>		

**Sweet Potato Slips, Greenhouse (Harvest slips 3 days after last application)**

PESTS	RATE OF APPLICATION	COMMENTS
Thrips Aphids Lygus bugs	0.3 g product/L (30 g product/100 L)	Foliar application. Apply sufficient volume to ensure good coverage, up to 1000 L/ha. The maximum volume should be used when plant foliage is dense. Do not apply more than 1 foliar application per crop cycle.
	30 mg product/m <sup>2</sup>	Apply using drip irrigation or drench by hand using sufficient water volume to ensure delivery of the product to the roots.

<sup>1</sup>See Chemigation Section of this label for additional information.

DO NOT follow greenhouse applications on sweet potato slips with any post-transplant field applications.

Apply before populations reach economic thresholds or as populations begin to increase but before damaging populations become established.

Allow a minimum of 7 days between applications.

Do not apply more than 2 applications per crop cycle (no more than 1 application per crop cycle may be foliar).



### Tomato, Greenhouse (0 day PHI)

PESTS	RATE OF APPLICATION	COMMENTS
Whiteflies	0.2 g product/L (20 g product/100 L)	Foliar application. Apply sufficient volume to ensure good coverage. The spray volume can vary from 500 to 1000 L/ha, depending on the size of the plants. The maximum volume should be used when plant foliage is dense. Use of this product on greenhouse tomatoes is restricted to plant growth stages for which thorough coverage can be achieved by spray volumes of 1000 L/ha or less.
<p>Apply before populations reach economic thresholds or as populations begin to increase but before damaging populations become established.            Allow a minimum of 7 days between applications.            Do not apply more than 2 applications per crop cycle.</p>		

### Lettuce, Greenhouse (0 day PHI)

PESTS	RATE OF APPLICATION	COMMENTS
Aphids	0.14 - 0.20 g product/L (14 - 20 g product/100L)	Foliar application. Apply sufficient volume to ensure good coverage, up to 1000 L/ha.
<p>Apply before populations reach economic thresholds or as populations begin to increase but before damaging populations become established. Use the lower rate for building populations and the higher rate for established populations and or dense foliage.</p> <p>Do not apply more than 1 application per crop cycle.</p>		

## Herbs (Crop Group 25) (7 day PHI)

**Crop Group 25 – Herbs: Agrimony (fresh leaves) Agrimony (dried leaves); Angelica (fresh leaves); Angelica (dried leaves); Angelica, fragrant (fresh leaves); Angelica, fragrant (dried leaves); Applemint (fresh leaves); Applemint (dried leaves); Avarum (fresh leaves); Avarum (dried leaves); Balm (fresh leaves); Balm (dried leaves); Balloon pea (fresh leaves); Balloon pea (dried leaves); Barrenwort (dried leaves); Basil (fresh leaves); Basil (dried leaves); Basil, American (fresh leaves); Basil, American (dried leaves); Basil, Greek (fresh leaves); Basil, Greek (dried leaves); Basil, holy (fresh leaves); Basil, holy (dried leaves); Basil, lemon (fresh leaves); Basil, lemon (dried leaves); Basil, Russian (fresh leaves); Basil, Russian (dried leaves); Bay (fresh leaves); Bay (dried leaves); Bisongrass (fresh leaves); Bisongrass (dried leaves); Blue mallow (fresh leaves); Boneset (fresh leaves); Boneset (dried leaves); Borage (fresh leaves); Borage (dried leaves); Borage, Indian (fresh leaves); Borage, Indian (dried leaves); Burnet (fresh leaves); Burnet (dried leaves); Burnet, garden (fresh leaves); Burnet, garden (dried leaves); Burnet, salad (fresh leaves); Burnet, salad (dried leaves); Butterbur (dried leaves); Calamint (fresh leaves); Calamint (dried leaves); Calamint, large-flower (fresh leaves); Calamint, large-flower (dried leaves); Calamint, lesser (fresh leaves); Calamint, lesser (dried leaves); Calendula (fresh leaves); Calendula (dried leaves); Caltrop (fresh leaves); Caltrop (dried leaves); Camomile, Chamomile (fresh leaves); Camomile, Chamomile (dried leaves); Camomile, Chamomile (German, fresh leaves); Camomile, Chamomile (German, dried leaves); Camomile, Chamomile (Roman, fresh leaves); Camomile, Chamomile (Roman, dried leaves); Caraway (fresh leaves); Caraway (dried leaves); Cat's claw (dried leaves); Catnip (fresh leaves); Catnip (dried leaves); Catnip, Japanese (fresh leaves); Catnip, Japanese (dried leaves); Celandine, greater (fresh leaves); Celandine, lesser (fresh leaves); Centaury (fresh leaves); Centaury (dried leaves); Chaste tree (fresh leaves); Chaste tree (dried leaves); Chervil (dried leaves); Chinese chastetree (dried leaves); Chinese foxglove (dried leaves); Chive (dried leaves); Chive, Chinese (dried leaves); Cicely, sweet (fresh leaves); Cicely, sweet (dried leaves); Cilantro (dried leaves); Clary (fresh leaves); Clary (dried leaves); Coriander, Bolivian (fresh leaves); Coriander, Bolivian (dried leaves); Coriander, Vietnamese (fresh leaves); Coriander, Vietnamese (dried leaves); Costmary (fresh leaves); Costmary (dried leaves); Creat (dried leaves); Culantro (fresh leaves); Culantro (dried leaves); Curry leaf (fresh leaves); Curry leaf (dried leaves); Curryplan (fresh leaves); Cut leaf (fresh leaves); Cut leaf (fresh leaves); Dillweed (dried leaves); Dokudami (fresh leaves); Echinacea (dried leaves); Epazote (fresh leaves); Epazote (dried leaves); Eucommia (dried leaves); Evening primrose (fresh leaves); Evening primrose (dried leaves); Fennel, common (fresh leaves); Fennel, common (dried leaves); Fennel, Florence (dried leaves); Fennel, Spanish (fresh leaves); Fennel, Spanish (dried leaves); Fenugreek (fresh leaves); Fenugreek (dried leaves); Feverfew (fresh leaves); Feverfew (dried leaves); Field pennycress (fresh leaves); Flowers, edible, fresh, multiple species; Flowers, edible, dried, multiple species; Fumitory (fresh leaves); Fumitory (dried leaves); Galbanum (dried leaves); Gambir (fresh leaves); Geranium (fresh leaves); Geranium (dried leaves); Geranium, lemon (fresh leaves); Geranium, lemon (dried leaves); Geranium, rose (fresh leaves); Geranium, rose (dried leaves); Germander, golden (fresh leaves); Germander, golden (dried leaves); Gotu kola (dried leaves); Gumweed (fresh leaves); Gumweed (dried leaves); Gymnema (dried leaves); Gypsywort (fresh leaves); Gypsywort (dried leaves); Heal-all (fresh leaves); Heal-all (dried leaves); Honewort (fresh leaves); Honeybush (dried leaves); Horehound (fresh leaves); Horehound (dried leaves); Horsemint (fresh leaves); Horsemint (dried leaves); Hyssop (fresh leaves); Hyssop (dried leaves); Hyssop, anise (fresh leaves); Hyssop, anise (dried**

leaves); Jasmine (fresh leaves); Jasmine (dried leaves); Labrador tea (fresh leaves); Labrador tea (dried leaves); Lavender (fresh leaves); Lavender (dried leaves); Lemongrass (fresh leaves); Lemongrass (dried leaves); Lemon verbena (fresh leaves); Lemon verbena (dried leaves); Lovage (fresh leaves); Lovage (dried leaves); Love-in-a-mist (fresh leaves); Love-in-a-mist (dried leaves); Mamaki (dried leaves); Marigold (fresh leaves); Marigold (dried leaves); Marigold, African (fresh leaves); Marigold, African (dried leaves); Marigold, Aztec (fresh leaves); Marigold, Aztec (dried leaves); Marigold, French (fresh leaves); Marigold, French (dried leaves); Marigold, Irish lace (fresh leaves); Marigold, Irish lace (dried leaves); Marigold, licorice (fresh leaves); Marigold, licorice (dried leaves); Marigold, Mexican mint (fresh leaves); Marigold, Mexican mint (dried leaves); Marigold, signet (fresh leaves); Marigold, signet (dried leaves); Marjoram, (fresh leaves); Marjoram (dried leaves); Marjoram, pot (fresh leaves); Marjoram, pot (dried leaves); Marjoram, sweet (fresh leaves); Marjoram, sweet (dried leaves); Marshmallow (fresh leaves); Marshmallow (dried leaves); Meadowsweet (fresh leaves); Meadowsweet (dried leaves); Mint (fresh leaves); Mint (dried leaves); Mint corn, (fresh leaves); Mint corn (dried leaves); Mint, Korean (fresh leaves); Mint, Korean (dried leaves); Monarda (fresh leaves); Monarda (dried leaves); Motherwort (fresh leaves); Motherwort (dried leaves); Mountainmint (fresh leaves); Mountainmint (dried leaves); Mountainmint, clustered (fresh leaves); Mountainmint, clustered (dried leaves); Mountainmint, hoary (fresh leaves); Mountainmint, hoary (dried leaves); Mountainmint, Virginia (fresh leaves); Mountainmint, Virginia (dried leaves); Mountainmint, whorled (fresh leaves); Mountainmint, whorled (dried leaves); Mugwort (fresh leaves); Mugwort (dried leaves); Mulberry, white (dried leaves); Mullein (fresh leaves); Mullein (dried leaves); Nasturtium (fresh leaves); Nasturtium (dried leaves); Nasturtium, bush (fresh leaves); Nasturtium, bush (dried leaves); Nasturtium, garden (fresh leaves); Nasturtium, garden (dried leaves); Nettle (fresh leaves); Nettle (dried leaves); Oregano (fresh leaves); Oregano (dried leaves); Oregano, Mexican (fresh leaves); Oregano, Mexican (dried leaves); Oregano, Puerto Rico (fresh leaves); Oregano, Puerto Rico (dried leaves); Oswego tea (fresh leaves); Oswego tea (dried leaves); Pandan leaf (fresh leaves); Pandan leaf (dried leaves); Pansy (fresh leaves); Pansy (dried leaves); Paracress (fresh leaves); Paracress (dried leaves); Parsley (dried leaves); Pennyroyal (fresh leaves); Pennyroyal (dried leaves); Peppermint (fresh leaves); Peppermint (dried leaves); Perilla (fresh leaves); Perilla (dried leaves); Rooibos (dried leaves); Rose (fresh leaves); Rose (dried leaves); Rosemary (fresh leaves); Rosemary (dried leaves); Sage (fresh leaves); Sage (dried leaves); Sage, Greek (fresh leaves); Sage, Greek (dried leaves); Sage, Spanish (fresh leaves); Sage, Spanish (dried leaves); Savory, summer (fresh leaves); Savory, summer (dried leaves); Savory, winter (fresh leaves); Savory, winter (dried leaves); Sorrel (fresh leaves); Sorrel, (dried leaves); Sorrel, French (fresh leaves); Sorrel, French (dried leaves); Sorrel, garden (fresh leaves); Sorrel, garden (dried leaves); Southernwood (fresh leaves); Southernwood (dried leaves); Spearmint (fresh leaves); Spearmint (dried leaves); Spearmint, Scotch (fresh leaves); Spearmint, Scotch (dried leaves); Spotted beebalm (fresh leaves); Spotted beebalm (dried leaves); Squaw vine (dried leaves); St. John's Wort (dried leaves); Stevia (dried leaves); Swamp leaf (fresh leaves); Tansy (fresh leaves); Tansy (dried leaves); Tarragon (fresh leaves); Tarragon (dried leaves); Thyme (fresh leaves); Thyme (dried leaves); Thyme, creeping (fresh leaves); Thyme, creeping (dried leaves); Thyme, lemon (fresh leaves); Thyme, lemon (dried leaves); Thyme, mastic (fresh leaves); Thyme, mastic (dried leaves); Toon, Chinese (fresh leaves); Toon, Chinese (dried leaves); Vasaka (dried leaves); Veronica (fresh leaves); Violet (fresh leaves); Violet (dried leaves); Watermint (fresh leaves); Watermint (dried leaves); Waterpepper (fresh leaves); Wild bergamot (fresh leaves); Wild bergamot (dried leaves); Wintergreen (fresh leaves); Wintergreen (dried leaves); Wood betony (dried leaves);

**Woodruff (fresh leaves); Woodruff (dried leaves); Wormwood (fresh leaves); Wormwood (dried leaves); Wormwood, Roman (fresh leaves); Wormwood, Roman (dried leaves); Yarrow (fresh leaves); Yarrow (dried leaves); Yellow gentian (fresh leaves); Yellow gentian (dried leaves); Yerba santa (fresh leaves); Yerba santa (dried leaves); Yomogi (fresh leaves); Yomogi (dried leaves); Also included are cultivars, varieties, and hybrids of these commodities.**

PESTS	RATE OF APPLICATION		COMMENTS
	Kilograms Beleaf <sup>®</sup> 50SG Insecticide per Hectare	Kilograms Active Ingredient per Hectare	
Aphids	0.12 to 0.16	0.06 to 0.08	
<p>Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use a minimum of 94 litres per hectare when applied by ground. Finished spray volumes should be increased when plant foliage is dense.</p> <p>Apply before populations reach economic thresholds or as populations begin to increase but before damaging populations become established. Scout fields and reapply if necessary.</p> <p>Allow a minimum of 14 days between applications.</p> <p>Do not apply more than 3 applications per year.</p> <p>Do not apply more than 0.16 kilograms per hectare Beleaf<sup>®</sup> 50SG Insecticide (0.08 kilograms a.i. per hectare) per application. Do not apply more than 0.48 kilograms per hectare Beleaf<sup>®</sup> 50SG Insecticide (0.24 kilograms a.i. per hectare) per year.</p>			

### **Greenhouse Grown Ornamentals including Cut Flowers**

APPLICATION METHOD	PESTS	RATE OF APPLICATION	COMMENTS
Foliar Spray	Thrips, Aphids	0.3 g product/L (30 g product/100 L)	Apply before populations reach economic thresholds or as populations begin to increase but before damaging populations become established. Apply sufficient volume to ensure good coverage, up to 1000 L/ha. The maximum volume should be used when plant foliage is dense. Do not apply more than 2 foliar applications per crop cycle.

	Whitefly	0.2 g product/L (20 g product/100 L)	Apply before populations reach economic thresholds or as populations begin to increase but before damaging populations become established. Apply sufficient volume to ensure good coverage, up to 1000 L/ha. The maximum volume should be used when plant foliage is dense. Do not apply more than 2 foliar applications per crop cycle.
	Mealybugs	0.16 to 0.3 g product/L (16 to 30 g/100L)	Treat when insects appear. Apply sufficient volume to ensure good coverage, up to 1000 L/ha. The maximum volume should be used when plant foliage is dense. Do not apply more than 2 applications per crop cycle.
Drench (Herbaceous Ornamentals Only, Including Cut Flowers)	Thrips, Aphids, Whiteflies, Lygus bugs	300 g product/ha	Apply before populations reach economic thresholds or as populations begin to increase but before damaging populations become established. Apply using drip irrigation <sup>1</sup> or drench by hand using sufficient water volume to ensure delivery of the product to the roots. Do not apply more than 2 applications per crop cycle.

<sup>1</sup>See Chemigation Section of this label for additional information.

Apply before populations reach economic thresholds or as populations begin to increase but before damaging populations become established.

**DO NOT** apply using handheld mistblower/airblast or handheld fogging equipment.

Do not apply more than 600 grams per hectare Beleaf® 50SG Insecticide (300 grams a.i. per hectare) using foliar and/or drench applications per crop cycle.

Allow a minimum of 7 days between applications.

\*\*\* Ornamental varieties may vary in their tolerance to pest control products, including Beleaf® 50SG Insecticide. Since not all ornamental varieties have been tested for tolerance to Beleaf® 50SG Insecticide, first use of Beleaf® 50SG Insecticide should be limited to a small area of each variety to confirm tolerance prior to adoption as a general field practice. Additionally, consult your seed or plant supplier for information on the tolerance of specific varieties of herbaceous ornamentals to Beleaf® 50SG Insecticide. Certain pansy cultivars have exhibited sensitivity to flonicamid. Prior to use on pansy, apply flonicamid at the appropriate use rate to a small portion of the pansy crop (about 10 units per cultivar) and visually assess impact 7 to 10 days after application.

## Ornamentals, Outdoor (excluding conifers)

PESTS	RATE OF APPLICATION		COMMENTS
Aphids	0.12 to 0.16 kg product/ha	0.06 to 0.08 kg active ingredient/ha	Apply before populations reach economic thresholds or as populations begin to increase but before damaging populations become established. Use higher rates for greater pest populations and/or dense foliage. Do not apply more than 3 applications per year.
Chilli Thrips ( <i>Scirtothrips dorsalis</i> Hood)	0.3 g product/L (30 g product/100 L)		Foliar application. Treat when insects appear. Do not apply more than 2 applications per year.
Mealybugs	0.16 to 0.3 g product/L (16 to 30 g product/100 L)		

Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use 94 to 1000 litres per hectare applied by ground equipment. Finished spray volumes should be increased when plant foliage is dense.

Allow a minimum of 7 days between applications.

Do not apply more than 600 grams per hectare Beleaf® 50SG Insecticide (300 grams a.i. per hectare) per year.

Do not apply more than 3 applications per year.

Do not apply more than 0.16 kilograms per hectare Beleaf® 50SG Insecticide (0.08 kilograms a.i. per hectare) per application. Do not apply more than 0.48 kilograms per hectare Beleaf® 50SG Insecticide (0.24 kilograms a.i. per hectare) per season.

Do not enter or allow worker entry into treated areas for the purposes of performing the re-entry activities of cut flowers hand harvesting, disbudding, or hand pruning during the restricted entry interval of 4 days.

**DO NOT** enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours to perform hand-set/hand-line irrigation related activities involving foliar contact.

Ornamental varieties may vary in their tolerance to pest control products, including Beleaf® 50SG Insecticide. Since not all ornamental varieties have been tested for tolerance to Beleaf® 50SG Insecticide, first use of Beleaf® 50SG Insecticide should be limited to a small area of each variety to confirm tolerance prior to adoption as a general field practice. Additionally, consult your seed or plant supplier for information on the tolerance of specific varieties of ornamentals to Beleaf® 50SG Insecticide.

Certain pansy cultivars have exhibited sensitivity to flonicamid. Prior to use on pansy, apply flonicamid at the appropriate use rate to a small portion of the pansy crop (about 10 units per cultivar) and visually assess impact 7 to 10 days after application.

**Low Growing Berry (0 day PHI)**

**Crop Subgroup 13-07G**

**Bearberry; Bilberry; Blueberry, lowbush; Cloudberry; Cranberry; Lingonberry; Muntries; Partridgeberry; Strawberry; cultivars, varieties and/or hybrids of these.**

PESTS	RATE OF APPLICATION		COMMENTS
	Kilograms Beleaf <sup>®</sup> 50SG Insecticide per Hectare	Kilograms Active Ingredient per Hectare	
Aphids	0.12 to 0.16	0.06 to 0.08	Apply before aphid populations reach economic thresholds or as populations begin to increase but before damaging populations become established. Scout fields and reapply if necessary. Use higher rates for greater pest populations and/or dense foliage.
Suppression of lygus bugs including tarnished plant bug	0.20	0.10	Apply when lygus bugs first appear in the field and before populations reach high levels. Beleaf <sup>®</sup> 50SG Insecticide will stop lygus bug feeding rapidly but it may take several days to see a reduction in lygus bug numbers. Reapply when new insects are detected.
Thrips	0.20 (3 applications) or 0.30 (2 applications)	0.10 (3 applications) or 0.15 (2 applications)	Apply when thrips first appear in the field and before populations reach high levels. Beleaf will stop thrips feeding rapidly but it may take several days to see a reduction in thrips numbers. Under heavy infestation, use the higher rate of 0.30 kg/ha. Multiple applications may be required.
<p>Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use a minimum of 94 litres per hectare when applied by ground. Finished spray volumes should be increased when plant foliage is dense.</p> <p>Allow a minimum of 7 days between applications.</p> <p>Do not apply more than 3 applications per year.</p>			

**For applications to control aphids:** Do not apply more than 0.16 kilograms per hectare Beleaf® 50SG Insecticide (0.08 kilograms a.i. per hectare) per application ( up to 3 applications per year).

**For applications to suppress lygus bugs:** Do not apply more than 0.20 kilograms per hectare Beleaf® 50SG Insecticide (0.10 kilograms a.i. per hectare) per application (up to 3 applications per year).

**For applications to control thrips:** Do not apply more than 0.30 kilograms per hectare Beleaf® 50SG Insecticide (0.15 kilograms a.i. per hectare) per application. Do not apply more than 0.60 kilograms per hectare Beleaf® 50SG Insecticide (0.30 kilograms a.i. per hectare) per year.

## Bushberries (0 day PHI)

### (Crop Subgroup 13-07B)

**Aronia berry; Blueberry, highbush; Blueberry, lowbush; Buffalo currant; Chilean guava; Currant, black; Currant, red; Elderberry; European barberry; Gooseberry; Highbush cranberry; Honeysuckle, edible; Huckleberry; Jostaberry; Juneberry (Saskatoon berry); Lingonberry; Native currant; Salal; Sea buckthorn; Cultivars, varieties and/or hybrids of these.**

PESTS	RATE OF APPLICATION		COMMENTS
	Kilograms Beleaf® 50SG Insecticide per Hectare	Kilograms Active Ingredient per Hectare	
Aphids	0.12 to 0.16	0.06 to 0.08	Use higher rates for greater pest populations and/or dense foliage.
<p>Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use a minimum of 94 litres per hectare when applied by ground. Finished spray volumes should be increased when plant foliage is dense.</p> <p>Apply before populations reach economic thresholds or as populations begin to increase but before damaging populations become established. Scout fields and reapply if necessary.</p> <p>Allow a minimum of 7 days between applications. Do not apply more than 3 applications per year.</p> <p>Do not apply more than 0.16 kilograms per hectare Beleaf® 50SG Insecticide (0.08 kilograms a.i. per hectare) per application. Do not apply more than 0.48 kilograms per hectare Beleaf® 50SG Insecticide (0.24 kilograms a.i. per hectare) per year.</p>			



**Strawberry, Greenhouse (0 day PHI)**

PESTS	RATE OF APPLICATION	COMMENTS
Aphids  Lygus bugs including tarnished plant bug ( <i>Lygus lineolaris</i> )	0.3 g product/L	Beleaf <sup>®</sup> 50SG Insecticide will stop Aphid and Lygus bug feeding rapidly but it may take several days to see a reduction in pest numbers.
<p>Thorough spray coverage of plant foliage is essential for optimum control. Use a maximum spray volume of 650 litres.</p> <p>Apply when Aphids and/or Lygus bugs first appear in the greenhouse and before populations reach high levels.</p> <p>Do not apply more than 1 application per crop cycle.</p> <p>Use Restrictions: DO NOT apply using fogging or ultra low volume equipment. DO NOT apply using handheld mistblowers or handheld airblast equipment.</p>		

**Legume Vegetables – Succulent or Dried, except soybeans (7 day PHI)**

**Crop Group 6**

**Bean (*Lipinus* spp.) (includes grain lupin, sweet lupin, white lupin and white sweet lupin), Bean (*Phaseolus* spp.) (includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean and wax bean); bean (*Vigna* spp.) (includes adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, Crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean and yardlong bean), Broadbean, (fava bean) (*Vicia faba*), Chickpea (garbanzo bean) (*Cicer arietinum*), Guar (*Cyamopsis tetragonoloba*), Jackbean (*Canavalia ensiformis*), Lablab bean (hyacinth bean) (*Lablab purpureus*), Lentil (*Lens esculenta*), Pea (*Pisum* spp.) (includes dwarf pea, edible-pod pea, English pea, field pea, garden pea, green pea, snow pea, sugar snap pea), Pigeon pea (*Cajanus cajan*), Sword bean (*Canavalia gladiata*)**

PESTS	RATE OF APPLICATION		COMMENTS
	Kilograms Beleaf <sup>®</sup> 50SG	Kilograms Active Ingredient per Hectare	

	Insecticide per Hectare		
Aphids	0.12 to 0.16	0.06 to 0.08	Apply before aphid populations reach economic thresholds or as populations begin to increase but before damaging populations become established. Scout fields and reapply if necessary. Use higher rates for greater pest populations and/or dense foliage.
Reduces the numbers of lygus bugs including tarnished plant bug	0.20	0.10	Apply when lygus bugs first appear in the field and before populations reach high levels. Beleaf <sup>®</sup> 50SG Insecticide will stop lygus bug feeding rapidly but it may take several days to see a reduction in lygus bug numbers. Reapply when new insects are detected.
<p>Thorough spray coverage of plant foliage is essential for optimum performance. Apply in sufficient water to ensure good coverage; use a minimum of 94 litres per hectare when applied by ground. Finished spray volumes should be increased when plant foliage is dense.</p> <p>Allow a minimum of 7 days between applications.</p> <p>Do not apply more than 3 applications per year.</p> <p><b>For applications to control aphids:</b> Do not apply more than 0.16 kilograms per hectare Beleaf<sup>®</sup> 50SG Insecticide (0.08 kilograms a.i. per hectare) per application (up to 3 applications per year).</p> <p><b>For applications to reduce the numbers of lygus bugs:</b> Do not apply more than 0.20 kilograms per hectare Beleaf<sup>®</sup> 50SG Insecticide (0.10 kilograms a.i. per hectare) per application (up to 3 applications per year).</p>			

## Balsam Fir and Fraser Fir, including Christmas Trees

PESTS	RATE OF APPLICATION		COMMENTS
	Kilograms Beleaf <sup>®</sup> 50SG Insecticide per Hectare	Kilograms Active Ingredient per Hectare	
Balsam twig aphids ( <i>Mindarus abietinus</i> )	0.12 to 0.16	0.06 to 0.08	Use higher rates for greater pest populations and/or dense foliage.
<p>Apply after the stem mothers have hatched but before the sexuparae (second generation) aphids are present, based on a degree-days model and when insect populations reach intervention threshold.</p> <p>Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use a minimum of 300 litres per hectare when applied by ground. Finished spray volumes should be increased when plant foliage is dense.</p> <p>Allow a minimum of 7 days between applications.</p> <p>Do not apply more than 2 applications per year.</p> <p>Do not apply more than 0.16 kilograms per hectare Beleaf<sup>®</sup> 50SG Insecticide (0.08 kilograms a.i. per hectare) per application. Do not apply more than 0.32 kilograms per hectare Beleaf<sup>®</sup> 50SG Insecticide (0.16 kilograms a.i. per hectare) per year.</p>			

**Sweet Corn (0 day PHI for mechanical harvest, 2 day PHI for hand harvest)**

PESTS	RATE OF APPLICATION		COMMENTS
	Kilograms Beleaf® 50SG Insecticide per Hectare	Kilograms Active Ingredient per Hectare	
Aphids	0.12 to 0.16	0.06 to 0.08	

Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water to ensure good coverage; use a minimum of 94 litres per hectare when applied by ground. Finished spray volumes should be increased when plant foliage is dense.

Apply before populations reach economic thresholds or as populations begin to increase but before damaging populations become established. Scout fields and reapply if necessary.

Allow a minimum of 7 days between applications.

Do not apply more than 3 applications per year.

Do not apply more than 0.16 kilograms per hectare Beleaf® 50SG Insecticide (0.08 kilograms a.i. per hectare) per application. Do not apply more than 0.48 kilograms per hectare Beleaf® 50SG Insecticide (0.24 kilograms a.i. per hectare) per season.

**DO NOT** enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 2 days to perform hand harvesting in corn.

**NOTICE TO USER**

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

Beleaf® is a registered trademark of Ishihara Sangyo Kaisha, Ltd.